

Daftar Pustaka

- Ayala-Ramírez, P., Serrano, N., Barrera, V., Bejarano, J. P., Silva, J. L., Martínez, R., Gil, F., Olaya-C, M., & García-Robles, R. (2020). Risk factors and fetal outcomes for preeclampsia in a Colombian cohort. *Heliyon*, *6*(9), e05079. <https://doi.org/10.1016/j.heliyon.2020.e05079>
- Bdolah, Y., Elchalal, U., Natanson-Yaron, S., Yechiam, H., Bdolah-Abram, T., Greenfield, C., Goldman-Wohl, D., Milwidsky, A., Rana, S., Karumanchi, S. A., Yagel, S., & Hochner-Celnikier, D. (2014). Relationship between nulliparity and preeclampsia may be explained by altered circulating soluble fms-like tyrosine kinase 1. *Hypertension in Pregnancy*, *33*(2), 250–259. <https://doi.org/10.3109/10641955.2013.858745>
- Berghella, V. (n.d.). *Maternal-Fetal Evidence Based Guidelines; Fourth Edition*. <https://www.crcpress.com/Series-in-Maternal-Fetal-Medicine/bookseries/>
- Bernardes, T. P., Mol, B. W., Ravelli, A. C. J., van den Berg, P., Marike Boezen, H., & Groen, H. (2020). Early and late onset pre-eclampsia and small for gestational age risk in subsequent pregnancies. *PLoS ONE*, *15*(3). <https://doi.org/10.1371/journal.pone.0230483>
- Catalano, P. M. (2010). Obesity, insulin resistance, and pregnancy outcome. In *Reproduction* (Vol. 140, Issue 3, pp. 365–371). <https://doi.org/10.1530/REP-10-0088>
- Cunningham, F. G., Leveno, K. J., Dashe, J. S., Hoffman, B. L., Spong, C. Y., & Casey, B. M. (2022). *Williams Obstetrics 26th ed.*
- de Logivière, V., Tsatsaris, V., Lepercq, J., Goffinet, F., & Girault, A. (2025). Evaluating the proteinuria/creatininuria ratio as a rapid prognostic tool for complications of preeclampsia: A comparison with 24-hour proteinuria. *Journal of Gynecology Obstetrics and Human Reproduction*, *54*(2). <https://doi.org/10.1016/j.jogoh.2024.102873>
- Delcroix-Gomez, C., Delcroix, M. H., Jamee, A., Gauthier, T., Marquet, P., & Aubard, Y. (2022). Fetal growth restriction, low birth weight, and preterm birth: Effects of active or passive smoking evaluated by maternal expired CO at delivery, impacts of cessation at different trimesters. *Tobacco Induced Diseases*, *20*(August). <https://doi.org/10.18332/tid/152111>
- Dianing Tyas, B., Lestari, P., Ilham, M., & Akbar, A. (2019). *Maternal Perinatal Outcomes Related to Advanced Maternal Age in Preeclampsia Pregnant Women* (Vol. 13, Issue 4). <http://jfrh.tums.ac.ir>
- Dong, X., Gou, W., Li, C., Wu, M., Han, Z., Li, X., & Chen, Q. (2017). Proteinuria in preeclampsia: Not essential to diagnosis but related to disease severity and fetal outcomes. *Pregnancy Hypertension*, *8*, 60–64. <https://doi.org/10.1016/j.preghy.2017.03.005>
- Duckitt, K., & Harrington, D. (2005). Risk factors for pre-eclampsia at antenatal booking: Systematic review of controlled studies. In *British Medical Journal* (Vol. 330, Issue 7491, pp. 565–567). <https://doi.org/10.1136/bmj.38380.674340.E0>

- Gathiram, P., & Moodley, J. (2016). Pre-eclampsia: Its pathogenesis and pathophysiology. In *Cardiovascular Journal of Africa* (Vol. 27, Issue 2, pp. 71–78). Clinics Cardive Publishing (PTY)Ltd. <https://doi.org/10.5830/CVJA-2016-009>
- Genc, S., Emeklioglu, C. N., Cingillioglu, B., Akturk, E., Turhan Ozkan, H., & Mihmanli, V. (2021). The effect of parity on obstetric and perinatal outcomes in pregnancies at the age of 40 and above: A retrospective study. *Croatian Medical Journal*, 62(2), 130–136. <https://doi.org/10.3325/cmj.2021.62.130>
- Genest, D. S., Dal Soglio, D., Girard, S., & Rey, E. (2021). Association between proteinuria and placental pathology in preeclampsia: A retrospective study. *SAGE Open Medicine*, 9. <https://doi.org/10.1177/20503121211058053>
- Gottardi, E., Lecarpentier, E., Villette, C., Berman, A., Redel, D., Tsatsaris, V., Goffinet, F., & Haddad, B. (2021). Preeclampsia before 26 weeks of gestation: Obstetrical prognosis for the subsequent pregnancy. *Journal of Gynecology Obstetrics and Human Reproduction*, 50(3), 102000. <https://doi.org/10.1016/j.jogoh.2020.102000>
- He, X. J., Dai, R. xue, & Hu, C. L. (2020). Maternal prepregnancy overweight and obesity and the risk of preeclampsia: A meta-analysis of cohort studies. In *Obesity Research and Clinical Practice* (Vol. 14, Issue 1, pp. 27–33). Elsevier Ltd. <https://doi.org/10.1016/j.orcp.2020.01.004>
- Hokken-Koelega, A. C. S., van der Steen, M., Boguszewski, M. C. S., Cianfarani, S., Dahlgren, J., Horikawa, R., Mericq, V., Rapaport, R., Alherbish, A., Braslavsky, D., Charmandari, E., Chernausek, S. D., Cutfield, W. S., Dauber, A., Deeb, A., Goedegebuure, W. J., Hofman, P. L., Isganatis, E., Jorge, A. A., ... Yau, M. (2023). International Consensus Guideline on Small for Gestational Age: Etiology and Management From Infancy to Early Adulthood. In *Endocrine Reviews* (Vol. 44, Issue 3, pp. 539–565). Endocrine Society. <https://doi.org/10.1210/endrev/bnad002>
- Komoróczy, B., Vánca, S., Váradi, A., Hegyi, P., Vágási, V., Baradács, I., Szabó, A., Nyirády, P., Benkő, Z., & Ács, N. (2025). Optimal Aspirin Dosage for the Prevention of Preeclampsia and Other Adverse Pregnancy Outcomes: A Systematic Review and Meta-Analysis of Randomized Controlled Trials. In *Journal of Clinical Medicine* (Vol. 14, Issue 7). Multidisciplinary Digital Publishing Institute (MDPI). <https://doi.org/10.3390/jcm14072134>
- Kovo, M., Schreiber, L., Ben-Haroush, A., Gold, E., Golan, A., & Bar, J. (2012). The placental component in early-onset and late-onset preeclampsia in relation to fetal growth restriction. *Prenatal Diagnosis*, 32(7), 632–637. <https://doi.org/10.1002/pd.3872>
- Kuitunen, I., Huttunen, T. T., Ponkilainen, V. T., & Kekki, M. (2022). Incidence of obese parturients and the outcomes of their pregnancies: A nationwide register study in Finland. *European Journal of Obstetrics and Gynecology and Reproductive Biology*, 274, 62–67. <https://doi.org/10.1016/j.ejogrb.2022.05.006>

- Lees, C. C., Stampalija, T., Baschat, A., da Silva Costa, F., Ferrazzi, E., Figueras, F., Hecher, K., Poon, L. C., Salomon, L. J., & Unterscheider, J. (2020). ISUOG Practice Guidelines: diagnosis and management of small-for-gestational-age fetus and fetal growth restriction. In *Ultrasound in Obstetrics and Gynecology* (Vol. 56, Issue 2, pp. 298–312). John Wiley and Sons Ltd. <https://doi.org/10.1002/uog.22134>
- Lewandowska, M., Więckowska, B., Sajdak, S., & Lubiński, J. (2020). Pre-pregnancy obesity vs. Other risk factors in probability models of preeclampsia and gestational hypertension. *Nutrients*, *12*(9), 1–19. <https://doi.org/10.3390/nu12092681>
- Li, X., Zhang, W., Lin, J., Liu, H., Yang, Z., Teng, Y., Duan, S., Li, Y., Xie, Y., Lin, X., Xie, L., Peng, Q., Huang, J., Chen, J., Duan, W., Luo, J., & Zhang, J. (2018). Preterm birth, low birthweight, and small for gestational age among women with preeclampsia: Does maternal age matter? *Pregnancy Hypertension*, *13*, 260–266. <https://doi.org/10.1016/j.preghy.2018.07.004>
- Magee, L. A., Brown, M. A., Hall, D. R., Gupte, S., Hennessy, A., Karumanchi, S. A., Kenny, L. C., McCarthy, F., Myers, J., Poon, L. C., Rana, S., Saito, S., Staff, A. C., Tsigas, E., & von Dadelszen, P. (2022). The 2021 International Society for the Study of Hypertension in Pregnancy classification, diagnosis & management recommendations for international practice. *Pregnancy Hypertension*, *27*, 148–169. <https://doi.org/10.1016/j.preghy.2021.09.008>
- Malik, A., Jee, B., & Gupta, S. K. (2019). Preeclampsia: Disease biology and burden, its management strategies with reference to India. In *Pregnancy Hypertension* (Vol. 15, pp. 23–31). Elsevier B.V. <https://doi.org/10.1016/j.preghy.2018.10.011>
- Melamed, N., Baschat, A., Yinon, Y., Athanasiadis, A., Mecacci, F., Figueras, F., Berghella, V., Nazareth, A., Tahlak, M., McIntyre, H. D., Da Silva Costa, F., Kihara, A. B., Hadar, E., McAuliffe, F., Hanson, M., Ma, R. C., Gooden, R., Sheiner, E., Kapur, A., ... Hod, M. (2021). FIGO (international Federation of Gynecology and obstetrics) initiative on fetal growth: best practice advice for screening, diagnosis, and management of fetal growth restriction. *International Journal of Gynecology and Obstetrics*, *152*(S1), 3–57. <https://doi.org/10.1002/ijgo.13522>
- Mitani, M., Matsuda, Y., Makino, Y., Akizawa, Y., & Ohta, H. (2009). Clinical features of fetal growth restriction complicated later by preeclampsia. *Journal of Obstetrics and Gynaecology Research*, *35*(5), 882–887. <https://doi.org/10.1111/j.1447-0756.2009.01120.x>
- Morikawa, M., Mayama, M., Saito, Y., Nakagawa-Akabane, K., Umazume, T., Chiba, K., Kawaguchi, S., Cho, K., & Watari, H. (2020). Severe proteinuria as a parameter of worse perinatal/neonatal outcomes in women with preeclampsia. *Pregnancy Hypertension*, *19*, 119–126. <https://doi.org/10.1016/j.preghy.2019.12.013>

- National Institute for Health and Care Excellence. (2024). *Hypertension in pregnancy: diagnosis and management NICE guideline*.
www.nice.org.uk/guidance/ng133
- Odibo, A. O., Nelson, D., Stamilio, D. M., Sehdev, H. M., & Macones, G. A. (2006). Advanced maternal age is an independent risk factor for intrauterine growth restriction. *American Journal of Perinatology*, 23(5), 325–328.
<https://doi.org/10.1055/s-2006-947164>
- Ogawa, K., Urayama, K. Y., Tanigaki, S., Sago, H., Sato, S., Saito, S., & Morisaki, N. (2017). Association between very advanced maternal age and adverse pregnancy outcomes: A cross sectional Japanese study. *BMC Pregnancy and Childbirth*, 17(1). <https://doi.org/10.1186/s12884-017-1540-0>
- Olisa, C. L., Nwosu, B. O., Eleje, G. U., Oguejiofor, C. B., Mbachu, I. I., Ogabido, C. A., Njoku, T. K., Okafor, C. C., Okechukwu, Z. C., Okeke, C. F., Okonkwo, I. O., Okaforcha, E. I., Enechukwu, C. I., Ilika, C. P., Nnabuchi, O. K., Osuafor, U. H., Ugwuoroko, H. C., Egwuatu, E. C., Andeh, M. C., & Okafor, C. G. (2024). Comparison of urine protein–creatinine ratio and urine dipstick test for significant proteinuria in preeclamptic women. *Therapeutic Advances in Reproductive Health*, 18.
<https://doi.org/10.1177/26334941241288841>
- Pasternak, Y., Lifshitz, D., Shulman, Y., Hirsch, L., Rimon, E., Kuperminc, M., Yogev, Y., & Ashwal, E. (2021). Diagnostic accuracy of random urinary protein-to-creatinine ratio for proteinuria in patients with suspected pre-eclampsia. *Archives of Gynecology and Obstetrics*, 304(1), 109–115.
<https://doi.org/10.1007/s00404-020-05937-0>
- Payne, B., Magee, L. A., Côté, A. M., Hutcheon, J. A., Li, J., Kyle, P. M., Menzies, J. M., Peter Moore, M., Parker, C., Pullar, B., von Dadelszen, P., Walters, B. N., Douglas, M. J., Walley, K. R., Russell, J. A., Lee, S. K., Gruslin, A., Smith, G. N., Moutquin, J. M., ... Widmer, M. (2011). PIERS Proteinuria: Relationship With Adverse Maternal and Perinatal Outcome. *Journal of Obstetrics and Gynaecology Canada*, 33(6), 588–597.
[https://doi.org/10.1016/S1701-2163\(16\)34907-6](https://doi.org/10.1016/S1701-2163(16)34907-6)
- Perkumpulan Obstetri dan Ginekologi Indonesia. (2016). *Pedoman Nasional Pelayanan Kedokteran Diagnosis dan Tata Laksana Pre-eklampsia*.
- Perry, H., Lehmann, H., Mantovani, E., Thilaganathan, B., & Khalil, A. (2020). Are maternal hemodynamic indices markers of fetal growth restriction in pregnancies with a small-for-gestational-age fetus? *Ultrasound in Obstetrics and Gynecology*, 55(2), 210–216.
<https://doi.org/10.1002/uog.20419>
- Phelan, L. K., Brown, M. A., Davis, G. K., & Mangos, G. (2004). A prospective study of the impact of automated dipstick urinalysis on the diagnosis of preeclampsia. *Hypertension in Pregnancy*, 23(2), 135–142.
<https://doi.org/10.1081/PRG-120028289>

- Poon, L. C., Shennan, A., Hyett, J. A., Kapur, A., Hadar, E., Divakar, H., McAuliffe, F., da Silva Costa, F., von Dadelszen, P., McIntyre, H. D., Kihara, A. B., Di Renzo, G. C., Romero, R., D'Alton, M., Berghella, V., Nicolaidis, K. H., & Hod, M. (2019). The International Federation of Gynecology and Obstetrics (FIGO) initiative on pre-eclampsia: A pragmatic guide for firsttrimester screening and prevention. *International Journal of Gynecology and Obstetrics*, *145*(S1), 1–33. <https://doi.org/10.1002/ijgo.12802>
- Rana, S., Lemoine, E., Granger, J., & Karumanchi, S. A. (2019). Preeclampsia: Pathophysiology, Challenges, and Perspectives. *Circulation Research*, *124*(7), 1094–1112. <https://doi.org/10.1161/CIRCRESAHA.118.313276>
- Redman, C. W. G., & Sargent, I. L. (2010). Immunology of Pre-Eclampsia. In *American Journal of Reproductive Immunology* (Vol. 63, Issue 6, pp. 534–543). Blackwell Publishing Ltd. <https://doi.org/10.1111/j.16000897.2010.00831.x>
- Sewell, M. F., Huston-Presley, L., Super, D. M., & Catalano, P. (2006). Increased neonatal fat mass, not lean body mass, is associated with maternal obesity. *American Journal of Obstetrics and Gynecology*, *195*(4), 1100–1103. <https://doi.org/10.1016/j.ajog.2006.06.014>
- Sohlberg, S., Mulic-Lutvica, A., Lindgren, P., Ortiz-Nieto, F., Wikström, A. K., & Wikström, J. (2014). Placental perfusion in normal pregnancy and early and late preeclampsia: A magnetic resonance imaging study. *Placenta*, *35*(3), 202–206. <https://doi.org/10.1016/j.placenta.2014.01.008>
- Sonkusare, S., Prabhu, L., Shetty, P., Naik, P., Gopal, N., & Kumari, S. (2025). Comparison of proteinuria assessment methods and their association with maternal-fetal outcomes in preeclampsia. *European Journal of Obstetrics and Gynecology and Reproductive Biology*, *311*. <https://doi.org/10.1016/j.ejogrb.2025.114010>
- Tanacan, A., Fadiloglu, E., & Beksac, M. S. (2019). The importance of proteinuria in preeclampsia and its predictive role in maternal and neonatal outcomes. *Hypertension in Pregnancy*, *38*(2), 111–118. <https://doi.org/10.1080/10641955.2019.1590718>
- Teeuw, H. M., Amoakoh, H. B., Ellis, C. A., Lindsley, K., & Browne, J. L. (2022). Diagnostic accuracy of urine dipstick tests for proteinuria in pregnant women suspected of preeclampsia: A systematic review and meta-analysis. In *Pregnancy Hypertension* (Vol. 27, pp. 123–130). Elsevier B.V. <https://doi.org/10.1016/j.preghy.2021.12.015>
- Umesawa, M., & Kobashi, G. (2017). Epidemiology of hypertensive disorders in pregnancy: Prevalence, risk factors, predictors and prognosis. In *Hypertension Research* (Vol. 40, Issue 3, pp. 213–220). Nature Publishing Group. <https://doi.org/10.1038/hr.2016.126>
- Valensise, H., Vasapollo, B., Gagliardi, G., & Novelli, G. P. (2008). Early and Late preeclampsia: Two different maternal hemodynamic states in the latent phase of the disease. *Hypertension*, *52*(5), 873–880. <https://doi.org/10.1161/HYPERTENSIONAHA.108.117358>

- Vasapollo, B., Valensise, H., Novelli, G. P., Altomare, F., Galante, A., & Arduini, D. (2004). Abnormal maternal cardiac function precedes the clinical manifestation of fetal growth restriction. *Ultrasound in Obstetrics and Gynecology*, 24(1), 23–29. <https://doi.org/10.1002/uog.1095>
- Vats, H., Saxena, R., Sachdeva, M. P., Walia, G. K., & Gupta, V. (2021). Impact of maternal pre-pregnancy body mass index on maternal, fetal and neonatal adverse outcomes in the worldwide populations: A systematic review and meta-analysis. In *Obesity Research and Clinical Practice* (Vol. 15, Issue 6, pp. 536–545). Elsevier Ltd. <https://doi.org/10.1016/j.orcp.2021.10.005>
- Vigil-De Gracia, P., Vargas, C., Sánchez, J., & Collantes-Cubas, J. (2023). Preeclampsia: Narrative review for clinical use. *Heliyon*, 9(3). <https://doi.org/10.1016/j.heliyon.2023.e14187>
- Wójtowicz, A., Zembala-Szczerba, M., Babczyk, D., Kołodziejczyk-Pietruszka, M., Lewaczyńska, O., & Huras, H. (2019). Early-and Late-Onset Preeclampsia: A Comprehensive Cohort Study of Laboratory and Clinical Findings according to the New ISHHP Criteria. *International Journal of Hypertension*, 2019. <https://doi.org/10.1155/2019/4108271>
- Xiao, J., Fan, W., Zhu, Q., & Shi, Z. (2022). Diagnosis of proteinuria using a random urine protein-creatinine ratio and its correlation with adverse outcomes in pregnancy with preeclampsia characterized by renal damage. *Journal of Clinical Hypertension*, 24(5), 652–659. <https://doi.org/10.1111/jch.14467>
- Yi, J., Chen, L., Meng, X., & Chen, Y. (2022). Risk factors and foetal growth restriction associated with expectant treatment of early-onset preeclampsia. *Annals of Medicine*, 54(1), 3250–3257. <https://doi.org/10.1080/07853890.2022.2144642>
- Yıldız, G. A., & Yılmaz, E. P. T. (2022). The association between protein levels in 24hour urine samples and maternal and neonatal outcomes of pregnant women with preeclampsia. *Journal of the Turkish German Gynecology Association*, 23(3), 190–198. <https://doi.org/10.4274/jtgga.galenos.2022.2022-4-3>
- Zhu, Y. C., Lin, L., Li, B. Y., Li, X. T., Chen, D. J., Zhao, X. L., Cui, S. H., Ding, H. J., Ding, G. F., Meng, H. X., Wei, H. W., Sun, X. T., Xin, H., & Yang, H. X. (2020). Incidence and Clinical Features of Fetal Growth Restriction in 4 451 Women with Hypertensive Disorders of Pregnancy. *Maternal-Fetal Medicine*, 2(4), 207–210. <https://doi.org/10.1097/FM9.0000000000000062>
- Zhuang, X., Shi, J., Deng, D. R., Zhang, W. S., Liu, X. H., & Lin, J. H. (2020). Initialonset symptoms of preeclampsia and their relation to pregnancy outcomes. *Journal of Obstetrics and Gynaecology*, 40(7), 947–952. <https://doi.org/10.1080/01443615.2019.1676210>