

DAFTAR PUSTAKA

- Buonacera, A. *et al.* (2022) 'Neutrophil to Lymphocyte Ratio: An Emerging Marker of the Relationships between the Immune System and Diseases', *International Journal of Molecular Sciences*, 23(7). doi: 10.3390/ijms23073636.
- Capone, M. *et al.* (2018) 'Baseline neutrophil-to-lymphocyte ratio (NLR) and derived NLR could predict overall survival in patients with advanced melanoma treated with nivolumab', pp. 1–7.
- Duan, J., Pan, L. and Yang, M. (2018) 'Preoperative elevated neutrophil-to-lymphocyte ratio (NLR) and derived NLR are associated with poor prognosis in patients with breast cancer A meta-analysis', *Medicine (United States)*, 97(49). doi: 10.1097/MD.00000000000013340.
- Giudici, N. *et al.* (2021) 'Characteristics of upper urinary tract urothelial carcinoma in the context of bladder cancer: a narrative review', 10(10), pp. 4036–4050. doi: 10.21037/tau-20-1472.
- Gontero, P. *et al.* (2024) 'EAU Guidelines on Upper Urinary Tract Urothelial Carcinoma'.
- Gonzalez, H., Hagerling, C. and Werb, Z. (2018) 'Roles of the immune system in cancer: From tumor initiation to metastatic progression', *Genes and Development*, 32(19–20). doi: 10.1101/GAD.314617.118.
- Huang, Y., Busuttill, R. A. and Boussioutas, A. (2021) 'The Role of Innate Immune Cells in Tumor Invasion and Metastasis', pp. 1–18.
- Janssen, L. M. E. *et al.* (2017) 'The immune system in cancer metastasis: Friend or foe?', *Journal for ImmunoTherapy of Cancer*, 5(1). doi: 10.1186/s40425-017-0283-9.
- Kim, H. S., Lee, H. S. and Ku, J. H. (2023) 'Prognostic impact of postoperative neutrophil-to-lymphocyte ratio on survival outcomes of patients treated with radical nephroureterectomy for upper urinary tract urothelial carcinoma: a single institution retrospective analysis using propensity score mat', *Journal of Cancer*, 14(7). doi: 10.7150/jca.76977.
- Klemm, J. *et al.* (2024) 'Upper Tract Urothelial Carcinoma: A Narrative Review of Current Surveillance Strategies for Non-Metastatic Disease', *Cancers*, 16(1). doi: 10.3390/cancers16010044.
- Li, X. *et al.* (2017) 'Prognostic value of neutrophil-to-lymphocyte ratio in urothelial carcinoma of the upper urinary tract and bladder: A systematic review and meta-analysis', *Oncotarget*, 8(37), pp. 62681–62692. doi: 10.18632/oncotarget.17467.
- Luo, Z. *et al.* (2022) 'What is the Role of the Preoperative Blood-Based Inflammation Biomarkers in the Prognosis of Upper Tract Urothelial Carcinoma With Radical

- Nephroureterectomy? A Single-Centre Retrospective Study', *Technology in Cancer Research and Treatment*, 21. doi: 10.1177/15330338221095667.
- Nishida, A. and Andoh, A. (2025) 'The Role of Inflammation in Cancer: Mechanisms of Tumor Initiation, Progression, and Metastasis', *Cells*, 14(7). doi: 10.3390/cells14070488.
- Petros, F. G. (2020) 'Epidemiology , clinical presentation , and evaluation of upper-tract urothelial carcinoma', 9(4), pp. 1794–1798. doi: 10.21037/tau.2019.11.22.
- Proctor, M. J. *et al.* (2012) 'A derived neutrophil to lymphocyte ratio predicts survival in patients with cancer', *British Journal of Cancer*, 107(4). doi: 10.1038/bjc.2012.292.
- Shao, Y. *et al.* (2020) 'Prognostic value of preoperative lymphocyte-related systemic inflammatory biomarkers in upper tract urothelial carcinoma patients treated with radical nephroureterectomy: a systematic review and meta-analysis', *World Journal of Surgical Oncology*, 18(1). doi: 10.1186/s12957-020-02048-7.
- Vartolomei, M. D. *et al.* (2018) 'Is neutrophil-to-lymphocytes ratio a clinical relevant preoperative biomarker in upper tract urothelial carcinoma? A meta-analysis of 4385 patients', *World Journal of Urology*, 36(7), pp. 1019–1029. doi: 10.1007/s00345-018-2235-5.
- Wood, G. *et al.* (2017) 'Derived neutrophil to lymphocyte ratio as a prognostic factor in patients with advanced colorectal cancer according to RAS and BRAF status: A post-hoc analysis of the MRC COIN study', *Anti-Cancer Drugs*, 28(5). doi: 10.1097/CAD.0000000000000488.
- Wu, S. *et al.* (2025) 'Association between systemic immune-inflammation index and the risk of all-cause, cancer and non-cancer mortality in the general population: results from national health and nutrition examination survey 2005–2018', *BMC Public Health*, 25(1), pp. 1–10. doi: 10.1186/s12889-025-21423-1.
- Yang, T. *et al.* (2021) 'Prognostic value of derived neutrophil-to-lymphocyte ratio (dNLR) in patients with non-small cell lung cancer receiving immune checkpoint inhibitors: A meta-analysis', *BMJ Open*, 11(9). doi: 10.1136/bmjopen-2021-049123.
- Zganjar, A. J., Thiel, D. D. and Lyon, T. D. (2023) 'Diagnosis, workup, and risk stratification of upper tract urothelial carcinoma', *Translational Andrology and Urology*, 12(9). doi: 10.21037/tau-23-45.
- Zhao, Z. *et al.* (2020) 'Preoperative risk classification using neutrophil-to-lymphocyte ratio and albumin for upper tract urothelial carcinoma treated with radical nephroureterectomy', *Cancer Management and Research*, 12. doi: 10.2147/CMAR.S274332.