

## DAFTAR PUSTAKA

- Abdurrosid, L. M. K., Maulana, A., Hapsari, Y. and Nandana, P. I. (2017) “Evaluasi Angka Bebas Batu pada Pasien Batu Ginjal yang Dilakukan ESWL Berdasarkan Letak dan Ukuran Batu di Rumah Sakit Harapan Keluarga Mataram Periode 2015-2016,” *Jurnal Kedokteran*, 6(3 SE-Research), p. 11. doi: 10.29303/jku.v6i3.140.
- Abou-Farha, M., El-Abd, A., Gameel, T., Eltatawy, H. and Nagla, S. (2022) “Efficacy of extracorporeal shockwave lithotripsy, with modified position of the machine head in the treatment of lower calyceal stones in obese patients.,” *Urology annals*, 14(1), pp. 81–84. doi: 10.4103/UA.UA\_176\_20.
- Al-Marhoon, M. S., Shareef, O., Al-Habsi, I. S., Al Balushi, A. S., Mathew, J. and Venkiteswaran, K. P. (2013) “Extracorporeal Shock-wave Lithotripsy Success Rate and Complications: Initial Experience at Sultan Qaboos University Hospital.,” *Oman medical journal*, 28(4), pp. 255–259. doi: 10.5001/omj.2013.72.
- Alelign, T. and Petros, B. (2018) “Kidney Stone Disease: An Update on Current Concepts,” *Advances in Urology*, 2018. doi: 10.1155/2018/3068365.
- Alić, J., Heljić, J., Hadžiosmanović, O., Kulovac, B., Lepara, Z., Spahović, H., *et al.* (2022) “The Efficiency of Extracorporeal Shock Wave Lithotripsy (ESWL) in the Treatment of Distal Ureteral Stones: An Unjustly Forgotten Option?,” *Cureus*, 14(9), p. e28671. doi: 10.7759/cureus.28671.
- Bilqisthi, A. R., Prasetyo, B. and Romadhoni, R. (2023) “Korelasi Ukuran Batu Ginjal dengan Jumlah Dilakukan ESWL di Rumah Sakit Islam Sultan Agung pada Tahun 2019: Correlation of Kidney Stone Size with the Frequency of ESWL Performed at the Sultan Agung Hospital in 2019,” *Jurnal Surya Medika (JSM)*, 9(2 SE-Articles), pp. 120–125. doi: 10.33084/jsm.v9i2.3618.
- Curhan, G. C., Willett, W. C., Knight, E. L. and Stampfer, M. J. (2004) “Dietary factors and the risk of incident kidney stones in younger women: Nurses’ Health Study II.,” *Archives of internal medicine*, 164(8), pp. 885–891. doi:

10.1001/archinte.164.8.885.

- Dede, O., Şener, N. C., Baş, O., Dede, G. and Bağbancı, M. Ş. (2015) “Does morbid obesity influence the success and complication rates of extracorporeal shockwave lithotripsy for upper ureteral stones?,” *Turkish journal of urology*, 41(1), pp. 20–23. doi: 10.5152/tud.2015.94824.
- El-Nahas, A. R., El-Assmy, A. M., Mansour, O. and Sheir, K. Z. (2007) “A prospective multivariate analysis of factors predicting stone disintegration by extracorporeal shock wave lithotripsy: the value of high-resolution noncontrast computed tomography.,” *European urology*, 51(6), pp. 1684–1688. doi: 10.1016/j.eururo.2006.11.048.
- Ferraro, P. M., Bargagli, M., Trinchieri, A. and Gambaro, G. (2020) “Risk of Kidney Stones: Influence of Dietary Factors, Dietary Patterns, and Vegetarian-Vegan Diets.,” *Nutrients*, 12(3). doi: 10.3390/nu12030779.
- Ferraro, P. M., Taylor, E. N., Eisner, B. H., Gambaro, G., Rimm, E. B., Mukamal, K. J., *et al.* (2013) “History of kidney stones and the risk of coronary heart disease.,” *JAMA*, 310(4), pp. 408–415. doi: 10.1001/jama.2013.8780.
- Lang, J., Narendrula, A., El-Zawahry, A., Sindhvani, P. and Ekwenna, O. (2022) “Global Trends in Incidence and Burden of Urolithiasis from 1990 to 2019: An Analysis of Global Burden of Disease Study Data,” *European Urology Open Science*, 35. doi: 10.1016/j.euros.2021.10.008.
- Liu, Y., Chen, Y., Liao, B., Luo, D., Wang, K., Li, H., *et al.* (2018) “Epidemiology of urolithiasis in Asia,” *Asian Journal of Urology*, 5(4). doi: 10.1016/j.ajur.2018.08.007.
- Maalouf, N. M., Sato, A. H., Welch, B. J., Howard, B. V, Cochrane, B. B., Sakhaee, K., *et al.* (2010) “Postmenopausal hormone use and the risk of nephrolithiasis: results from the Women’s Health Initiative hormone therapy trials.,” *Archives of internal medicine*, 170(18), pp. 1678–1685. doi: 10.1001/archinternmed.2010.342.
- Massoud, A. M., Abdelbary, A. M., Al-Dessoukey, A. A., Moussa, A. S., Zayed, A. S. and Mahmmoud, O. (2014) “The success of extracorporeal shock-wave

- lithotripsy based on the stone-attenuation value from non-contrast computed tomography.” *Arab journal of urology*, 12(2), pp. 155–161. doi: 10.1016/j.aju.2014.01.002.
- Mohamed, H. H. A., Abdelmaksoud, M. M., El Gharbawy, M. S. and Badreldin, M. R. (2022) “Effect of Hounsfield unit of renal stones and skin-to-stone distance detected by computed tomography on extracorporeal shock wave lithotripsy outcome,” *Menoufia Medical Journal*, 35(3), pp. 1604–1607.
- Muñoz, R. D., Tirolien, P. P., Belhamou, S., Desta, M., Grimberg, R., Dulys, P., *et al.* (2003) “[Treatment of reno-ureteral lithiasis with ESWL in obese patients. Apropos of 150 patients].” *Archivos espanoles de urologia*, 56(8), pp. 933–938.
- Naghii, M. R., Babaei, M. and Hedayati, M. (2014) “Androgens involvement in the pathogenesis of renal stones formation.” *PloS one*, 9(4), p. e93790. doi: 10.1371/journal.pone.0093790.
- Natalin, R., Xavier, K., Okeke, Z. and Gupta, M. (2009) “Impact of obesity on ureteroscopic laser lithotripsy of urinary tract calculi.” *International braz j urol : official journal of the Brazilian Society of Urology*, 35(1), pp. 32–36. doi: 10.1590/s1677-55382009000100006.
- Oliveira, B., Teixeira, B., Magalhães, M., Vinagre, N., Fraga, A. and Cavadas, V. (2024) “Extracorporeal shock wave lithotripsy: retrospective study on possible predictors of treatment success and revisiting the role of non-contrast-enhanced computer tomography in kidney and ureteral stone disease,” *Urolithiasis*, 52(1), p. 65. doi: 10.1007/s00240-024-01570-7.
- Partin, A. W., Wein, A. J., Kavoussi, L. R., Peters, C. A. and Dmochowski, R. R. (2020) *Campbell Walsh Wein Urology, E-Book*. Elsevier Health Sciences.
- Roy, P., Sarkar, D., Jalan, V. and Pal, D. K. (2021) “Comparative Study of Extracorporeal Shock Wave Lithotripsy Versus Mini Percutaneous Nephrolithotomy for the Treatment of Nonlower Calyceal 10–20 mm Size Kidney Stone,” *Urological Science*, 32(2). Available at: [https://journals.lww.com/ursc/fulltext/2021/32020/comparative\\_study\\_of\\_ext](https://journals.lww.com/ursc/fulltext/2021/32020/comparative_study_of_ext)

racorporeal\_shock\_wave.8.aspx.

- Sakhaee, K., Maalouf, N. M. and Sinnott, B. (2012) “Clinical review. Kidney stones 2012: pathogenesis, diagnosis, and management.,” *The Journal of clinical endocrinology and metabolism*, 97(6), pp. 1847–1860. doi: 10.1210/jc.2011-3492.
- Scales, C. D. J., Smith, A. C., Hanley, J. M. and Saigal, C. S. (2012) “Prevalence of kidney stones in the United States.,” *European urology*, 62(1), pp. 160–165. doi: 10.1016/j.eururo.2012.03.052.
- Skolarikos, A., Jung, H., Neisius, A., Petřík, A., Somani, B., Tailly, T., *et al.* (2024) “EAU Guidelines on Urolithiasis,” *European Association of Urology 2021*, (March), pp. 1–87. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/26344917> [http://uroweb.org/wp-content/uploads/22-Urolithiasis\\_LR\\_full.pdf](http://uroweb.org/wp-content/uploads/22-Urolithiasis_LR_full.pdf).
- Snicorius, M., Drevinskaitė, M., Miglinas, M., Čekauskas, A., Urbonienė, V., Bandzevičiūtė, R., *et al.* (2023) “A Prospective Study on the Impact of Clinical Factors and Adjusted Triple D System for Success Rate of ESWL,” *Medicina*. doi: 10.3390/medicina59101827.
- Stamatelou, K. and Goldfarb, D. S. (2023) “Epidemiology of Kidney Stones,” *Healthcare (Switzerland)*, 11(3). doi: 10.3390/healthcare11030424.
- Taylor, E. N., Stampfer, M. J. and Curhan, G. C. (2005) “Obesity, weight gain, and the risk of kidney stones.,” *JAMA*, 293(4), pp. 455–462. doi: 10.1001/jama.293.4.455.
- Wang, L.-J., Wong, Y.-C., Chuang, C.-K., Chu, S.-H., Chen, C.-S., See, L.-C., *et al.* (2005) “Predictions of outcomes of renal stones after extracorporeal shock wave lithotripsy from stone characteristics determined by unenhanced helical computed tomography: a multivariate analysis.,” *European radiology*, 15(11), pp. 2238–2243. doi: 10.1007/s00330-005-2742-9.
- Yan, C.-P., Wang, X.-K., Jiang, K., Yin, C., Xiang, C., Wang, Y., *et al.* (2022) “ $\beta$ -Ecdysterone Enhanced Bone Regeneration Through the BMP-2/SMAD/RUNX2/Osterix Signaling Pathway.,” *Frontiers in cell and*

*developmental biology*, 10, p. 883228. doi: 10.3389/fcell.2022.883228.

Yoon, J. H., Park, Sejun, Kim, S. C., Park, Sungchan, Moon, K. H., Cheon, S. H., *et al.* (2021) “Outcomes of extracorporeal shock wave lithotripsy for ureteral stones according to ESWL intensity,” *Translational Andrology and Urology; Vol 10, No 4 (April 26, 2021): Translational Andrology and Urology*. Available at: <https://tau.amegroups.org/article/view/63248>.