

DAFTAR PUSTAKA

- Antunes, D.E. *et al.* (2013) 'Identification of clinical, epidemiological and laboratory risk factors for leprosy reactions during and after multidrug therapy.', *Memorias do Instituto Oswaldo Cruz*, 108(7), pp. 901–908. Available at: <https://doi.org/10.1590/0074-0276130222>.
- Arunraghav, P. and Herakal, K. (2021) 'Leprosy in Elderly and Children among New Cases - A 3-Year Retrospective Study.', *Indian dermatology online journal*, 12(2), pp. 294–297. Available at: https://doi.org/10.4103/idoj.IDOJ_177_18.
- Bakshi, N., Rao, S. and Batra, R. (2017) 'Bullous Erythema Nodosum Leprosum as the First Manifestation of Multibacillary Leprosy: A Rare Phenomenon.', *The American Journal of dermatopathology*, 39(11), pp. 857–859.
- Benites-Zapata, V.A. *et al.* (2015) 'Usefulness of Neutrophil-to-Lymphocyte Ratio in Risk Stratification of Patients With Advanced Heart Failure', *The American Journal of Cardiology*, 115(1), pp. 57–61. Available at: <https://doi.org/https://doi.org/10.1016/j.amjcard.2014.10.008>.
- Chaudhary, D.R. and Modi, D.K. (2015) 'Azathioprine as a Steroid Sparing Agent in Recalcitrant Type 2 Reaction', in. Available at: <https://api.semanticscholar.org/CorpusID:47000094>.
- Dunn, G.P. *et al.* (2002) 'Cancer immunoediting: from immunosurveillance to tumor escape', *Nature Immunology*, 3(11), pp. 991–998. Available at: <https://doi.org/10.1038/ni1102-991>.
- Gomes, L.T. *et al.* (2020) 'Diagnostic value of neutrophil-to-lymphocyte ratio in patients with leprosy reactions', *Heliyon*, 6(2). Available at: <https://doi.org/10.1016/j.heliyon.2020.e03369>.
- Gupta, S. *et al.* (2017) 'Clinical profile of type II reaction in leprosy - A cross sectional study', *Indian Journal of Leprosy*, 89, pp. 191–195.
- Kahawita, I.P. and Lockwood, D.N.J. (2008) 'Towards understanding the pathology of erythema nodosum leprosum.', *Transactions of the Royal Society of Tropical Medicine and Hygiene*, 102(4), pp. 329–337.
- Kar, H.K. and Kumar, B. (2010) *IAL textbook of leprosy, TA - TT* -. New Delhi, India SE - xix, 624 pages : color illustrations, color maps ; 28 cm: Jaypee Brothers Medical Publishers. Available at: <https://doi.org/LK> - <https://worldcat.org/title/715111442>.

- Kumar, B., Dogra, S. and Kaur, I. (2004) 'Epidemiological characteristics of leprosy reactions: 15 years experience from north India.', *International journal of leprosy and other mycobacterial diseases : official organ of the International Leprosy Association*, 72(2), pp. 125–133.
- Lee, D.J., Rea, T.H. and Modlin, R.L. (2012) 'Chapter 186. Leprosy', in L.A. Goldsmith et al. (eds) *Fitzpatrick's Dermatology in General Medicine*, 8e. New York, NY: The McGraw-Hill Companies.
- Lockwood, D.N.J. and Suneetha, S. (2005) 'Leprosy: too complex a disease for a simple elimination paradigm.', *Bulletin of the World Health Organization*, 83(3), pp. 230–235.
- Lubis, R.S. et al. (2022) 'Epidemiology of Leprosy in Indonesia: a Retrospective Study', *Berkala Ilmu Kesehatan Kulit dan Kelamin*, 34(1 SE-Research), pp. 29–35. Available at: <https://doi.org/10.20473/bikk.V34.1.2022.29-35>.
- Made, L. et al. (2024) 'Neutrophil-to-lymphocyte ratio (RNL) as a severity predictor of leprosy reaction', *Bali Medical Journal*, 13(2), pp. 791–795. Available at: <https://doi.org/10.15562/BMJ.V13I2.5164>.
- Mittal, D. et al. (2014) 'New insights into cancer immunoediting and its three component phases—elimination, equilibrium and escape', *Current Opinion in Immunology*, 27, pp. 16–25. Available at: <https://doi.org/https://doi.org/10.1016/j.coi.2014.01.004>.
- Motta, A.C.F. et al. (2012) 'Leprosy reactions: coinfections as a possible risk factor.', *Clinics (Sao Paulo, Brazil)*, 67(10), pp. 1145–1148. Available at: [https://doi.org/10.6061/clinics/2012\(10\)05](https://doi.org/10.6061/clinics/2012(10)05).
- Nayak, D. et al. (2024) 'Risk Factors for Erythema Nodosum Leprosum: A Case Control Study in a Tertiary Hospital of Western Odisha, India', *Indian Journal of Leprosy*, pp. 17–25.
- Papa, A. et al. (2008) 'Predictive value of elevated neutrophil–lymphocyte ratio on cardiac mortality in patients with stable coronary artery disease', *Clinica Chimica Acta*, 395(1), pp. 27–31.
- Riley, L.K. and Rupert, J. (2015) 'Evaluation of Patients with Leukocytosis.', *American family physician*, 92(11), pp. 1004–1011.
- Sardana, K. and Khurana, A. (2023) 'Jopling's Handbook of Leprosy 7th Edition'. New Delhi: CBS Publishers & Distributors PP - New Delhi.
- Scollard, D.M. et al. (2006) 'The continuing challenges of leprosy.', *Clinical microbiology reviews*, 19(2), pp. 338–381. Available at: <https://doi.org/10.1128/CMR.19.2.338-381.2006>.



- Singal, A. (2020) 'Current Concepts and Challenges in the Management of Erythema Nodosum Leprosum', *Indian Dermatology Online Journal*, 11(4), p. 479. Available at: https://doi.org/10.4103/IDJ.IDJ_69_20.
- Song, M. *et al.* (2021) 'Neutrophil-to-lymphocyte ratio and mortality in the United States general population.', *Scientific reports*, 11(1), p. 464. Available at: <https://doi.org/10.1038/s41598-020-79431-7>.
- Suryawati, N. and Saputra, H. (2018) 'Erythema nodosum leprosum presenting as sweet's syndrome-like reaction in a borderline lepromatous leprosy patient', *International Journal of Mycobacteriology*, 7, pp. 191–194. Available at: <https://api.semanticscholar.org/CorpusID:49184993>.
- Tanojo, N. *et al.* (2022) 'Diagnostic Value of Neutrophil-to-Lymphocyte Ratio, Lymphocyte-to-Monocyte Ratio, and Platelet-to-Lymphocyte Ratio in the Diagnosis of Erythema Nodosum Leprosum: A Retrospective Study', *Tropical Medicine and Infectious Disease*, 7(3), p. 39.
- Voorend, C.G.N. and Post, E.B. (2013) 'A Systematic Review on the Epidemiological Data of Erythema Nodosum Leprosum, a Type 2 Leprosy Reaction', *PLoS Neglected Tropical Diseases*, 7(10), p. e2440.
- Walker, S. *et al.* (2015) 'ENLIST 1: An International Multi-centre Cross-sectional Study of the Clinical Features of Erythema Nodosum Leprosum', *PLoS neglected tropical diseases*, 9, p. e0004065. Available at: <https://doi.org/10.1371/journal.pntd.0004065>.
- World Health Organization (2023) *Global leprosy (Hansen disease) update, 2023: accelerating reduction of disease burden*. Available at: <https://iris.who.int/handle/10665/373164>.
- Yamanaka, T. *et al.* (2008) 'The Baseline Ratio of Neutrophils to Lymphocytes Is Associated with Patient Prognosis in Advanced Gastric Cancer', *Oncology*, 73(3–4), pp. 215–220. Available at: <https://doi.org/10.1159/000127412>.
- Zahorec, R. (2021) 'Neutrophil-to-lymphocyte ratio, past, present and future perspectives.', *Bratislavske lekarske listy*, 122(7), pp. 474–488. Available at: https://doi.org/10.4149/BLL_2021_078.