

Daftar Pustaka

- Abd-Alla, M. H., Al-Amri, S. M., & El-Enany, A. W. E. (2023). Enhancing rhizobium–legume symbiosis and reducing nitrogen fertilizer use are potential options for mitigating climate change. *Agriculture (Switzerland)*, *13*(11). <https://doi.org/10.3390/agriculture13112092>
- Adams, M. A., Buckley, T. N., Salter, W. T., Buchmann, N., Blessing, C. H., Turnbull, T. L., & Adams, M. A. (2018). *Contrasting responses of crop legumes and cereals to nitrogen availability*. 1475–1483. <https://doi.org/10.1111/nph.14918>
- Adie, M. M., & Krisnawati, A. (2016). *Biologi tanaman kedelai*. Balai Penelitian Tanaman Kacang-kacangan dan Umbi-umbian.
- Adisarwanto. (2005). *Budidaya kedelai dengan pemupukan yang efektif dan pengoptimalan peran bintil Akar*. Penebar Swadaya.
- Ahmed, S., Raza, M. A., Yuan, X., Du, Y., Iqbal, N., Chachar, Q., Soomro, A. A., Ibrahim, F., Hussain, S., Wang, X., Liu, W., & Yang, W. (2020). Optimized planting time and co-growth duration reduce the yield difference between intercropped and sole soybean by enhancing soybean resilience toward size-asymmetric competition. *Food and Energy Security*, *9*, e226. <https://doi.org/10.1002/fes3.226>
- Ahmed, S., Raza, M., Zhou, T., Hussain, S., Khalid, M., Feng, L., Wasaya, A., Iqbal, N., Ahmed, A., Liu, W., & Yang, W. (2018). Responses of soybean dry matter production, phosphorus accumulation, and seed yield to sowing time under relay intercropping with maize. *Agronomy*, *8*(12), 282. <https://doi.org/https://doi.org/10.3390/agronomy8120282>
- Akchaya, K., Parasuraman, P., Pandian, K., Vijayakumar, S., Thirukumaran, K., Mustaffa, M. R. A. F., Rajpoot, S. K., & Choudhary, A. K. (2025). Boosting resource use efficiency, soil fertility, food security, ecosystem services, and climate resilience with legume intercropping: a review. *Frontiers in Sustainable Food Systems*, *9*, 1–22. <https://doi.org/10.3389/fsufs.2025.1527256>
- Andrews, D. J., & Kassam, A. H. (1976). The importance of multiple cropping in increasing world food supplies. In R. I. Papendick, P. A. Sanchez, & G. B. Triplett (Eds.), *Multiple cropping* (pp. 1–10). ASA Special Publications. <https://doi.org/10.2134/asaspecpub27.c1>
- Andrianto, T. ., & Indarto, N. (2004). *Budidaya dan analisis usaha tani kedelai, kacang hijau, kacang panjang*. Absolut.
- Araújo, W. A. de, Santana, R. S., Mauad, M., & da Silva, R. S. (2021). Dry matter accumulation and nutrient uptake in determinate and indeterminate soybeans. *Journal of Plant Nutrition*, *44*(4), 508–522. <https://doi.org/10.1080/01904167.2020.1845371>
- Ashworth, A. J., West, C. P., Allen, F. L., Keyser, P. D., Weiss, S. A., Tyler, D. D., Taylor, A. M., Warwick, K. L., & Beamer, K. P. (2015). Biologically fixed nitrogen in legume intercropped systems: Comparison of nitrogen-difference and nitrogen-15 enrichment techniques. *Agronomy Journal*, *107*(6), 2419–2430. <https://doi.org/10.2134/agronj14.0639>
- Astutik, D., Yudono, P., & Waluyo, S. (2018). The growth of two cultivars mung bean under different sweet corn shelter density in sandy soil coastal area. *Ilmu Pertanian (Agricultural Science)*, *2*(3), 106. <https://doi.org/10.22146/ipas.26703>
- Badan Pusat Statistik. (2021). *Analisis produktivitas jagung dan kedelai di Indonesia, 2021*.

BPS-RI.

- Bakal, H., Gulluoglu, L., Onat, B., & Arioglu, H. (2017). The effect of growing seasons on some agronomic and quality characteristics of Soybean varieties in mediterranean region in Turkey. *Turkish Journal of Field Crops*, 22(2), 187–196. <https://doi.org/10.17557/tjfc.356213>
- Balitkabi. (2021). *Denasa 1 dan Denasa 2 varietas unggul baru kedelai toleran naungan dan berbiji besar*. <https://balitkabi.litbang.pertanian.go.id/infotek/denasa-1-dan-denasa-2-varietas-unggul-baru-kedelai-toleran-naungan-dan-berbiji-besar/>
- Becana, M., & Sprent, J. I. (1987). Nitrogen fixation and nitrate reduction in the root nodules of legumes. *Physiologia Plantarum*, 70(4), 757–765. <https://doi.org/10.1111/j.1399-3054.1987.tb04335.x>
- Bedoussac, L., Journet, E. P., Hauggaard-Nielsen, H., Naudin, C., Corre-Hellou, G., Jensen, E. S., Prieur, L., & Justes, E. (2015). Ecological principles underlying the increase of productivity achieved by cereal-grain legume intercrops in organic farming. A review. *Agronomy for Sustainable Development*, 35(3), 911–935. <https://doi.org/10.1007/s13593-014-0277-7>
- Bertham, Y. H., Aini, N., Gonggo Murcitro, B., & Dipo Nusantara, A. (2018). Uji Coba Empat Varietas Kedelai di Kawasan Pesisir Berbasis Biokompos. *Biogenesis: Jurnal Ilmiah Biologi*, 6(1), 36–42. <https://doi.org/10.24252/bio.v6i1.4144>
- Boote, K. J., Prasad, V., Allen, L. H., Singh, P., & Jones, J. W. (2018). Modeling sensitivity of grain yield to elevated temperature in the DSSAT crop models for peanut, soybean, dry bean, chickpea, sorghum, and millet. *European Journal of Agronomy*, 100, 99–109. <https://doi.org/10.1016/j.eja.2017.09.002>
- Bouchet, A. S., Laperche, A., Bissuel-Belaygue, C., Snowdon, R., Nesi, N., & Stahl, A. (2016). Nitrogen use efficiency in rapeseed. A review. *Agronomy for Sustainable Development*, 36(2). <https://doi.org/10.1007/s13593-016-0371-0>
- Boyd, E. S., & Peters, J. W. (2013). New insights into the evolutionary history of biological nitrogen fixation. *Frontiers in Microbiology*, 4, 1–12. <https://doi.org/10.3389/fmicb.2013.00201>
- Brooker, R. W., Bennett, A. E., Cong, W. F., Daniell, T. J., George, T. S., Hallett, P. D., Hawes, C., Iannetta, P. P. M., Jones, H. G., Karley, A. J., Li, L., Mckenzie, B. M., Pakeman, R. J., Paterson, E., Schöb, C., Shen, J., Squire, G., Watson, C. A., Zhang, C., ... White, P. J. (2015). Improving intercropping: A synthesis of research in agronomy, plant physiology and ecology. *New Phytologist*, 206, 107–117. <https://doi.org/10.1111/nph.13132>
- Budiyanto, G. (2014). *Manajemen Sumber Daya Lahan*. LP3M UMY. <http://repository.umy.ac.id/handle/123456789/6405>
- Caddel, J., Redfean, D., & Edwards, J. (2017). Forage legumes and nitrogen production. *Oklohoma Cooperative Extension Service*, 1–6. <http://pods.dasnr.okstate.edu/docushare/dsweb/Get/Document-3101/PSS-2590webcolor.pdf>
- Cahyono. (2007). *Kedelai*. CV. Aneka Ilmu.
- Canadell, P., Tian, H., Patra, P., & Thomson, R. (2019). *Nitrogen fertilisers are incredibly efficient, but they make climate change a lot worse*. <https://theconversation.com/nitrogen-fertilisers-are-incredibly-efficient-but-they-make-climate-change-a-lot-worse-127103>

- Carley, C. N., Zubrod, M. J., Dutta, S., & Singh, A. K. (2023). Using machine learning enabled phenotyping to characterize nodulation in three early vegetative stages in soybean. *Crop Science*, 63(1), 204–226. <https://doi.org/10.1002/csc2.20861>
- Carranca, C. (2014). *Legumes: Properties and symbiosis* (Issue January 2013).
- Casal, J. J. (2012). Shade Avoidance. *The Arabidopsis Book*, 10, e0157. <https://doi.org/10.1199/tab.0157>
- Chang, X., Wei, D., Zeng, Y., Zhao, X., Hu, Y., Wu, X., Song, C., Gong, G., Chen, H., Yang, C., Zhang, M., Liu, T., Chen, W., & Yang, W. (2022). Maize-soybean relay strip intercropping reshapes the rhizosphere bacterial community and recruits beneficial bacteria to suppress Fusarium root rot of soybean. *Frontiers in Microbiology*, 13, 1009689. <https://doi.org/10.3389/fmicb.2022.1009689>
- Chang, X., Yan, L., Naeem, M., Khaskheli, M. I., Zhang, H., Gong, G., Zhang, M., Song, C., Yang, W., Liu, T., & Chen, W. (2020). Maize/soybean relay strip intercropping reduces the occurrence of fusarium root rot and changes the diversity of the pathogenic fusarium species. *Pathogens*, 9(3), 211. <https://doi.org/10.3390/pathogens9030211>
- Chapagain, T., & Riseman, A. (2014). Barley-pea intercropping: Effects on land productivity, carbon and nitrogen transformations. *Field Crops Research*, 166, 18–25. <https://doi.org/10.1016/j.fcr.2014.06.014>
- Chen, P., Du, Q., Liu, X., Zhou, L., Hussain, S., Lei, L., Song, C., Wang, X., Liu, W., Yang, F., Shu, K., Liu, J., Du, J., Yang, W., & Yong, T. (2017). Effects of reduced nitrogen inputs on crop yield and nitrogen use efficiency in a long-term maize-soybean relay strip intercropping system. *Plos One*, 12(9), e0184503. <https://doi.org/10.1371/journal.pone.0184503>
- Chen, P., Song, C., Liu, X. ming, Zhou, L., Yang, H., Zhang, X., Zhou, Y., Du, Q., Pang, T., Fu, Z. dan, Wang, X. chun, Liu, W. guo, Yang, F., Shu, K., Du, J., Liu, J., Yang, W., & Yong, T. (2019). Yield advantage and nitrogen fate in an additive maize-soybean relay intercropping system. *Science of the Total Environment*, 657, 987–999. <https://doi.org/10.1016/j.scitotenv.2018.11.376>
- Chen, X., Chen, J., & Cao, J. (2023). Intercropping increases soil N-targeting enzyme activities: A meta-analysis. *Rhizosphere*, 26, 100686. <https://doi.org/10.1016/j.rhisph.2023.100686>
- Cheng, B., Wang, L., Liu, R., Wang, W., Yu, R., Zhou, T., Ahmad, I., Raza, A., Jiang, S., Xu, M., Liu, C., Yu, L., Wang, W., Jing, S., Liu, W., & Yang, W. (2022). Shade-Tolerant Soybean Reduces Yield Loss by Regulating Its Canopy Structure and Stem Characteristics in the Maize–Soybean Strip Intercropping System. *Frontiers in Plant Science*, 13(March), 1–16. <https://doi.org/10.3389/fpls.2022.848893>
- Cheng, Z., Meng, L., Yin, T., Li, Y., Zhang, Y., & Li, S. (2023). Changes in Soil Rhizobia Diversity and Their Effects on the Symbiotic Efficiency of Soybean Intercropped with Maize. *Agronomy*, 13(4). <https://doi.org/10.3390/agronomy13040997>
- Ciampitti, I. A., de Borja Reis, A. F., Córdova, S. C., Castellano, M. J., Archontoulis, S. V., Correndo, A. A., Antunes De Almeida, L. F., & Moro Rosso, L. H. (2021). Revisiting biological nitrogen fixation dynamics in soybeans. *Frontiers in Plant Science*, 12, 1–11. <https://doi.org/10.3389/fpls.2021.727021>
- Cigelske, B. D., Kandel, H., & DeSutter, T. M. (2020). Soybean Nodulation and Plant Response to Nitrogen and Sulfur Fertilization in the Northern US. *Agricultural Sciences*, 11(06), 592–607. <https://doi.org/10.4236/as.2020.116037>

- De Costa, W. A. J. M., & Perera, M. K. K. W. (1998). Effects of bean population and row arrangement on the productivity of chilli/dwarf bean (*Capsicum annuum*/Phaseolus vulgaris L.) intercropping in Sri Lanka. *Journal of Agronomy and Crop Science*, 180(1), 53–58. <https://doi.org/10.1111/j.1439-037x.1998.tb00369.x>
- de Souza, C. H. E., Ribeiro, V. G. S., Goncalves, L. L., de Borba, M. G., & dos Anjos Reis, R. (2022). Phosphorus fertilizer with increased efficiency affects soybean yields. *Australian Journal of Crop Science*, 16(7), 893–898. <https://doi.org/10.21475/ajcs.22.16.07.p3538>
- Díaz-Pérez, J. C. (2013). Bell pepper (*Capsicum annuum* L.) crop as affected by Shade level: Microenvironment, plant growth, leaf gas exchange, and leaf mineral nutrient concentration. *HortScience*, 48(2), 175–182. <https://doi.org/10.21273/hortsci.48.2.175>
- Distan Yogyakarta. (2009). *Standard operating procedure (SOP): Budidaya cabai merah Kulonprogo [Standard operating procedure (SOP): Red chili cultivation in Kulonprogo]*. Dinas Pertanian Daerah Istimewa Yogyakarta.
- Du, Q., Chen, P., Zheng, B., Hu, Y., Yang, W., & Yong, T. (2022). Screening soybean for adaptation to relay intercropping systems: associations between reproductive organ abscission and yield. *Agronomy*, 12(10), 2379. <https://doi.org/10.3390/agronomy12102379>
- Du, Q., Zhou, L., Chen, P., Liu, X., Song, C., Yang, F., Wang, X., Liu, W., Sun, X., Du, J., Liu, J., Shu, K., Yang, W., & Yong, T. (2020). Relay-intercropping soybean with maize maintains soil fertility and increases nitrogen recovery efficiency by reducing nitrogen input. *The Crop Journal*, 8(1), 140–152. <https://doi.org/10.1016/j.cj.2019.06.010>
- Duchene, O., Vian, J. F., & Celette, F. (2017). Intercropping with legume for agroecological cropping systems: Complementarity and facilitation processes and the importance of soil microorganisms. A review. *Agriculture, Ecosystems & Environment*, 240, 148–161. <https://doi.org/10.1016/J.AGEE.2017.02.019>
- Ebbisa, A. (2022). Mechanisms underlying cereal/legume intercropping as nature-based biofortification: A review. In *Food Production, Processing and Nutrition* (Vol. 4, Issue 1). <https://doi.org/10.1186/s43014-022-00096-y> LK - <https://doi.org/10.1186/s43014-022-00096-y>
- Elhady, A., Hallmann, J., & Heuer, H. (2020). Symbiosis of Soybean With Nitrogen Fixing Bacteria Affected by Root Lesion Nematodes in a Density-Dependent Manner. *Scientific Reports*, 10(1). <https://doi.org/10.1038/s41598-020-58546-x>
- Fageria, N. K., & Baligar, V. C. (2005). Enhancing Nitrogen Use Efficiency in Crop Plants. *Advances in Agronomy*, 88(05), 97–185. [https://doi.org/10.1016/S0065-2113\(05\)88004-6](https://doi.org/10.1016/S0065-2113(05)88004-6)
- Fan, Y., Chen, J., Cheng, Y., Raza, M. A., Wu, X., Wang, Z., Liu, Q., Wang, R., Wang, X., Yong, T., Liu, W., Liu, J., Du, J., Shu, K., Yang, W., & Yang, F. (2018). Effect of shading and light recovery on the growth, leaf structure, and photosynthetic performance of soybean in a maize-soybean relay-strip intercropping system. *PLoS ONE*, 13(5), 1–15. <https://doi.org/10.1371/journal.pone.0198159>
- Faozi, K., Yudono, P., Indradewa, D., & Maas, A. (2018). Banana stem bokashi and its effect to increase soybean yield (*Glycine max* L. Merrill) in coastal sands Area. *Agrotechnology*, 7(2), 1–7. <https://doi.org/10.4172/2168-9881.1000184>
- Fattah, A., Idaryani, Herniwati, Yasin, M., Suriani, S., Salim, Nappu, M. B., Mulia, S., Irawan Hannan, M. F., Wulanningtyas, H. S., Saenong, S., Dewayani, W., Suriyani, Winanda, E., Manwan, S. W., Asaad, M., Warda, Nurjanani, Nurhafsah, ... Ella, A. (2024).

- Performance and morphology of several soybean varieties and responses to pests and diseases in South Sulawesi. *Heliyon*, 10(5), e25507. <https://doi.org/10.1016/j.heliyon.2024.e25507>
- Feng, L., Hu, Y., Shi, K., Tang, H., Pu, T., Wang, X., & Yang, W. (2024). Synergistic effects of crop aboveground growth and root traits guarantee stable yield of strip relay intercropping maize. *Agronomy*, 14(3), 527. <https://doi.org/10.3390/agronomy14030527> LK
- Feng, L., Raza, M. A., Chen, Y., Khalid, M. H. Bin, Meraj, T. A., Ahsan, F., Fan, Y., Du, J., Wu, X., Song, C., Liu, C., Bawa, G., Zhang, Z., Yuan, S., Yang, F., & Yang, W. (2019). Narrow-wide row planting pattern improves the light environment and seed yields of intercrop species in relay intercropping system. *PLOS ONE*, 14(2), e0212885. <https://doi.org/10.1371/journal.pone.0212885>
- Feng, L., Yang, W., Zhou, Q., Tang, H., Ma, Q., Huang, G., & Wang, S. (2021). Effects of interspecific competition on crop yield and nitrogen utilisation in maize-soybean intercropping system. *Plant, Soil and Environment*, 67(8), 460–467. <https://doi.org/10.17221/665/2020-PSE>
- Franklin, K. A. (2008). Shade avoidance. *New Phytologist*, 179(4), 930–944. <https://doi.org/10.1111/j.1469-8137.2008.02507.x>
- Gecaitė, V., Arlauskienė, A., & Cesevičienė, J. (2021). Competition Effects and Productivity in Oat–Forage Legume Relay Intercropping Systems under Organic Farming Conditions. *Agriculture*, 11(2), 99. <https://doi.org/10.3390/agriculture11020099>
- Gitari, H. I., Karanja, N. N., Gachene, C. K. K., Kamau, S., Sharma, K., & Schulte-Geldermann, E. (2018). Nitrogen and phosphorous uptake by potato (*Solanum tuberosum* L.) and their use efficiency under potato-legume intercropping systems. *Field Crops Research*, 222, 78–84. <https://doi.org/10.1016/j.fcr.2018.03.019>
- Glaze-Corcoran, S., Hashemi, M., Sadeghpour, A., Jahanzad, E., Keshavarz Afshar, R., Liu, X., & Herbert, S. J. (2020). Understanding intercropping to improve agricultural resiliency and environmental sustainability. In *Advances in Agronomy* (1st ed., Vol. 162, pp. 199–256). Elsevier Inc. <https://doi.org/10.1016/bs.agron.2020.02.004>
- Goh, C. H., Nicotra, A. B., & Mathesius, U. (2019). Genes controlling legume nodule numbers affect phenotypic plasticity responses to nitrogen in the presence and absence of rhizobia. *Plant Cell and Environment*, 42(5), 1747–1757. <https://doi.org/10.1111/pce.13498>
- Gommers, C. M. M., Visser, E. J. W., Onge, K. R. S., Voesenek, L. A. C. J., & Pierik, R. (2013). Shade tolerance: when growing tall is not an option. *Trends in Plant Science*, 18(2), 65–71. <https://doi.org/10.1016/j.tplants.2012.09.008>
- Gong, W. Z., Jiang, C. D., Wu, Y. S., Chen, H. H., Liu, W. Y., & Yang, W. Y. (2015). Tolerance vs. avoidance: two strategies of soybean (*Glycine max*) seedlings in response to shade in intercropping. *Photosynthetica*, 53(2), 259–268. <https://doi.org/10.1007/s11099-015-0103-8>
- Govindasamy, P., Muthusamy, S. K., Bagavathiannan, M., Mowrer, J., Jagannadham, P. T. K., Maity, A., Halli, H. M., G. K. S., Vadivel, R., T. K. D., Raj, R., Pooniya, V., Babu, S., Rathore, S. S., L. M., & Tiwari, G. (2023). Nitrogen use efficiency—a key to enhance crop productivity under a changing climate. *Frontiers in Plant Science*, 14(3), 1–19. <https://doi.org/10.3389/fpls.2023.1121073>
- Handriawan, A., Respatie, D. W., & Tohari. (2017). Pengaruh intensitas naungan terhadap pertumbuhan dan hasil tiga kultivar kedelai (*Glycine max* (L.) Merrill) di lahan pasir

- pantai Bugel, Kulon Progo [Effect of shade intensity on growth and yield of three soybean (*Glycine max* (L.) Merrill) cultivars on co. *Vegetalika*, 5(3), 1–14. <https://doi.org/10.22146/veg.25346>
- Hani, A. (2015). Produktivitas Kedelai pada Pola Agroforestri Nyamplung (*Callophylum inophyllum*) di Lahan Berpasir Pangandaran, Jawa Barat. *Jurnal Silvikultur Tropika*, 06(2), 78–82.
- Hassan, A., Dresbøll, D. B., Rasmussen, C. R., Lyhne-Kjærbye, A., Nicolaisen, M. H., Stokholm, M. S., Lund, O. S., & Thorup-Kristensen, K. (2021). Root distribution in intercropping systems—a comparison of DNA based methods and visual distinction of roots. *Archives of Agronomy and Soil Science*, 67(1), 15–28. <https://doi.org/10.1080/03650340.2019.1675872>
- He, Z. Q., Zhuang, J., Tang, H. R., & Huang, Z. (2012). Different vegetables crops in response to allelopathic of hot pepper root exudates. *World Applied Sciences Journal*, 19(9), 1289–1294. <https://doi.org/10.5829/idosi.wasj.2012.19.09.1886>
- Herawati, A., Mujiyo, Herdiansyah, G., Syamsiyah, J., & Lestari, A. (2023). Automasi irigasi di lahan pasir pantai untuk meningkatkan produktivitas tanaman budidaya. *Jurnal Sains Dan Teknologi*, 12(3), 572–582. <https://doi.org/10.23887/jstundiksha.v12i3.56096>
- Herlina, N., Hariyono, D., & Tri Margawati, D. (2017). Pengaruh Waktu Tanam Kubis (*Brassica oleraceae* L. var *capitata*) dan Cabai (*Capsicum annum* L.) Terhadap Efisiensi Penggunaan Lahan Pada Sistem Tumpangsari. *Jurnal Hortikultura Indonesia*, 8(2), 111. <https://doi.org/10.29244/jhi.8.2.111-119>
- Ho, R., Mo, O., Ti, O., & Ra, A. (2022). Cowpea *Vigna unguiculata* (L.) walp and chilli pepper *Capsicum frutescens* (L.) intercropping as affected by planting pattern and insect pest management in the rain forest area of Nigeria. *Journal of Entomology and Zoology Studies*, 10(2), 01–07. <https://doi.org/10.22271/j.ento.2022.v10.i2a.8960>
- Homulle, Z., George, T. S., & Karley, A. J. (2022). Root traits with team benefits: understanding belowground interactions in intercropping systems. *Plant and Soil*, 471(1–2), 1–26. <https://doi.org/10.1007/s11104-021-05165-8>
- Houlton, B. Z., Almaraz, M., Aneja, V., Austin, A. T., Bai, E., Cassman, K. G., Compton, J. E., Davidson, E. A., Erisman, J. W., Galloway, J. N., Gu, B., Yao, G., Martinelli, L. A., Scow, K., Schlesinger, W. H., Tomich, T. P., Wang, C., & Zhang, X. (2019). A World of Cobenefits: Solving the Global Nitrogen Challenge. *Earth's Future*, 7(8), 865–872. <https://doi.org/10.1029/2019EF001222>
- Ibrahim Abdel-Wahab, T. (2016). Response of Some Soybean Cultivars to Low Light Intensity under Different Intercropping Patterns with Maize. *International Journal of Applied Agricultural Sciences*, 2(2), 21. <https://doi.org/10.11648/j.ijaas.20160202.11>
- Indradewa, D. (2021). Potensi lahan pasir pantai sebagai lahan pertanian produktif [Potential of coastal sandy land as productive agricultural land]. In D. Indradewa, S. N. H. Utami, V. D. S. Handayani, & S. N. R. Irwan (Eds.), *Inovasi teknologi agronomi di lahan pasir pantai [Innovations in agronomic technology on coastal sandy land]* (pp. 1–30). Deepublish.
- Iqbal, N., Fatma, M., Gautam, H., Sehar, Z., Rasheed, F., Khan, M. I. R., Sofo, A., & Khan, N. A. (2022). Salicylic Acid Increases Photosynthesis of Drought Grown Mustard Plants Effectively with Sufficient-N via Regulation of Ethylene, Abscisic Acid, and Nitrogen-Use Efficiency. *Journal of Plant Growth Regulation*, 41(5), 1966–1977. <https://doi.org/10.1007/s00344-021-10565-2>

- Istiyanti, E., Khasanah, U., & Anjarwati, A. (2015). Pengembangan usahatani cabai merah di lahan pasir pantai kecamatan Temon Kabupaten Kulonprogo [Development of red chilli farming in coastal land of Temon, Kulonprogo]. *Agraris: Journal of Agribusiness and Rural Development Research*, 1(1), 6–11. <https://doi.org/10.18196/agr.112>
- Jensen, E. S., Carlsson, G., & Hauggaard-Nielsen, H. (2020). Intercropping of grain legumes and cereals improves the use of soil N resources and reduces the requirement for synthetic fertilizer N: a global-scale analysis. *Agronomy for Sustainable Development*, 40, 5. <https://doi.org/10.1007/s13593-020-0607-x>
- Jensen, E. S., Chongtham, I. R., Dhamala, N. R., Rodriguez, C., Carton, N., & Carlsson, G. (2020). Diversifying European agricultural systems by intercropping. *International Journal of Agriculture and Natural Resources*, 47(3), 174–186. <https://doi.org/10.7764/ijanr.v47i3.2241>
- Jumrani, K., Bhatia, V. S., Hussain, S., Kataria, S., Yang, X., & Brestic, M. (2024). Effect of shading on leaf anatomical structure, photosynthesis characteristics and chlorophyll fluorescence of soybean (*Glycine max*). *Journal of Agronomy and Crop Science*, 210(e12783). <https://doi.org/10.1111/jac.12783>
- Kebede, E. (2021). Contribution, Utilization, and Improvement of Legumes-Driven Biological Nitrogen Fixation in Agricultural Systems. *Frontiers in Sustainable Food Systems*, 5(November), 1–18. <https://doi.org/10.3389/fsufs.2021.767998>
- Kertonegoro, B. D. (2001). Gumuk Pasir Pantai di D.I. Yogyakarta: Potensi dan Pemanfaatannya Untuk Pertanian Berkelanjutan. *Prosiding Seminar Nasional Pemanfaatan Sumberdaya Lokal Untuk Pembangunan Pertanian Berkelanjutan*, 46–54.
- Khalid, M. H. B., Raza, M. A., Yu, H. Q., Sun, F. A., Zhang, Y. Y., Lu, F. Z., Si, L., Iqbal, N., Khan, I., Fu, F. L., & Li, W. C. (2019). Effect of shade treatments on morphology, photosynthetic and chlorophyll fluorescence characteristics of soybeans (*Glycine max* L. Merr.). *Applied Ecology and Environmental Research*, 17(2), 2551–2569. https://doi.org/10.15666/aeer/1702_25512569
- Kim, J., Song, Y., Kim, D. W., Fiaz, M., & Kwon, C. H. (2017). Enhancing Yield and Nutritive Value of Forage for Livestock Feeding Through Corn Soybean Intercropping Strategy with Several Pre-sowing Soybean Seed Coatings. *Journal of The Korean Society of Grassland and Forage Science*, 37(1), 50–55. <https://doi.org/10.5333/KGFS.2017.37.1.50>
- Klimek-Kopyra, A., Baciór, M., Lorenc-Kozik, A., Neugschwandtner, R. W., & Zajac, T. (2020). The Intraspecific competition as a driver for true production potential of soybean. *Italian Journal of Agronomy*, 16(1), 1709. <https://doi.org/10.4081/ija.2020.1709>
- Kosslak, R. M., & Bohlool, B. Ben. (1984). Suppression of Nodule Development of One Side of a Split-Root System of Soybeans Caused by Prior Inoculation of the Other Side. *Plant Physiology*, 75(1), 125–130. <https://doi.org/10.1104/pp.75.1.125>
- Larcher, W. (2003). Physiological Plant Ecology. In *Physiological Plant Ecology* (Fourth). Springer Berlin Heidelberg. <https://doi.org/10.1007/978-3-662-05214-3>
- Latati, M., Benlahrech, S., Lazali, M., Sihem, T., Kaci, G., Takouachet, R., Alkama, N., Hamdani, F. Z., Hafnaoui, E. A., Belarbi, B., Ounane, G., & Ounane, S. M. (2016). Intercropping Promotes the Ability of Legume and Cereal to Facilitate Phosphorus and Nitrogen Acquisition through Root- Induced Processes. In *Grain Legumes* (pp. 127–139). <https://doi.org/10.5772/63438>

- Lawson, T., & Matthews, J. (2020). Guard Cell Metabolism and Stomatal Function. *Annual Review of Plant Biology*, 71, 273–302. <https://doi.org/10.1146/annurev-arplant-050718-100251>
- Legba, E. C., Dossou, L., Honfoga, J., Pawera, L., & Srinivasan, R. (2025). Productivity and Profitability of Maize-Mungbean and Maize-Chili Pepper Relay Intercropping Systems for Income Diversification and Soil Fertility in Southern Benin. *Sustainability (Switzerland)*, 17(3). <https://doi.org/10.3390/su17031076>
- Li, C., Hoffland, E., Kuyper, T. W., Yu, Y., Zhang, C., Li, H., Zhang, F., & van der Werf, W. (2020). Syndromes of production in intercropping impact yield gains. *Nature Plants*, 6(6), 653–660. <https://doi.org/10.1038/s41477-020-0680-9>
- Li, L., Li, S. M., Sun, J. H., Zhou, L. L., Bao, X. G., Zhang, H. G., & Zhang, F. S. (2007). Diversity enhances agricultural productivity via rhizosphere phosphorus facilitation on phosphorus-deficient soils. *Proceedings of the National Academy of Sciences of the United States of America*, 104(27), 11192–11196. <https://doi.org/10.1073/pnas.0704591104>
- Li, L., Tilman, D., Lambers, H., & Zhang, F. S. (2014). Plant diversity and overyielding: Insights from belowground facilitation of intercropping in agriculture. *New Phytologist*, 203(1), 63–69. <https://doi.org/10.1111/nph.12778>
- Liang, Y., Liu, Q., Zeng, J., Xiong, F., Guo, J., Li, G., & Lu, D. (2024). Optimizing Nitrogen Input Increased Yield and Efficiency in Maize-Soybean Strip Intercropping System. *Agronomy*, 14(11). <https://doi.org/10.3390/agronomy14112472>
- Liao, C., Fang, S., Zhang, H., Liu, L., Xie, J., Li, X., Cao, F., Chen, J., & Huang, M. (2024). Grain Yield and Yield Attributes of Currently Popular Hybrid Rice Varieties Compared to Representative Super Hybrid Rice Varieties in Subtropical Environments. *Agronomy*, 14(318). <https://doi.org/10.3390/agronomy14020318>
- Lin, P., Wang, J., Chen, P., Fu, Z., Luo, K., Li, Y., Pu, T., Wang, X., Yong, T., & Yang, W. (2024). Relay intercropped soybean promotes nodules development and nitrogen fixation by root exudates deposition. *Frontiers in Plant Science*, 15. <https://doi.org/10.3389/fpls.2024.1447447>
- Liu, B., Gao, P., & Zhang, S. (2022). Photosynthesis and Fine Root Growth Dynamics of Soybean in Walnut-Soybean Agroforestry System. *Computational Intelligence and Neuroscience*, 2022(1), 1–8. <https://doi.org/10.1155/2022/2246824>
- Liu, C., Tu, B., Li, Y., Tian, B., Zhang, Q., Liu, X., & Herbert, S. J. (2017). Potassium application affects key enzyme activities of sucrose metabolism during seed filling in vegetable soybean. *Crop Science*, 57(5), 2707–2717. <https://doi.org/10.2135/cropsci2016.08.0648>
- Liu, L., Knight, J. D., Lemke, R. L., & Farrell, R. E. (2024). Quantifying the contribution of above- and below-ground residues of chickpea, faba bean, lentil, field pea and wheat to the nitrogen nutrition of a subsequent wheat crop. *Field Crops Research*, 313, 109412. <https://doi.org/10.1016/j.fcr.2024.109412>
- Liu, R., Zhou, G. peng, Cang, D. na, Gao, S. juan, Han, M., Zhang, J. dong, Sun, X. feng, & Cao, W. dong. (2022). Transfer characteristics of nitrogen fixed by leguminous green manure crops when intercropped with maize in northwestern China. *Journal of Integrative Agriculture*, 21(4), 1177–1187. [https://doi.org/10.1016/S2095-3119\(21\)63674-2](https://doi.org/10.1016/S2095-3119(21)63674-2)
- Liu, S., Wang, L., Chang, L., Khan, I., Nadeem, F., Rehman, A., & Suo, R. (2023). Evaluating the influence of straw mulching and intercropping on nitrogen uptake, crop

- growth, and yield performance in maize and soybean. *Frontiers in Plant Science*, *14*. <https://doi.org/10.3389/fpls.2023.1280382>
- Liu, X., Meng, L., Yin, T., Wang, X., Zhang, S., Cheng, Z., Ogundeji, A. O., & Li, S. (2023). Maize/soybean intercrop over time has higher yield stability relative to matched monoculture under different nitrogen-application rates. *Field Crops Research*, *301*, 109015. <https://doi.org/10.1016/j.fcr.2023.109015>
- Liu, X., Rahman, T., Song, C., Su, B., Yang, F., Yong, T., Wu, Y., Zhang, C., & Yang, W. (2017). Changes in light environment, morphology, growth and yield of soybean in maize-soybean intercropping systems. *Field Crops Research*, *200*, 38–46. <https://doi.org/10.1016/j.fcr.2016.10.003>
- Liu, X., Rahman, T., Yang, F., Song, C., Yong, T., Liu, J., Zhang, C., & Yang, W. (2017). PAR Interception and Utilization in Different Maize and Soybean Intercropping Patterns. *PLOS ONE*, *12*(1), e0169218. <https://doi.org/10.1371/journal.pone.0169218>
- Lo Presti, E., Kavamura, V. N., Abadie, M., Romeo, M., Reid, T. E., Heuer, S., Monti, M., & Mauchline, T. H. (2024). Phosphorus availability drives the effect of legume-wheat intercropping on prokaryotic community interactions. *Applied Soil Ecology*, *199*, 105414. <https://doi.org/10.1016/j.apsoil.2024.105414>
- Long, G., Li, L., Wang, D., Zhao, P., Tang, L., Zhou, Y., & Yin, X. (2021). Nitrogen levels regulate intercropping-related mitigation of potential nitrate leaching. *Agriculture, Ecosystems & Environment*, *319*, 107540. <https://doi.org/10.1016/j.agee.2021.107540>
- Lopes, E. C. S., Rodrigues, W. P., Fraga, K. R., Filho, J. A. M., da Silva, J. R., de Assis-Gomes, M. M., Figueiredo, F. A. M. M. A., Gresshoff, P. M., & Campostrini, E. (2019). Hypernodulating soybean mutant line nod4 lacking 'Autoregulation of Nodulation' (AON) has limited root-to-shoot water transport capacity. *Annals of Botany*, *124*(6), 979–991. <https://doi.org/10.1093/aob/mcz040>
- Luo, C., Guo, Z., Xiao, J., Dong, K., & Dong, Y. (2021). Effects of Applied Ratio of Nitrogen on the Light Environment in the Canopy and Growth, Development and Yield of Wheat When Intercropped. *Frontiers in Plant Science*, *12*, 1–12. <https://doi.org/10.3389/fpls.2021.719850>
- Lv, Q., Dai, J., Ding, K., He, N., Li, Z., Zhang, D., Xu, S., Li, C., Chi, B., Zhang, Y., & Dong, H. (2024). Managing interspecific competition to enhance productivity through selection of soybean varieties and sowing dates in a cotton-soybean intercropping system. *Field Crops Research*, *316*, 109513. <https://doi.org/10.1016/j.fcr.2024.109513>
- Mahmud, K., Makaju, S., Ibrahim, R., & Missaoui, A. (2020). Current progress in nitrogen fixing plants and microbiome research. *Plants*, *9*(1), 1–17. <https://doi.org/10.3390/plants9010097>
- Mahmud, K., Panday, D., Mergoum, A., & Missaoui, A. (2021). Nitrogen losses and potential mitigation strategies for a sustainable agroecosystem. *Sustainability (Switzerland)*, *13*(4), 1–23. <https://doi.org/10.3390/su13042400>
- Mao, L. L., Zhang, L. Z., Zhang, S. P., Evers, J. B., van der Werf, W., Wang, J. J., Sun, H. Q., Su, Z. C., & Spiertz, H. (2015). Resource use efficiency, ecological intensification and sustainability of intercropping systems. *Journal of Integrative Agriculture*, *14*(8), 1542–1550. [https://doi.org/10.1016/S2095-3119\(15\)61039-5](https://doi.org/10.1016/S2095-3119(15)61039-5)
- Marschner, P. (2012). Marschner's Mineral Nutrition of Higher Plants: Third Edition. In *Marschner's Mineral Nutrition of Higher Plants: Third Edition*. Elsevier Inc. <https://doi.org/10.1016/C2009-0-63043-9>

- Martínez-Dalmau, J., Berbel, J., & Ordóñez-Fernández, R. (2021). Nitrogen fertilization. A review of the risks associated with the inefficiency of its use and policy responses. *Sustainability (Switzerland)*, 13(10), 1–15. <https://doi.org/10.3390/su13105625>
- Matimati, I., Anthony Verboom, G., & Cramer, M. D. (2014). Do hydraulic redistribution and nocturnal transpiration facilitate nutrient acquisition in *Aspalathus linearis*? *Oecologia*, 175(4), 1129–1142. <https://doi.org/10.1007/s00442-014-2987-6>
- Mead, R., & Willey, R. W. (1980). The concept of a 'land equivalent ratio' and advantages in yields from intercropping. *Experimental Agriculture*, 16(3), 217–228. <https://doi.org/10.1017/S0014479700010978>
- Meinarti, N. S., & Jauhari, S. (2008). Penerapan Irigasi Mikro, Tumpangsari Dan Mulsa Untuk Mengantisipasi Kehilangan Hasil Cabai Merah Pada Penanaman Di Musim Kemarau. *Jurnal Agromet Indonesia*, 22(1), 13–21.
- Meitasari, A. D., & Wicaksono, K. P. (2017). Inokulasi rhizobium dan perimbangan nitrogen pada tanaman kedelai (*Glycine max* (L) merrill) varietas Wilis. *Plantropica Journal of Agricultural Science*, 2(1), 55–63.
- Mndzebele, B., Ncube, B., Fessehazion, M., Mabhaudhi, T., Amoo, S., du Plooy, C., Venter, S., & Modi, A. (2020). Effects of cowpea-amaranth intercropping and fertiliser application on soil phosphatase activities, available soil phosphorus, and crop growth response. *Agronomy*, 10(1). <https://doi.org/10.3390/agronomy10010079>
- Moekasan, T. K., & Prabaningrum, L. (2012). Penggunaan Rumah Kasa untuk Mengatasi Serangan Organisme Pengganggu Tumbuhan pada Tanaman Cabai Merah di Dataran Rendah. *Jurnal Hortikultura*, 22(1), 66. <https://doi.org/10.21082/jhort.v22n1.2012.p65-75>
- Mu, X., & Chen, Y. (2021). The physiological response of photosynthesis to nitrogen deficiency. *Plant Physiology and Biochemistry*, 158, 76–82. <https://doi.org/10.1016/j.plaphy.2020.11.019>
- Munawar. (2011). *Kesuburan Tanah dan Nutrisi Tanaman*. IPB Pres.
- Munawaroh, L., Kalsum, U., Laksono, P. B., & Siallagan, I. (2018). Respon Tanaman Kedelai Varietas Ceneng Pada Intensitas Cahaya Berbeda. *Jurnal Pertanian Presisi (Journal of Precision Agriculture)*, 2(2), 98–112. <https://doi.org/10.35760/jpp.2018.v2i2.2669>
- Munda, E. (2017). Effect of intercropping and phosphorous application on the growth and yield of sweetpotato, groundnut and soybean. In *Stellenbosch University*. Stellenbosch University.
- Nasar, J., Wang, G. Y., Zhou, F. J., Gitari, H., Zhou, X. B., Tabl, K. M., Hasan, M. E., Ali, H., Waqas, M. M., Ali, I., & Jahan, M. S. (2022). Nitrogen fertilization coupled with foliar application of iron and molybdenum improves shade tolerance of soybean under maize-soybean intercropping. *Frontiers in Plant Science*, 13. <https://doi.org/10.3389/fpls.2022.1014640>
- Nasar, J., Zhao, C. J., Khan, R., Gul, H., Gitari, H., Shao, Z., Abbas, G., Haider, I., Iqbal, Z., Ahmed, W., Rehman, R., Liang, Q. P., Zhou, X. B., & Yang, J. (2023). Maize-soybean intercropping at optimal N fertilization increases the N uptake, N yield and N use efficiency of maize crop by regulating the N assimilatory enzymes. *Frontiers in Plant Science*, 13. <https://doi.org/10.3389/fpls.2022.1077948>
- Ngwenya, Z. D., Mohammed, M., & Dakora, F. D. (2024). Monocropping and Intercropping of Maize with Six Food Legumes at Malkerns in Eswatini: Their Effects on Plant

- Growth, Grain Yield and N₂ Fixation, Measured using the ¹⁵N Natural Abundance and Ureide Techniques. *Symbiosis*, 92(2), 257–269. <https://doi.org/10.1007/s13199-024-00971-x>
- Ningrum, S. M., Tohari, T., & Respatie, D. W. (2020). Pengaruh Tingkat Naungan dan Takaran Pupuk Kandang Kambing Etawa terhadap Pertumbuhan dan Hasil Kedelai (*Glycine max* (L.) Merrill) di Lahan Pasir Pantai. *Vegetalika*, 9(2), 373. <https://doi.org/10.22146/veg.34876>
- Nugroho, N. C., & Caturatmi, A. (2017). Inovasi spesifik lokasi dalam pengembangan lahan pasir pantai sebagai lahan pertanian. *Prosiding Seminar Nasional Mewujudkan Kedaulatan Pangan Pada Lahan Sub Optimal Melalui Inovasi Teknologi Pertanian Spesifik Lokasi*, 1, 264–268.
- Ofori, F., & Stern, W. R. (1987). Cereal–Legume Intercropping Systems. *Advances in Agronomy*, 41, 41–90. [https://doi.org/10.1016/S0065-2113\(08\)60802-0](https://doi.org/10.1016/S0065-2113(08)60802-0)
- Omollo, J., Semu, E., Msaky, J., & Owuor, P. (2016). Effects of Lime and Nitrogen on Properties of an Acidic Soil and Nutrient Content of Sugarcane under Sugarcane – Soybean Intercropping in Kenya. *American Journal of Experimental Agriculture*, 13(6), 1–15. <https://doi.org/10.9734/AJEA/2016/28153>
- Partoyo. (2005). Analisis Indeks Kualitas Tanah Pertanian Di Lahan Pasir Pantai Samas Yogyakarta Analysis of Soil Quality Index for Sand Dune Agriculture Land At Samas Yogyakarta. *Agricultural Science (Ilmu Pertanian)*, 12(2), 140–151. [http://eprints.upnyk.ac.id/13510/1/Partoyo%2C 2005_Indeks Kualitas Tanah Samas_Ilmu Pertanian_Vol 12 No 2_140-151%281%29.pdf%0Ahttp://agrisci.ugm.ac.id/vol12_2.htm](http://eprints.upnyk.ac.id/13510/1/Partoyo%2C%202005_Indeks_Kualitas_Tanah_Samas_Ilmu_Pertanian_Vol_12_No_2_140-151%281%29.pdf%0Ahttp://agrisci.ugm.ac.id/vol12_2.htm)
- Pelzer, E., Hombert, N., Jeuffroy, M. H., & Makowski, D. (2014). Meta-analysis of the effect of nitrogen fertilization on annual cereal–legume intercrop production. *Agronomy Journal*, 106(5), 1775–1786. <https://doi.org/10.2134/agronj13.0590>
- Pemkab Bantul. (2021). *RPJMD Kabupaten Bantul 2021-2026*. Pemkab Bantul.
- Peng, X., Thevathasan, N. V, Gordon, A. M., Mohammed, I., & Gao, P. (2015). Photosynthetic Response of Soybean to Microclimate in 26-Year-Old Tree-Based Intercropping Systems in Southern Ontario, Canada. *PLOS ONE*, 10(6), e0129467. <https://doi.org/10.1371/journal.pone.0129467>
- Permanasari, I., Sulistyaningsih, E., Kurniasih, B., & Indradewa, D. (2021). Soybean varieties tolerance to intercropping with maize. *IOP Conference Series: Earth and Environmental Science*, 883(1), 012033. <https://doi.org/10.1088/1755-1315/883/1/012033>
- Permanasari, I., Sulistyaningsih, E., Kurniasih, B., & Indradewa, D. (2023). Morphophysiological and yield traits of soybean varieties tolerant of intercropping with maize. *Biodiversitas*, 24(7), 3872–3880. <https://doi.org/10.13057/biodiv/d240726>
- Pierozan Junior, C., Favarin, J. L., Lago, B. C., de Almeida, R. E. M., de Oliveira, S. M., Trivelin, P. C. O., Oliveira, F. B., & Gilabel, A. P. (2020). Nitrogen Fertilizer Recovery and Partitioning Related to Soybean Yield. *Journal of Soil Science and Plant Nutrition*, 20(4), 2566–2578. <https://doi.org/10.1007/s42729-020-00322-x>
- Poorter, H., Niklas, K. J., Reich, P. B., Oleksyn, J., Poot, P., & Mommer, L. (2012). Biomass allocation to leaves, stems and roots: Meta-analyses of interspecific variation and environmental control. *New Phytologist*, 193(1), 30–50. <https://doi.org/10.1111/j.1469-8137.2011.03952.x>

- Prabawardani, S., Puadi, L., Noya, A. I., & Syaranamual, S. (2021). Respon Pertumbuhan dan Hasil Jagung (*Zea mays* L.) Dalam Sistem Tumpangsari dengan Beberapa Jenis Tanaman Semusim. *Peningkatan Produktivitas Pertanian Era Society 5.0 Pasca Pandemi*, 121–132. <https://doi.org/10.25047/agropross.2021.214>
- Prasetyo, T., Subagiono, & Hasnelly. (2020). Produktivitas tanaman cabai rawit (*Capsicum Frutescens* L) dalam pola tumpangsari cabai dan jenis legum Triman. *Jurnal Sains Agro*, 9(2), 117–126.
- Pratiwi, H., & Artari, D. R. (2018). Respon Morfo-Fisiologi Genotipe Kedelai terhadap Naungan Jagung dan Ubikayu. *Jurnal Agronomi Indonesia (Indonesian Journal of Agronomy)*, 46(1), 48. <https://doi.org/10.24831/jai.v46i1.15441>
- Pratiwi, H., & Purwaningrahayu, R. D. (2021). Potensi Hasil Kedelai sebagai Tanaman Sela pada Pertanaman Cabai Rawit. *Proceedings Series on Physical & Formal Sciences*, 2, 124–130. <https://doi.org/10.30595/pspfs.v2i.180>
- Purnamasari, E. T., Yudono, P., & Rogomulyo, R. (2016). Pertumbuhan gulma dan hasil kedelai hitam (*Glycine max* (L.) Merrill) 'Mallika' terhadap pengolahan tanah dan jarak tanam di lahan pasir pantai [The weeds growth and yield of 'Mallika' black soybean (*Glycine max* (L.) Merrill) against to tillage and spac. *Vegetalika*, 5(1), 23–31. <https://doi.org/10.22146/VEG.24652>
- Pusdatin. (2024). *Analisis kinerja perdagangan kedelai*. Pusat Data dan Sistem Informasi Pertanian Kementerian Pertanian.
- Putra, F. P., Yudono, P., & Waluyo, S. (2018). Perubahan komposisi gulma pada sistem tumpangsari padi gogo dengan kedelai di lahan pasir pantai [Weeds composition change under intercropping system of upland rice with soybean in coastal sandy Area]. *Jurnal Agronomi Indonesia*, 46(1), 33–39. <https://doi.org/10.24831/jai.v46i1.17093>
- Putra, S. S., Putra, E. T. S., & Widada, J. (2020). The effects of types of manure and mycorrhizal applications on sandy soils on the growth and yield of curly red chili (*Capsicum annum* L.). *Caraka Tani: Journal of Sustainable Agriculture*, 35(2), 258–267. <https://doi.org/10.20961/carakatani.v35i2.34971>
- Putra, T. K., Afany, M. R., & Widodo, R. A. (2020). Pengaruh Bahan Organik Dan Tanah Vertisol Sebagai Pembena Tanah Terhadap Ketersediaan Dan Pelindian Kalium Di Tanah Regosol Pasir Pantai. *Tanah Dan Air*, 17(1), 20–25. <https://doi.org/10.31315/jta.v17i1.4007>
- Qin, H., Ma, Q., Chen, Y., Tian, B., Xu, L., Bai, Y., Chen, W., & Li, X. (2020). Variation in Rhizosphere Microbial Communities and Its Association With the Symbiotic Efficiency of Rhizobia in Soybean. *The Isme Journal*, 14(8), 1915–1928. <https://doi.org/10.1038/s41396-020-0648-9>
- Querejeta, J. I., Prieto, I., Armas, C., Casanoves, F., Diémé, J. S., Diouf, M., Yossi, H., Kaya, B., Pugnaire, F. I., & Rusch, G. M. (2022). Higher leaf nitrogen content is linked to tighter stomatal regulation of transpiration and more efficient water use across dryland trees. *New Phytologist*, 235(4), 1351–1364. <https://doi.org/10.1111/nph.18254>
- Rahman, T., Ye, L., Liu, X., Iqbal, N., Du, J., Gao, R., Liu, W., Yang, F., & Yang, W. (2017). Water use efficiency and water distribution response to different planting patterns in maize–soybean relay strip intercropping systems. *Experimental Agriculture*, 53(2), 159–177. <https://doi.org/10.1017/S0014479716000260>
- Rajiman. (2014). Pengaruh Bahan Pembena Tanah di Lahan Pasir Pantai Terhadap Kualitas Tanah. *Prosiding Seminar Nasional Lahan Suboptimal 2014*.

- Rasmussen, J., Gylfadóttir, T., Loges, R., Eriksen, J., & Helgadóttir, Á. (2013). Spatial and temporal variation in N transfer in grass-white clover mixtures at three Northern European field sites. *Soil Biology and Biochemistry*, *57*, 654–662. <https://doi.org/10.1016/j.soilbio.2012.07.004>
- Raza, A., Asghar, M. A., Ahmad, B., Bin, C., Iftikhar Hussain, M., Li, W., Iqbal, T., Yaseen, M., Shafiq, I., Yi, Z., Ahmad, I., Yang, W., & Weiguo, L. (2020). Agro-Techniques for Lodging Stress Management in Maize-Soybean Intercropping System—A Review. *Plants*, *9*(11), 1592. <https://doi.org/10.3390/plants9111592>
- Raza, M. A., Bin Khalid, M. H., Zhang, X., Feng, L. Y., Khan, I., Hassan, M. J., Ahmed, M., Ansar, M., Chen, Y. K., Fan, Y. F., Yang, F., & Yang, W. (2019). Effect of planting patterns on yield, nutrient accumulation and distribution in maize and soybean under relay intercropping systems. *Scientific Reports*, *9*, 4947. <https://doi.org/10.1038/s41598-019-41364-1>
- Raza, M. A., Din, A. M. U., Zhiqi, W., Gul, H., Ur Rehman, S., Bukhari, B., Haider, I., Rahman, M. H. U., Liang, X., Luo, S., El Sabagh, A., Qin, R., & Zhongming, M. (2023). Spatial differences influence nitrogen uptake, grain yield, and land-use advantage of wheat/soybean relay intercropping systems. *Scientific Reports*, *13*, 16916. <https://doi.org/10.1038/s41598-023-43288-3>
- Raza, M. A., Feng, L. Y., Iqbal, N., Ahmed, M., Chen, Y. K., Khalid, M. H. Bin, Mohi Ud Din, A., Khan, A., Ijaz, W., Hussain, A., Jamil, M. A., Naeem, M., Bhutto, S. H., Ansar, M., Yang, F., & Yang, W. (2019). Growth and development of soybean under changing light environments in relay intercropping system. *PeerJ*, *7*, e7262. <https://doi.org/10.7717/peerj.7262>
- Raza, M. A., Feng, L. Y., van der Werf, W., Cai, G. R., Khalid, M. H. Bin, Iqbal, N., Hassan, M. J., Meraj, T. A., Naeem, M., Khan, I., Rehman, S. ur, Ansar, M., Ahmed, M., Yang, F., & Yang, W. (2019). Narrow-Wide-Row Planting Pattern Increases The Radiation Use Efficiency and Seed Yield of Intercrop Species in Relay-Intercropping System. *Food and Energy Security*, *8*(3), 1–12. <https://doi.org/10.1002/fes3.170>
- Raza, M. A., Gul, H., Wang, J., Yasin, H. S., Qin, R., Bin Khalid, M. H., Naeem, M., Feng, L. Y., Iqbal, N., Gitari, H., Ahmad, S., Battaglia, M., Ansar, M., Yang, F., & Yang, W. (2021). Land productivity and water use efficiency of maize-soybean strip intercropping systems in semi-arid areas: A case study in Punjab Province, Pakistan. *Journal of Cleaner Production*, *308*, 127282. <https://doi.org/10.1016/j.jclepro.2021.127282>
- Richards, J. H., & Caldwell, M. M. (1987). Hydraulic lift: Substantial nocturnal water transport between soil layers by *Artemisia tridentata* roots. *Oecologia*, *73*(4), 486–489. <https://doi.org/10.1007/BF00379405>
- Richardson, A. E., Hocking, P. J., Simpson, R. J., & George, T. S. (2009). Plant mechanisms to optimise access to soil phosphorus. *Crop and Pasture Science*, *60*(2), 124–143. <https://doi.org/10.1071/CP07125>
- Rodriguez, C., Carlsson, G., Englund, J.-E., Flöhr, A., Pelzer, E., Jeuffroy, M.-H., Makowski, D., & Jensen, E. S. (2020). Grain legume-cereal intercropping enhances the use of soil-derived and biologically fixed nitrogen in temperate agroecosystems. A meta-analysis. *European Journal of Agronomy*, *118*, 126077. <https://doi.org/10.1016/j.eja.2020.126077>
- Römheld, V., & Kirkby, E. A. (2010). Research on potassium in agriculture: Needs and prospects. *Plant and Soil*, *335*(1), 155–180. <https://doi.org/10.1007/S11104-010-0520-1>

- Sai, E. J., . R., & Kumari, M. S. (2024). Maximizing Water Use Efficiency (WUE), Land Utilization Efficiency (LUE) and Competitive Indices of Maize and Soybean Crops Through Intercropping Pattern with Straw Mulching. *Indian Journal Of Agricultural Research*, 58(5), 800–805. <https://doi.org/10.18805/IJARE.A-6244>
- Sakoda, K., Watanabe, T., Sukemura, S., Kobayashi, S., Nagasaki, Y., Tanaka, Y., & Shiraiwa, T. (2019). Genetic Diversity in Stomatal Density among Soybeans Elucidated Using High-throughput Technique Based on an Algorithm for Object Detection. *Scientific Reports*, 9(1), 1–9. <https://doi.org/10.1038/s41598-019-44127-0>
- Salgado, G. C., Ambrosano, E. J., Rossi, F., Otsuk, I. P., Ambrosano, G. M. B., Santana, C. A., Muraoka, T., & Trivelin, P. C. O. (2021). Biological n fixation and n transfer in an intercropping system between legumes and organic cherry tomatoes in succession to green corn. *Agriculture (Switzerland)*, 11(8). <https://doi.org/10.3390/agriculture11080690>
- Salinas-Roco, S., Morales-González, A., Espinoza, S., Pérez-Díaz, R., Carrasco, B., del Pozo, A., & Cabeza, R. A. (2024). N₂ Fixation, N Transfer, and Land Equivalent Ratio (LER) in Grain Legume–Wheat Intercropping: Impact of N Supply and Plant Density. *Plants*, 13(7). <https://doi.org/10.3390/plants13070991>
- Salvagiotti, F., Cassman, K. G., Specht, J. E., Walters, D. T., Weiss, A., & Dobermann, A. (2008). Nitrogen uptake, fixation and response to fertilizer N in soybeans: A review. *Field Crops Research*, 108(1), 1–13. <https://doi.org/10.1016/j.fcr.2008.03.001>
- Sánchez-Navarro, V., Marcos-Pérez, M., Contreras, J., & Zornoza, R. (2024). Biological nitrogen fixation and intra- and interspecific competition in two vegetable/legume intercropping systems and their relationship to crop yield. *Scientia Horticulturae*, 337(113502). <https://doi.org/10.1016/j.scienta.2024.113502>
- Sapalina, F., Ginting, E. N., & Hidayat, F. (2022). Bakteri penambat nitrogen sebagai agen biofertilizer. *Warta Pusat Penelitian Kelapa Sawit*, 27(1), 41–50.
- Selosse, M. A., Richard, F., He, X., & Simard, S. W. (2006). Mycorrhizal networks: des liaisons dangereuses? *Trends in Ecology and Evolution*, 21(11), 621–628. <https://doi.org/10.1016/j.tree.2006.07.003>
- Setiawan, A. N., Isnawan, B. H., & Aini, L. N. (2015). *Sistem Pengelolaan Lahan Pasir Pantai Untuk Pengembangan Pertanian*.
- Setiawati, W., Muharam, A., Susanto, A., Boes, E., & Hudayya, A. (2019). Penerapan Teknologi Input Luar Rendah Pada Budidaya Cabai Merah untuk Mengurangi Penggunaan Pupuk dan Pestisida Sintetik (Implementation of Low External Input Technology for Chili Pepper Cultivation to Reduce Fertilizer and Synthetic Pesticide). *Jurnal Hortikultura*, 28(1), 113. <https://doi.org/10.21082/jhort.v28n1.2018.p113-122>
- Sevirasari, N., Sulistyarningsih, E., Kurniasih, B., Suryanti, S., Wibowo, A., & Joko, T. (2022). Effects of relay intercropping model and application of biological agents on the growth and yield of hot pepper. *Ilmu Pertanian (Agricultural Science)*, 7(1), 35. <https://doi.org/10.22146/ipas.69078>
- Shah, A., Nazari, M., Antar, M., Msimbira, L. A., Naamala, J., Lyu, D., Rabileh, M., Zajonc, J., & Smith, D. L. (2021). PGPR in Agriculture: A Sustainable Approach to Increasing Climate Change Resilience. *Frontiers in Sustainable Food Systems*, 5, 1–22. <https://doi.org/10.3389/fsufs.2021.667546>
- Shen, X., Liu, Y., Li, X., & Li, L. (2024). Effect of intercropping with legumes at different rates on the yield and soil physicochemical properties of *Cyperus esculentus* L. in arid land.

- Frontiers in Plant Science*, 15, 1–12. <https://doi.org/10.3389/fpls.2024.1351843>
- Shine, M. B., Gao, Q., Chowda-Reddy, R. V., Singh, A. K., Kachroo, P., & Kachroo, A. (2019). Glycerol-3-phosphate mediates rhizobia-induced systemic signaling in soybean. *Nature Communications*, 10(1), 5303. <https://doi.org/10.1038/s41467-019-13318-8>
- Sirait, M. H. A., & Karyawati, A. S. (2019). Pengaruh Naungan pada Pertumbuhan dan Hasil Beberapa Varietas Kedelai (*Glycine max* (L.) Merr). *Jurnal Produksi Tanaman*, 7(7), 1304–1313. <http://protan.studentjournal.ub.ac.id/index.php/protan/article/view/1179>
- Sitadz, S. A., & Kabirun, S. (2007). Pengembangan lahan marginal pesisir pantai dengan bioteknologi masukan rendah. *Ilmu Tanah Dan Lingkungan*, 7(2), 83–92.
- Somantri, R. U., Syahri, S., & Thamrin, T. (2019). Keragaan Agronomis dan Kelayakan Usahatani Kedelai yang Dibudidayakan Secara Monokultur dan Polikultur di Sumatera Selatan. *Jurnal Lahan Suboptimal: Journal of Suboptimal Lands*, 8(2), 159–172. <https://doi.org/10.33230/jlso.8.2.2019.426>
- Song, C., Wang, Q., Zhang, X., Sarpong, C. K., Wang, W., Yong, T., Wang, X., Wang, Y., & Yang, W. (2020). Crop Productivity and Nutrients Recovery in Maize–Soybean Additive Relay Intercropping Systems Under Subtropical Regions in Southwest China. *International Journal of Plant Production*, 14(2), 373–387. <https://doi.org/10.1007/s42106-020-00090-9>
- Soumare, A., Diedhiou, A. G., Thuita, M., Hafidi, M., Ouhdouch, Y., Gopalakrishnan, S., & Kouisni, L. (2020). Exploiting Biological Nitrogen Fixation: A Route Towards a Sustainable Agriculture. *Plants* 2020, Vol. 9, Page 1011, 9(8), 1011. <https://doi.org/10.3390/PLANTS9081011>
- Souza, J. a., Fratoni, M. M. J., Moraes, L. A. C., & Moreira, A. (2018). Boron and Amino Acid Foliar Application on Wheat-Soybean Intercropping in a Non-Tillage System. *Communications in Soil Science and Plant Analysis*, 49(13), 1638–1649. <https://doi.org/10.1080/00103624.2018.1474902>
- Su, B., Liu, X., Cui, L., Xiang, B., & Yang, W. (2018). Suppression of Weeds and Increases in Food Production in Higher Crop Diversity Planting Arrangements: A Case Study of Relay Intercropping. *Crop Science*, 58(4), 1729–1739. <https://doi.org/10.2135/cropsci2017.11.0670> LK
- Su, B. Y., Song, Y. X., Song, C., Cui, L., Yong, T. W., & Yang, W. Y. (2014). Growth and photosynthetic responses of soybean seedlings to maize shading in relay intercropping system in Southwest China. *Photosynthetica*, 52(3), 332–340. <https://doi.org/10.1007/s11099-014-0036-7>
- Su, Y., Yang, H., Wu, Y., Gong, W., Gul, H., Yan, Y., & Yang, W. (2023). Photosynthetic acclimation of shade-grown soybean seedlings to a high-light environment. *Plants*, 12, 2324. <https://doi.org/10.3390/plants12122324>
- Subagiono, S., Syarif, A., Syarif, Z., & Satria, B. (2019). Tumpangsari Berbasis Legum : a Review. *Jurnal Sains Agro*, 4(2), 1–7. <https://ojs.umb-bungo.ac.id/index.php/saingro/article/view/282/344>
- Subrata, B. A. G., Yudono, P., & Waluyo, S. (2016). Effect of Plant Density and Sowing Time on Growth and Yield of Mung Bean (*VignaRadiata* (L.) Wilczek) in Upland Rice-Mung Bean Intercropping on Land Sand Beach Samas Indonesia. *International Journal of Science and Research (IJSR)*, 5(8), 914–917. <https://doi.org/10.21275/v5i8.art2016986>

- Sudiono dan Purnomo. (2009). Hubungan Antara Populasi Kutu Kebul. *J. HPT Tropika*, 9(2), 115–120.
- Sukri, M. Z., Sari, V. K., & Firgiyanto, R. (2018). Improving Soil Fertilizer Through Application of Organic Fertilizer Humid Acid and Mikoriza in Supporting Growth and Production of Chilli Plants in Sand Land. *IOP Conference Series: Earth and Environmental Science*, 207(1). <https://doi.org/10.1088/1755-1315/207/1/012056>
- Sumarno, & Manshuri, A. G. (2013). Persyaratan tumbuh dan wilayah produksi kedelai di Indonesia [Requirements for soybean growth and production areas in Indonesia]. In Sumarno, Suyamto, A. Widjono, Hermanto, & H. Kasim (Eds.), *Kedelai: Teknik produksi dan pengembangan [Soybeans: Production techniques and development]* (pp. 74–103). Puslitbangtan.
- Sun, B., Wang, Y., Yang, Q., Gao, H., Niu, H., Li, Y., Ma, Q., Huan, Q., Qian, W., & Ren, B. (2023). A high-resolution transcriptomic atlas depicting nitrogen fixation and nodule development in soybean. *Journal of Integrative Plant Biology*, 65(6), 1536–1552. <https://doi.org/10.1111/jipb.13495>
- Sun, X., Wang, P., & Mi, G. (2025). *Genotypic Differences in Maize Root Morphology in Response to Low-Nitrogen Stress*.
- Sun, Y., Zang, H., Splettstößer, T., Kumar, A., Xu, X., Kuzyakov, Y., & Pausch, J. (2021). Plant intraspecific competition and growth stage alter carbon and nitrogen mineralization in the rhizosphere. *Plant Cell and Environment*, 44(4), 1231–1242. <https://doi.org/10.1111/pce.13945>
- Sundari, T., & Mutmaidah, D. S. (2018). Kesesuaian Galur-Galur Harapan Kedelai untuk Tumpangsari Jagung + Kedelai. *Jurnal Agronomi Indonesia (Indonesian Journal of Agronomy)*, 46(1), 40. <https://doi.org/10.24831/jai.v46i1.14880>
- Suriyagoda, L. D. B., Ryan, M. H., Renton, M., & Lambers, H. (2014). Plant Responses to Limited Moisture and Phosphorus Availability: A Meta-Analysis. *Advances in Agronomy*, 124, 143–200. <https://doi.org/10.1016/B978-0-12-800138-7.00004-8>
- Sutardi, & Astri, C. (2017). Sistem Usahatani Cabai Merah Pada Lahan Pasir Di Yogyakarta. *Pengkajian Dan Pengembangan Teknologi Pertanian*, 20(2), 125–140.
- Sutrisno, S., & Wijanarko, A. (2020). Perbedaan lokasi dan waktu tanam terhadap pertumbuhan dan hasil kedelai di lahan sawah Nusa Tenggara Barat [Effect of planting time and location on soybean growth and yield in lowland rice fields of West Nusa Tenggara]. *Agritech*, 21(2), 82–91. <https://doi.org/10.30595/agritech.v21i2.5579>
- Taiz, L., Zeiger, E., Møller, I. M., & Murphy, A. (2015). *Plant Physiology and Development* (6th ed.). Sinauer Associates Inc.
- Tang, Q. xiang, Tewolde, H., Liu, H. bin, Ren, T. zhi, Jiang, P. an, Zhai, L. mei, Lei, B. kun, Lin, T., & Liu, E. ke. (2018). Nitrogen Uptake and Transfer in Broad Bean and Garlic Strip Intercropping Systems. *Journal of Integrative Agriculture*, 17(1), 220–230. [https://doi.org/10.1016/S2095-3119\(17\)61772-6](https://doi.org/10.1016/S2095-3119(17)61772-6)
- Tanveer, M., Anjum, S. A., Hussain, S., Cerdà, A., & Ashraf, U. (2017). Relay cropping as a sustainable approach: problems and opportunities for sustainable crop production. *Environmental Science and Pollution Research*, 24(8), 6973–6988. <https://doi.org/10.1007/s11356-017-8371-4>
- Taufiq, A., & Sundari, T. (2012). Respon Tanaman Kedelai Terhadap Lingkungan Tumbuh. *Buletin Palawija*, 23, 13–26.
- Tcherkez, G., & Limami, A. M. (2019). Net photosynthetic CO₂ assimilation: more than just

- CO₂ and O₂ reduction cycles. *New Phytologist*, 223(2), 520–529. <https://doi.org/10.1111/nph.15828>
- Te, X., Din, A. M. U., Cui, K., Raza, M. A., Fraz Ali, M., & Xiao, J. (2023). Inter-specific root interactions and water use efficiency of maize/soybean relay strip intercropping. *Field Crops Research*, 291, 108793. <https://doi.org/10.1016/j.fcr.2022.108793>
- Telkar, S. G., Singh, A. K., & Dey, J. K. (2017). Effect of Population Proportion of Component Crops on Groth, Yield and Nutrient Uptake of Component Crops in Maize+Soybean Intercropping. *International Journal of Bio-Resource and Stress Management*, 8(6), 779–783. <https://doi.org/10.23910/IJBMSM/2017.8.6.3C0363>
- Teshome, S. (2019). A Review on Strategy of Developing Intercropping Practices. *International Journal of Current Research and Academic Review*, 7(1), 61–67. <https://doi.org/10.20546/ijcrar.2019.701.007>
- Thenveetil, N., Bheemanahalli, R., Reddy, K. N., Gao, W., & Reddy, K. R. (2024). Temperature and elevated CO₂ alter soybean seed yield and quality, exhibiting transgenerational effects on seedling emergence and vigor. *Frontiers in Plant Science*, 15(July), 1–18. <https://doi.org/10.3389/fpls.2024.1427086>
- Thilakarathna, M. S., McElroy, M. S., Chapagain, T., Papadopoulos, Y. A., & Raizada, M. N. (2016). Belowground nitrogen transfer from legumes to non-legumes under managed herbaceous cropping systems. A review. *Agronomy for Sustainable Development*, 36(4). <https://doi.org/10.1007/s13593-016-0396-4>
- Ton, A. (2021). Advantages of Grain Legume-Cereal Intercropping in Sustainable Agriculture. *Turkish Journal of Agriculture - Food Science and Technology*, 9(8), 1560–1566. <https://doi.org/10.24925/turjaf.v9i8.1560-1566.4481>
- Tsialtas, I. T., Baxevanos, D., Vlachostergios, D. N., Dordas, C., & Lithourgidis, A. (2018). Cultivar complementarity for symbiotic nitrogen fixation and water use efficiency in pea-oat intercrops and its effect on forage yield and quality. *Field Crops Research*, 226, 28–37. <https://doi.org/10.1016/j.fcr.2018.07.005>
- Van Der Heijden, M. G. A., & Horton, T. R. (2009). Socialism in soil? the importance of mycorrhizal fungal networks for facilitation in natural ecosystems. *Journal of Ecology*, 97(6), 1139–1150. <https://doi.org/10.1111/j.1365-2745.2009.01570.x>
- Van Rhijn, P., & Vanderleyden, J. (1995). The Rhizobium-plant symbiosis. *Microbiological Reviews*, 59(1), 124–142. <https://doi.org/10.1128/mmbr.59.1.124-142.1995>
- Vandenbussche, F., Pierik, R., Millenaar, F. F., Voeselek, L. A. C. J., & Van Der Straeten, D. (2005). Reaching out of the shade. *Current Opinion in Plant Biology*, 8(5), 462–468. <https://doi.org/10.1016/j.pbi.2005.07.007>
- von Caemmerer, S., & Farquhar, G. D. (1981). Some relationships between the biochemistry of photosynthesis and the gas exchange of leaves. *Planta*, 153(4), 376–387. <https://doi.org/10.1007/BF00384257>
- Vorobey, N., Kukol, K., Pukhtaievych, P., & Kots, T. (2022). Symbiotic and physiological indicators of soybean inoculated of Bradyrhizobium japonicum single-strain in 7 days before sowing. *Acta Agriculturae Slovenica*, 118(2), 1–11. <https://doi.org/10.14720/aas.2022.118.2.1867>
- Wang, Q., Li, S., Li, J., & Huang, D. (2024). The Utilization and Roles of Nitrogen in Plants. *Forests*, 15(7). <https://doi.org/10.3390/f15071191>
- Wang, Q., Sun, Z., Bai, W., Zhang, D., Zhang, Yu., Wang, R., Van Der Werf, W., Evers, Jochem, B., Stomph, T.-J., Guo, J., & Zhang, L. (2021). Light interception and use

- efficiency differ with maize plant density in maize-peanut intercropping. *Frontiers of Agricultural Science and Engineering*, 8(3), 432–446. <https://doi.org/10.15302/J-FASE-2021403>
- Wang, T. (2024). *Bacillus* Suppresses Nitrogen Efficiency of Soybean–rhizobium Symbiosis Through Regulation of Nitrogen-related Transcriptional and Microbial Patterns. *Plant Cell & Environment*, 47(11), 4305–4322. <https://doi.org/10.1111/pce.15023>
- Wang, X., Li, X., Zhou, Q., Song, S., & Dong, S. (2023). Comparison and Evaluation of Low-Temperature Tolerance of Different Soybean Cultivars during the Early-Growth Stage. In *Agronomy* (Vol. 13, Issue 7, p. 1716). <https://doi.org/10.3390/agronomy13071716> LK - <https://doi.org/10.3390/agronomy13071716>
- Wang, Y., Qin, Y., Chai, Q., Feng, F., Zhao, C., & Yu, A. (2018). Interspecies interactions in relation to root distribution across the rooting profile in wheat-maize intercropping under different plant densities. *Frontiers in Plant Science*, 9(April), 1–17. <https://doi.org/10.3389/fpls.2018.00483>
- Wen, B. xiao, Hussain, S., Yang, J. yue, Wang, S., Zhang, Y., Qin, S. si, Xu, M., Yang, W. yu, & Liu, W. guo. (2020). Rejuvenating soybean (*Glycine max* L.) growth and development through slight shading stress. *Journal of Integrative Agriculture*, 19(10), 2439–2450. [https://doi.org/10.1016/S2095-3119\(20\)63159-8](https://doi.org/10.1016/S2095-3119(20)63159-8)
- Widodo, A. S. (2015). Pendapatan dan Produksi Potensial Usahatani Konservasi Lahan Pantai di Kabupaten Bantul. *AGRARIS: Journal of Agribusiness and Rural Development Research*, 1(1), 1–5. <https://doi.org/10.18196/agr.111>
- Wu, Y., Gong, W., Yang, F., Wang, X., Yong, T., Liu, J., Pu, T., Yan, Y., & Yang, W. (2022). Dynamic of recovery growth of intercropped soybean after maize harvest in maize–soybean relay strip intercropping system. *Food and Energy Security*, 11(1), 1–14. <https://doi.org/10.1002/fes3.350>
- Wysokinski, A., Wysokińska, A., Noulas, C., & Wysokińska, A. (2024). Optimal Nitrogen Fertilizer Rates for Soybean Cultivation. *Agronomy*, 14(7), 1–12. <https://doi.org/10.3390/agronomy14071375>
- Xia, H. Y., Zhao, J. H., Sun, J. H., Bao, X. G., Christie, P., Zhang, F. S., & Li, L. (2013). Dynamics of root length and distribution and shoot biomass of maize as affected by intercropping with different companion crops and phosphorus application rates. *Field Crops Research*, 150, 52–62. <https://doi.org/10.1016/j.fcr.2013.05.027>
- Xiao, J., Yin, X., Ren, J., Zhang, M., Tang, L., & Zheng, Y. (2018). Complementation drives higher growth rate and yield of wheat and saves nitrogen fertilizer in wheat and faba bean intercropping. *Field Crops Research*, 221, 119–129. <https://doi.org/10.1016/j.fcr.2017.12.009>
- Xiao, Y., Yang, D., Zhang, S.-B., Mo, Y.-X., Dong, Y.-Y., Wang, K.-F., He, L.-Y., Dong, B., Dossa, G. G. O., & Zhang, J.-L. (2024). Nitrogen-fixing and non-nitrogen-fixing legume plants differ in leaf nutrient concentrations and relationships between photosynthetic and hydraulic traits. *Tree Physiology*, 44(5), tpae048. <https://doi.org/10.1093/treephys/tpae048>
- Yang, F., Feng, L., Liu, Q., Wu, X., Fan, Y., Raza, M. A., Cheng, Y., Chen, J., Wang, X., Yong, T., Liu, W., Liu, J., Du, J., Shu, K., & Yang, W. (2018). Effect of interactions between light intensity and red-to- far-red ratio on the photosynthesis of soybean leaves under shade condition. *Environmental and Experimental Botany*, 150(February), 79–87. <https://doi.org/10.1016/j.envexpbot.2018.03.008>

- Yang, F., Liao, D., Wu, X., Gao, R., Fan, Y., Raza, M. A., Wang, X., Yong, T., Liu, W., Liu, J., Du, J., Shu, K., & Yang, W. (2017). Effect of aboveground and belowground interactions on the intercrop yields in maize-soybean relay intercropping systems. *Field Crops Research*, 203, 16–23. <https://doi.org/10.1016/j.fcr.2016.12.007>
- Yin, W., Chai, Q., Zhao, C., Yu, A., Fan, Z., Hu, F., Fan, H., Guo, Y., & Coulter, J. A. (2020). Water utilization in intercropping: A review. *Agricultural Water Management*, 241, 106335. <https://doi.org/10.1016/j.agwat.2020.106335>
- Yong, T. wen, Chen, P., Dong, Q., Du, Q., Yang, F., Wang, X. chun, Liu, W. guo, & Yang, W. yu. (2018). Optimized nitrogen application methods to improve nitrogen use efficiency and nodule nitrogen fixation in a maize-soybean relay intercropping system. *Journal of Integrative Agriculture*, 17(3), 664–676. [https://doi.org/10.1016/S2095-3119\(17\)61836-7](https://doi.org/10.1016/S2095-3119(17)61836-7)
- Yu, Y., Stomph, T. J., Makowski, D., & van der Werf, W. (2015). Temporal niche differentiation increases the land equivalent ratio of annual intercrops: A meta-analysis. *Field Crops Research*, 184, 133–144. <https://doi.org/10.1016/j.fcr.2015.09.010>
- Zhang, F., & Li, L. (2003). Using competitive and facilitative interactions in intercropping systems enhances crop productivity and nutrient-use efficiency. *Plant and Soil*, 248(1–2), 305–312. <https://doi.org/10.1023/A:1022352229863>
- Zhang, G., Hou, Y., Zhang, H., Fan, H., Wen, X., Han, J., & Liao, Y. (2022). Optimizing planting pattern and nitrogen application rate improves grain yield and water use efficiency for rain-fed spring maize by promoting root growth and reducing redundant root growth. *Soil and Tillage Research*, 220, 105385. <https://doi.org/10.1016/j.still.2022.105385>
- Zhang, H., Zeng, F., Zou, Z., Zhang, Z., & Li, Y. (2017). Nitrogen uptake and transfer in a soybean/maize intercropping system in the karst region of southwest China. In *Ecology and Evolution* (Vol. 7, Issue 20, pp. 8419–8426). <https://doi.org/10.1002/ece3.3295> LK - <https://doi.org/10.1002/ece3.3295>
- Zhang, M. cong, Sun, W. xiang, Liu, Y. ying, Luo, S. guo, Zhao, J., Wu, Q., Wu, Z. yu, & Jiang, Y. (2014). Timing of N Application Affects Net Primary Production of Soybean with Different Planting Densities. *Journal of Integrative Agriculture*, 13(12), 2778–2787. [https://doi.org/10.1016/S2095-3119\(13\)60535-3](https://doi.org/10.1016/S2095-3119(13)60535-3)
- Zhang, R., Wang, C., Teng, W., Wang, J., Lyu, X., Dong, S., Kang, S., Gong, Z., & Ma, C. (2020). Accumulation and distribution of fertilizer nitrogen and nodule-fixed nitrogen in soybeans with dual root systems. *Agronomy*, 10(3). <https://doi.org/10.3390/agronomy10030397>
- Zhang, T., Tang, H., Peng, P., Ge, S., Liu, Y., Feng, Y., & Wang, J. (2023). Sugarcane/soybean intercropping with reduced nitrogen addition promotes photosynthesized carbon sequestration in the soil. *Frontiers in Plant Science*, 14(December), 1–14. <https://doi.org/10.3389/fpls.2023.1282083>
- Zhang, W. F., Dou, Z. X., He, P., Ju, X. T., Powlson, D., Chadwick, D., Norse, D., Lu, Y. L., Zhang, Y., Wu, L., Chen, X. P., Cassman, K. G., & Zhang, F. S. (2013). New technologies reduce greenhouse gas emissions from nitrogenous fertilizer in China. *Proceedings of the National Academy of Sciences of the United States of America*, 110(21), 8375–8380. <https://doi.org/10.1073/pnas.1210447110>
- Zhang, Y., Sun, Z., Su, Z., Du, G., Bai, W., Wang, Q., Wang, R., Nie, J., Sun, T., Feng, C., Zhang, Z., Yang, N., Zhang, X., Evers, J. B., van der Werf, W., & Zhang, L. (2022).

- Root plasticity and interspecific complementarity improve yields and water use efficiency of maize/soybean intercropping in a water-limited condition. *Field Crops Research*, 282, 108523. <https://doi.org/10.1016/j.fcr.2022.108523>
- Zhang, Z., Yu, Z., Zhang, Y., & Shi, Y. (2022). Impacts of fertilization optimization on soil nitrogen cycling and wheat nitrogen utilization under water-saving irrigation. *Frontiers in Plant Science*, 13, 1–13. <https://doi.org/10.3389/fpls.2022.878424>
- Zhao, Y., Tian, Y., Li, X., Song, M., Fang, X., Jiang, Y., & Xu, X. (2022). Nitrogen fixation and transfer between legumes and cereals under various cropping regimes. *Rhizosphere*, 22(May), 100546. <https://doi.org/10.1016/j.rhisph.2022.100546>
- Zhao, Z., Li, Z., Li, Y., Yu, L., Gu, X., & Cai, H. (2024). Supplementary irrigation and reduced nitrogen application improve the productivity, water and nitrogen use efficiency of maize-soybean intercropping system in the semi-humid drought-prone region of China. *Agricultural Water Management*, 305, 109126. <https://doi.org/10.1016/j.agwat.2024.109126>
- Zheng, B. chuan, Zhou, Y., Chen, P., Zhang, X. na, Du, Q., Yang, H., Wang, X. chun, Yang, F., Xiao, T., Li, L., Yang, W. yu, & Yong, T. wen. (2022). Maize–legume intercropping promote N uptake through changing the root spatial distribution, legume nodulation capacity, and soil N availability. *Journal of Integrative Agriculture*, 21(6), 1755–1771. [https://doi.org/10.1016/S2095-3119\(21\)63730-9](https://doi.org/10.1016/S2095-3119(21)63730-9)
- Zhou, J., Beche, E., Vieira, C. C., Yungbluth, D., Zhou, J., Scaboo, A., & Chen, P. (2022). Improve Soybean Variety Selection Accuracy Using UAV-Based High-Throughput Phenotyping Technology. *Frontiers in Plant Science*, 12. <https://doi.org/10.3389/fpls.2021.768742>