



## DAFTAR PUSTAKA

- Akinbomi, J. and Taherzadeh, M. J. (2015) 'Evaluation of Fermentative Hydrogen Production from Single and Mixed Fruit Wastes', 504, pp. 4253–4272. doi: 10.3390/en8054253.
- Aguilar R.M.A., Guelfo, F., Gallego, A., Garcia, R. (2012) " Effect of HRT on hydrogen production and organic matter solubilization in acidogenic anaerobic digestion of OFMSW", Chemical Engineering Journal, 219, 443-449.
- Ali S, Khan, Ahmed, Musaddiq, Khaja S Ahmed, H Polasa, Venkateswar Rao, Habibullah, Sechi, and Ahmed (2005), "Antimicrobial activities of Eugenol and Cinnamaldehyde against the human gastric pathogen *Helicobacter pylori*. *Ann Clin Microbiol Antimicrob*
- Ameekan, Y. (2014)"Pengaruh Perbedaan Asal Inokulum Terhadap Produksi Hidrogen Dari Limbah Buah Melon (Cucumis melo. L") pada fermentor batch", Skripsi : Universitas Gadjah mada.M
- Anonim (2017) "Indonesia - Finlandia bahas kerjasama pengelolaan sampah menjadi energi". Available at : <http://www.menlhk.go.id/berita-189-indonesia-finlandia-bahas-kerjasama-pengelolaan-sampah-menjadi-energi.html> (Accessed: 10 juli 2017)
- Anonim (2017) "Basic Report : Nutrient data for 09181, Melons, cantaloupe, raw". Available at : <https://ndb.nal.usda.gov/ndb/foods/show/2274> (Accessed; 21 juli 2017)
- Aubert C and Pitrat M.(2006)"Volatile compounds in the skin and pulp of Queen Anne's pocket melon", J. Agric. Food Chem. 54(21) 8177–8182
- Balachandar, G., Khanna, N. and Das, D. (2013) "Biohydrogen Production from Organic Wastes by Dark Fermentation", 1st edn, Biohydrogen. 1st edn. Elsevier B.V. doi: 10.1016/B978-0-444-59555-3.00006-4.
- Baghchehsaraee, B. (2009)"Batch and Continuous Biohydrogen Production Using Mixed Microbial Culture", Dissertation, Engineering Science, Department of Chemical and Biochemical Engineering, The University of Western Ontario, Canada
- Beaulieu, J. C. et al. (2003) 'Sugar and organic acid variations in commercial cantaloupes and their inbred parents', *Journal of the American Society for Horticultural Science*, 128(4), pp. 531–536.
- Cahyari K, Hidayat M, Sarto, Syamsiah Siti, (2013)"Effect of Initial Substrate Concentration on Biohydrogen Production from Orange and Melon Fruit Waste through Dark Fermentation", 6<sup>th</sup> regional Conf on Chemical Eng, Manila Philipines,



Cahyari K, Sarto, Syamsiah S, Prasetya A.(2017) "Performance of continuous stirred tank reactor (CSTR) on fermentative biohydrogen production from melon waste", 2<sup>nd</sup> Int conf of Chem Eng, IOP Conf. Series: Materials Science and Engineering 162 (2017) 012013 doi:10.1088/1757-899X/162/1/012013

Catherine, A., Deepika, H., and Negi.P. (2012) "Antibacterial activity of eugenol and peppermint oil in model food systems", Journal of Essential Oil Research Volume 24, 2012, pp. 481-486,

Das, D. and Veziroglu, T. N.(2001)"Hydrogen Production by Biological Processes: A Survey of Literature", Int. J. Hydrogen Energy, 26, 13-28.

Das, D., Khanna, N. and Veziroglu, T. N. (2008) 'Recent developments in biological sequence databases.', Current opinion in biotechnology, 9(2), pp. 54–58. doi: 10.2298/CICEQ0802057D

Fan, Y., Li, C., Lay J.J., Hou, H., Zhang, G. (2004) " Optimization of initial susbtrate and pH level for germination of sporing hydrogen-producing anaerobes in cow dung compost", Biosource Tecnology, 91, 189-193.

Fang, H.H.P., Liu, H. (2002),"Effect of pH on hydrogen production from glucose by a mixed culture", Bioresour. Technol.82, 87–93.

Fasogbon, S.K. (2015), "Melon oil methyl ester : An environmentatlly friendly fuel", Journal of Natural resources and Development 05 : 47-53

Ghimire, A. *et al.* (2015) 'A review on dark fermentative biohydrogen production from organic biomass : Process parameters and use of by-products', *Applied Energy*. Elsevier Ltd, 144, pp. 73–95. doi: 10.1016/j.apenergy.2015.01.045.

Gustavsson J, Cederberg C, Sonesson U, Van Otterdijk R and Meybeck A.(2011) Global food losses and food waste *Food Agric. Organ. U. N. Rom*

Hallenbeck, P.C., Benemann, J.R. (2002), Biological hydrogen production; fundamentals and limiting processes. Int. J. Hydrogen Energy 27, 1185–1193.

Hawkes, F. R., Hussy, I., Kyazze, G., Dinsdale, R. and Hawkes, D. L.(2007), "Continuous Dark Fermentative Hydrogen Production by Mesophilic Microfolora: Principles and Progress", Int. J. Hydrogen Energy, 32 (2), 172-84

Kapdan, I.K., and Kargi, F. (2006), "Bio-hydrogen Production from Waste Materials", Enzyme and Microbial Technology, vol 38, 569–582



- Khanal, S.K, Chen, W.H, Li, L., and Sung, S. (2004) "Biological hydrogen production : Effect of pH and intermediate product", Int. Journal hydrogen energy, 29(11), 1123-1131.
- Kim, D.H., Kim, S. H., Jung, K. W., Kim, M. S., Shin, H.S. , (2011), "Effect of initial pH independent of operational pH on hydrogen fermentation of food waste". Biosour. Technol, 102, 8646-8652
- Kim SH, Han SK, Shin HS. (2008), "Optimization of continuous hydrogen fermentation of food waste as a function of solids retention time independent of hydraulic retention time, Process Biochemistry, 43 pp. 213-218.
- Kim, M.-S., Cha, J. and Kim, D.-H. (2013) Fermentative Biohydrogen Production from Solid Wastes. 1st edn, Biohydrogen. 1st edn. Elsevier B.V. doi: 10.1016/B978-0-444-59555-3.00011-8.
- Kim, J.K., Oh, B.R., Chun, Y.N., Kim, S.W. (2006) " Effect of temperature and hydraulic retention time on anaerobic digestion of food waste, J. Bioscience and bioengineering, 102, 328-332.
- Koskinen, A.H. Kaksonen, J.A. Puhakka, (2007, "The relationship between instability of H<sub>2</sub> production and compositions of bacterial communities within a dark fermentation fluidized-bed bioreactor", Biotechnol. Bioeng. 97 742–758, doi:<http://dx.doi.org/10.1002/bit.21299>.
- Levin, D. B., Pitt, L. and Love, M. (2004) 'Biohydrogen production: Prospects and limitations to practical application', International Journal of Hydrogen Energy, 29(2), pp. 173–185. doi: 10.1016/S0360-3199(03)00094-6.
- Lo, Y.C., Su, Y.C., Chen, C.Y., Chen, Y.M., Lee, K.S., Chang, J.S (2009) "Biohydrogen production from cellulosic hydrolysate produced via temperature-shift-enhanced bacterial cellulose hydrolysis. Biosource Technology, 100(23), 5802-5807.
- Logan, B. E. (2004)"Extracting Hydrogen and Electricity from Renewable Resources", Environmental Sci. Tech., 38 (9), 160-7.
- Maier, R.M. (2009), " Bacterial Growth", Environmental Microbiology, pp 37-54
- Mohan, S. V. and Pandey, A. (2013) 'Biohydrogen Production: An Introduction', Biohydrogen, pp. 1–24. doi: 10.1016/B978-0-444-59555-3.00001-5.
- Moon, C., et al. (2005) "Effect of the accuracy of pH control on hydrogen fermentation", Biosource Technology, 179, 595-601.
- Mu, Y., Wang, G., and Yu, H. Q., (2006), "kinetic modelling of batch hydrogen production process by mixed anaerobic cultures", Biosource Technology, 97 (11), 1302-1307



Mullai, P., Sridevi, K., (2014), "Cell growth and product formation kinetic of biohydrogen production using mixed consortia by batch process", Int. J. of Chem Tech Research, 6 (12), 5125-5130

Nurkholis (2017) "Pengaruh organic loading rate (olr) dan frekuensi pengumpanan pada produksi biohidrogen dari sampah buah melon (cucumis melo l.) dengan pengumpanan secara berkala (periodic feeding) menggunakan reaktor alir pipa", Tesis, Universitas Gadjah Mada, Yogyakarta

Nurrihadini, O. D. (2009) *Karakterisasi Sampah Pasar Buah Gemah Ripah Yogyakarta sebagai Bahan Baku Produksi Biogas*, Skripsi di Jurusan Teknologi Pangan dan Hasil Pertanian Fakultas Teknologi Pertanian Universitas Gadjah Mada, Yogyakarta.

Oh, K.Y., Kim, H.S., Kim, S.M., Park, S., (2004) "Thermophilic Biohydrogen production from glucose with trickling biofilter", Biotechnology and bioengineering, 88 (6), 690-698.

Pachauri, R.K and Reisinger, A. (2008) "PCC, 2007: Climate Change 2007: Synthesis Report. Contribution of Working Groups I, II and III to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change (eds.)]", IPCC, Geneva, Switzerland

Ren, N., Wang, A., Cao, G., Xu, J., Gao, L. (2009) Bioconversion of lignocellulosic biomass to hydrogen: Potential and challenges. Biotechnol. Adv. 27, 1051–1060.

Ruggeri et al. (2013) "Biohydrogen & Biomethane Through Anaerobic Digestion", Green Energy and Technology, DOI 10.1007/978-1-4471-6431-9\_225

Saratale, G. D., Saratale, R. G. and Chang, J. S. (2013) 'Biohydrogen from Renewable Resources', *Biohydrogen*, pp. 185–221. doi: 10.1016/B978-0-444-59555-3.00009-X.

Shah B, Davidson P M and Zhong Q (2013)"Nanodispersed eugenol has improved antimicrobial activity against Escherichia coli O157: H7 and Listeria monocytogenes in bovine milk" Int. J. Food Microbiol. 161(1) 53–59

Shin, H.S., Youn, J.H., Kim, S.H. (2004)"Hydrogen production from food waste in anaerobic mesophilic and thermophilic acidogenesis", Int. J. Hydrogen Energy 29, 1355–1363.

Shuller, M and Kargi,F. (2002) " Bioprocess Engineering Basic Concept 2<sup>nd</sup> Ed.", Prentice Hall, New Jersey

Stepansky, A. et al. (1999) 'Variation in sugar levels and invertase activity in mature fruit representing a broad spectrum of Cucumis melo genotypes', Genetic Resources and Crop Evolution, 46(1), pp. 53–62. doi: 10.1023/A:1008636732481.



- Swinnen, I. A.M., Bernaerts, K., Dens, E.J.J., Geereard, A.H., van Impe, JF., (2004) " Predictive modelling of the microbial lag phase : a view", Int. J. Food Microbiol. 94, 137-159
- Vijayaraghavan, K., Ahmad, D. and Soning, C. (2007) 'Bio-hydrogen generation from mixed fruit peel waste using anaerobic contact filter Bio-hydrogen generation from mixed fruit peel waste using anaerobic contact filter', (April). doi: 10.1016/j.ijhydene.2007.07.001.
- Wang, J., Wan, W. (2009) Factors influencing fermentative hydrogen production: A review. Int. J. Hydrogen Energy. 34, 799–811.
- Widhyasih, Kunti (2016) "Evaluasi waktu start up dan pengaruh Hydraulic Retention Time pada proses peruraian stillage secara anaerobik menggunakan reaktor fluidized bed kontinyu dengan zeolit sebagai media immobilisasi", Tesis, Universitas Gadjah Mada, Yogyakarta
- Zealand, A.M., Roskilly, A.P. and Graham, D.W. (2017) " Effect of feeding frequency and organic loading rate on beomethane production in the anaerobic digestion of rice straw", Applied Energy, the authors, 207, 156-165
- Zhang, M.L., Y.T. Fan, Y.Y. Xing, C.M. Pan, G.S. Zhang and J.J. Lay (2007) "Enhanced biohydrogen production from cornstalk wastes with acidification pretreatment by mixed anaerobic cultures", Biom. Bioen., 31: 250-254
- Zhang, B., Zhang, L.L., Zhang, S.C., Shi, H.Z., Cai, W.M. (2005) "The influence of pH on hydrolysis and acidogenesis of kitchen wastes in two-phase anaerobic digestion", Environ. Technol. 26, 329–339.