

Daftar Pustaka

- Amin, M.B., Greene, F.L., Edge, S.B., Compton, C.C., Gershenwald, J.E., Brookland, R.K., *et al.* (2017). The Eighth Edition AJCC Cancer Staging Manual: Continuing to build a bridge from a population-based to a more “personalized” approach to cancer staging. *CA Cancer J Clin* 67 : 93–99.
- Arroyo, V., García-Martinez, R., & Salvatella, X. (2014). Human serum albumin, systemic inflammation, and cirrhosis. *Journal of Hepatology* 61 : 396–407.
- Brkic, F.F., Kadletz, L., Jank, B., Mayer, C., Heiduschka, G., & Brunner, M. (2020). Impact of pretherapeutic neutrophil-to-lymphocyte ratio, serum albumin, body-mass index, and advanced lung cancer inflammation index on clinical outcome in sinonasal squamous cell carcinoma. *Journal of Cranio-Maxillofacial Surgery* 48 : 33–37.
- Brkic, F.F., Stoiber, S., Al-Gboore, S., Quint, C., Schnoell, J., Scheiflinger, A., *et al.* (2022). Evaluation of the Prognostic Capacity of a Novel Survival Marker in Patients with Sinonasal Squamous Cell Carcinoma. *Nutrients* 14 : 4337.
- Chen, Y.-P., Chan, A.T.C., Le, Q.-T., Blanchard, P., Sun, Y., & Ma, J. (2019). Nasopharyngeal carcinoma. *The Lancet* 394 : 64–80.
- Dahlan, M.S. (2010). Besar Sampel dan Cara Pengambilan Sampel dalam Penelitian Kedokteran dan Kesehatan, 3th edition. ed. Jakarta : Salemba Medika.
- Dwijayanti, F., Prabawa, A., Besral and Herawati, C. (2020). The Five-Year Survival Rate of Patients with Nasopharyngeal Carcinoma Based on Tumor Response after Receiving Neoadjuvant Chemotherapy, Followed by Chemoradiation, in Indonesia: A Retrospective Study. *Oncology*, [online] 98(3), pp.154–160.
- Eisenhauer, E.A., Therasse, P., Bogaerts, J., Schwartz, L.H., Sargent, D., Ford, R., *et al.* (2009). New response evaluation criteria in solid tumours: Revised RECIST guideline (version 1.1). *European Journal of Cancer* 45 : 228–247.
- Fles, R., Indrasari, S.R., Herdini, C., Martini, S., Isfandiari, A., Romdhoni, A.C., *et al.* (2016). Effectiveness of a multicentre nasopharyngeal carcinoma awareness programme in Indonesia. *BMJ Open* 6 : e008571.
- Friedl, M., Stoiber, S., Brkic, F.F., & Kadletz-Wanke, L. (2022). Pretherapeutic Serum Albumin as an Outcome Prognosticator in Head and Neck Adenoid-Cystic Carcinoma. *Biomedicines* 10 : 191.
- Fuhrman, M.P., Charney, P., & Mueller, C.M. (2004). Hepatic proteins and nutrition assessment. *Journal of the American Dietetic Association* 104 : 1258–1264.

- Goldenberg, D., & Goldstein, B.J. (Eds.) (2018). Handbook of otolaryngology: head and neck surgery, Second edition. ed. New York Stuttgart Delhi Rio de Janeiro : Thieme.
- Gullett, N., Rossi, P., Kucuk, O., & Johnstone, P.A.S. (2009). Cancer-induced cachexia: a guide for the oncologist. *Journal of the Society for Integrative Oncology* 7 : 155–169.
- Hegmans, J.P., & Aerts, J.G. (2014). Immunomodulation in cancer. *Current Opinion in Pharmacology* 17 : 17–21.
- Hua, X., Chen, J., Wu, Y., Sha, J., Han, S., & Zhu, X. (2019). Prognostic role of the advanced lung cancer inflammation index in cancer patients: a meta-analysis. *World J Surg Onc* 17 : 177.
- Huang, P.-Y., Wang, C.-T., Cao, K.-J., Guo, X., Guo, L., Mo, H.-Y., *et al.* (2013). Pretreatment body mass index as an independent prognostic factor in patients with locoregionally advanced nasopharyngeal carcinoma treated with chemoradiotherapy: Findings from a randomised trial. *European Journal of Cancer* 49 : 1923–1931.
- Hutajulu, S.H., Howdon, D., Taroeno-Hariadi, K.W., Hardianti, M.S., Purwanto, I., Indrasari, S.R., *et al.* (2021). Survival outcome and prognostic factors of patients with nasopharyngeal cancer in Yogyakarta, Indonesia: A hospital-based retrospective study. *PLoS ONE* 16 : e0246638.
- Iwakiri, D., Sheen, T.-S., Chen, J.-Y., Huang, D.P., & Takada, K. (2005). Epstein–Barr virus-encoded small RNA induces insulin-like growth factor 1 and supports growth of nasopharyngeal carcinoma-derived cell lines. *Oncogene* 24 : 1767–1773.
- Jank, B.J., Kadletz, L., Schnöll, J., Selzer, E., Perisanidis, C., & Heiduschka, G. (2019). Prognostic value of advanced lung cancer inflammation index in head and neck squamous cell carcinoma. *Eur Arch Otorhinolaryngol* 276 : 1487–1492.
- Jin, X., Hu, R., Guo, H., Ding, C., Pi, G., & Tian, M. (2022). Pretreatment Body Mass Index (BMI) as an Independent Prognostic Factor in Nasopharyngeal Carcinoma Survival: A Systematic Review and Meta-Analysis. *Nutrition and Cancer* 74 : 3457–3467.
- Jin, Y., Cai, X.Y., Cai, Y.C., Cao, Y., Xia, Q., Tan, Y.T., *et al.* (2012). To build a prognostic score model containing indispensable tumour markers for metastatic nasopharyngeal carcinoma in an epidemic area. *European Journal of Cancer* 48 : 882–888.
- Johnson, J.T., Rosen, C.A., & Newlands, S. (2014). Bailey’s Head and Neck Surgery - Otolaryngology, Fifth Edition. ed. Lippincott Williams & Wilkins.
- Leliefeld, P.H.C., Koenderman, L., & Pillay, J. (2015). How Neutrophils Shape Adaptive Immune Responses. *Front. Immunol.* 6.

- Lian, C.-L., Zhou, R., Zhou, Y., Zhou, P. and Wu, S.-G. (2023). Assessment of Response to Different Induction Chemotherapy Regimens in Locally Advanced Nasopharyngeal Carcinoma. *Drug Design Development and Therapy*, Volume 17, pp.551–562.
- Liu, Q., Ma, L., Ma, H., Yang, L. and Xu, Z. (2022). Establishment of a prognostic nomogram for patients with locoregionally advanced nasopharyngeal carcinoma incorporating clinical characteristics and dynamic changes in hematological and inflammatory markers. *Frontiers in Oncology*, 12.
- Lund, V.J., Howard, D.J., Bs, M., & Wei, W.I. (2014). Tumors of the Nose, Sinuses and Nasopharynx.
- Ma, J., Lee, N.Y., & Lu, J.J. (Eds.) (2021). *Nasopharyngeal Cancer: A Practical Guide on Diagnosis and Treatment, Practical Guides in Radiation Oncology*. Cham : Springer International Publishing.
- Ma, S.J., Khan, M., Chatterjee, U., Santhosh, S., Hashmi, M., Gill, J., *et al.* (2023). Association of Body Mass Index With Outcomes Among Patients With Head and Neck Cancer Treated With Chemoradiotherapy. *JAMA Netw Open* 6 : e2320513.
- Mantovani, A., Allavena, P., Sica, A., & Balkwill, F. (2008). Cancer-related inflammation. *Nature* 454 : 436–444.
- Mariani, P., Russo, D., Maisto, M., Troiano, G., Caponio, V.C.A., Annunziata, M., *et al.* (2022). Pre-treatment neutrophil-to-lymphocyte ratio is an independent prognostic factor in head and neck squamous cell carcinoma: Meta-analysis and trial sequential analysis. *J Oral Pathology Medicine* 51 : 39–51.
- Mei, Z., Shi, L., Wang, B., Yang, J., Xiao, Z., Du, P., *et al.* (2017). Prognostic role of pretreatment blood neutrophil-to-lymphocyte ratio in advanced cancer survivors: A systematic review and meta-analysis of 66 cohort studies. *Cancer Treatment Reviews* 58 : 1–13.
- Members of the WHO expert consultation (2004). Appropriate body-mass index for Asian populations and its implications for policy and intervention strategies. *The Lancet* 363 : 157–163.
- Minici, R., Siciliano, M.A., Ammendola, M., Santoro, R.C., Barbieri, V., Ranieri, G., *et al.* (2022). Prognostic Role of Neutrophil-to-Lymphocyte Ratio (NLR), Lymphocyte-to-Monocyte Ratio (LMR), Platelet-to-Lymphocyte Ratio (PLR) and Lymphocyte-to-C Reactive Protein Ratio (LCR) in Patients with Hepatocellular Carcinoma (HCC) undergoing Chemoembolizations (TACE) of the Liver: The Unexplored Corner Linking Tumor Microenvironment, Biomarkers and Interventional Radiology. *Cancers* 15 : 257.
- Mujaddid, I., Pratama, A.A., & Alimah, Y. (2025). Characteristics Of Nasopharyngeal Carcinoma Patients In Indonesia 16.

NCCN (2024). NCCN Guidelines for Head and Neck Cancers.

OuYang, P.-Y., Zhang, L.-N., Tang, J., Lan, X.-W., Xiao, Y., Gao, Y.-H., *et al.* (2016). Evaluation of Body Mass Index and Survival of Nasopharyngeal Carcinoma by Propensity-Matched Analysis: An Observational Case-Control Study. *Medicine* 95 : e2380.

Romdhoni, A., Rejeki, P., Guo, H., Milla, C., Melbiarta, R., Visuddho, V., *et al.* (2023). Risk Factors Associated with Nasopharyngeal Cancer Incidences in Indonesia: A Systematic Review and Meta-Analysis. *Asian Pac J Cancer Prev* 24 : 1105–1111.

Roxburgh, C.S.D., & McMillan, D.C. (2014). Cancer and systemic inflammation: treat the tumour and treat the host. *Br J Cancer* 110 : 1409–1412.

Rueda Domínguez, A., Cirauqui, B., García Castaño, A., Alvarez Cabellos, R., Carral Masada, A., Castelo Fernández, B., *et al.* (2022). SEOM-TTCC clinical guideline in nasopharynx cancer (2021). *Clin Transl Oncol* 24 : 670–680.

Shah, A.B., & Nagalli, S. (2024). Nasopharyngeal Carcinoma, in: *StatPearls*. Treasure Island (FL) : StatPearls Publishing.

Shah, J.P., Patel, S.G., Singh, B., & Wong, R.J. (2020). Jatin Shah's head and neck surgery and oncology, Fifth edition. ed. Edinburgh : Elsevier.

Shen, G.-P., Xu, F.-H., He, F., Ruan, H.-L., Cui, C., Chen, L.-Z., *et al.* (2012). Pretreatment Lifestyle Behaviors as Survival Predictors for Patients with Nasopharyngeal Carcinoma. *PLoS ONE* 7 : e36515.

Song, Q., Wang, G., Chu, Y., Zhou, L., Jiang, M., He, Q., *et al.* (2013). TNF- α upregulates cellular inhibitor of apoptosis protein 2 (c-IAP2) via c-Jun N-terminal kinase (JNK) pathway in nasopharyngeal carcinoma. *International Immunopharmacology* 16 : 148–153.

Stan, D.J., Niculet, E., Lungu, M., Onisor, C., Rebegea, L., Vesa, D., *et al.* (2021). Nasopharyngeal carcinoma: A new synthesis of literature data (Review). *Exp Ther Med* 23 : 136.

Su, Z.Y., Siak, P.Y., Leong, C.-O., & Cheah, S.-C. (2022). Nasopharyngeal Carcinoma and Its Microenvironment: Past, Current, and Future Perspectives. *Front. Oncol.* 12 : 840467.

Sun, W., Liu, D.-B., Li, W.-W., Zhang, L.-L., Long, G.-X., Wang, J.-F., *et al.* (2014). Interleukin-6 promotes the migration and invasion of nasopharyngeal carcinoma cell lines and upregulates the expression of MMP-2 and MMP-9. *International Journal of Oncology* 44 : 1551–1560.

Tan, L., & Loh, T. (2010). Benign and Malignant Tumors of the Nasopharynx, in: *Cummings Otolaryngology - Head and Neck Surgery*. pp. 1348–1357, Elsevier.

Topkan, E., Ozdemir, Y., Kucuk, A., Guler, O.C., Sezer, A., Besen, A.A., *et al.*

- (2020). Low Advanced Lung Cancer Inflammation Index Predicts Poor Prognosis in Locally Advanced Nasopharyngeal Carcinoma Patients Treated with Definitive Concurrent Chemoradiotherapy. *Journal of Oncology* 2020 : 1–10.
- Wong, K.C.W., Hui, E.P., Lo, K.-W., Lam, W.K.J., Johnson, D., Li, L., *et al.* (2021). Nasopharyngeal carcinoma: an evolving paradigm. *Nat Rev Clin Oncol* 18 : 679–695.
- Workman, A.D., Palmer, J.N., & Adappa, N.D. (2017). Posttreatment surveillance for sinonasal malignancy. *Current Opinion in Otolaryngology & Head & Neck Surgery* 25 : 86–92.
- Xu, Y., Wu, Z., Ye, W., Xiao, Y., Zheng, W., Chen, Q., Bai, P., Lin, Z. and Chen, C. (2021). Prognostic value of serum uric acid and tumor response to induction chemotherapy in locally advanced nasopharyngeal carcinoma. *BMC Cancer*, [online] 21(1).
- Yang, H., Wang, K., Liang, Z., Guo, S., Zhang, P., Xu, Y., *et al.* (2020). Prognostic role of pre-treatment serum albumin in patients with nasopharyngeal carcinoma: A meta-analysis and systematic review. *Clinical Otolaryngology* 45 : 167–176.
- Yin, J., Qin, Y., Luo, Y., Feng, M., & Lang, J. (2017). Prognostic value of neutrophil-to-lymphocyte ratio for nasopharyngeal carcinoma: A meta-analysis. *Medicine* 96 : e7577.
- Zhang, G., Tsang, C.M., Deng, W., Yip, Y.L., Lui, V.W.-Y., Wong, S.C.C., *et al.* (2013). Enhanced IL-6/IL-6R Signaling Promotes Growth and Malignant Properties in EBV-Infected Premalignant and Cancerous Nasopharyngeal Epithelial Cells. *PLoS ONE* 8 : e62284.
- Zhang, N., Liang, S.-B., Deng, Y.-M., Lu, R.-L., Chen, H.-Y., Zhao, H., *et al.* (2014). Primary tumor regression speed after radiotherapy and its prognostic significance in nasopharyngeal carcinoma: a retrospective study. *BMC Cancer* 14 : 136.