

DAFTAR PUSTAKA

- Aarjane, M., Slassi, S., and Amine, A., 2021, Synthesis, antibacterial evaluation and computational studies of new acridone-1,2,3-triazole hybrids, *J. Mol. Struct.*, 1241, 130636.
- Abdullahi, M., 2023, 3D-QSAR, homology modelling of influenza hemagglutinin receptor (StrainA/WS/1933), molecular dynamics, DFT, and ADMET studies for newly designed inhibitors, *J. Indian Chem. Soc.*, 100, 100975.
- Abdullahi, M., Uzairu, A., Shallangwa, G.A., Mamza, P.A., and Ibrahim, M.T., 2022a, 2D-QSAR, 3D-QSAR, molecular docking and ADMET prediction studies of some novel 2-((1H-indol-3-yl)thio)-N-phenyl-acetamide derivatives as anti-influenza A virus, *Egypt. J. Basic Appl. Sci.*, 9, 510–532.
- Abdullahi, M., Uzairu, A., Shallangwa, G.A., Mamza, P.A., and Ibrahim, M.T., 2022b, In silico modelling studies of 5-benzyl-4-thiazolinone derivatives as influenza neuraminidase inhibitors via 2D-QSAR, 3D-QSAR, molecular docking, and ADMET predictions, *Heliyon*, 8, 10101.
- Acharya, C., Coop, A., Polli, J.E., and MacKerell, A.D., 2010, Recent advances in ligand-based drug design: relevance and utility of the conformationally sampled pharmacophore approach, *Curr. Comput. Aided-Drug Des.*, 7, 10–22.
- Afantitis, A., Melagraki, G., Sarimveis, H., Koutentis, P.A., Igglessi, M.O., and Kollias, G., 2010, A combined LS-SVM & MLR QSAR workflow for predicting the inhibition of CXCR3 receptor by quinazolinone analogs, *Mol. Divers.*, 14, 225–235.
- Afsar, J., Brij, K.S., and Vishnu, D.S., 2021, Quantitative structure-activity relationship study on the MMP-13 inhibitory activity of fused pyrimidine derivatives possessing a 1,2,4-Triazol-3-yl group as a ZBG, *GSC Biol. Pharm. Sci.*, 16, 251–265.
- Aissouq, A.E., Bouachrine, M., Ouammou, A., and Khalil, F., 2022, Computational investigation of unsaturated ketone derivatives as MAO-B inhibitors by using QSAR, ADME-Tox, molecular docking, and molecular dynamics simulations, *Turkish J. Chem.*, 46, 687–703.
- Aissouq, A.E., Chedadi, O., Bouachrine, M., Ouammou, A., and Khalil, F., 2023, Development of novel monoamine oxidase B (MAO-B) inhibitors by combined application of docking-based alignment, 3D-QSAR, ADMET prediction, molecular dynamics simulation, and MM_GBSA binding free energy, *J. Biomol. Struct. Dyn.*, 41, 4667–4680.
- Alder, B.J. and Wainwright, T.E., 1959, Studies in molecular dynamics. I. General method, *J. Chem. Phys.*, 31, 459–466.
- Alsigar, M.T.Y., 2018, Review of docking and drug-likeness as theoretical and computer aided techniques, *Adv. Res. J.*, 2, 1–12.
- Aly, A.S.I., Vaughan, A.M., and Kappe, S.H.I., 2009, Malaria parasite development in the mosquito and infection of the Mammalian Host, *Annu. Rev. Microbiol.*, 63, 195–221.
- Amini, Z., Fatemi, M.H., and Gharaghani, S., 2016, Hybrid docking-QSAR studies of DPP-IV inhibition activities of a series of aminomethyl-piperidones,

Comput. Biol. Chem., 64, 335–345.

- Amusengeri, A., Tata, R.B., and Bishop, O.T., 2020, Understanding the pyrimethamine drug resistance mechanism via combined molecular dynamics and dynamic residue network analysis, *Molecules*, 25, 1–24.
- Anderson, A.C., 2025, Targeting DHFR in parasitic protozoa, *Drug Discov. Today*, 10, 121–128.
- Antypenko, L., Antypenko, O., Fominichenko, A., Karnaukh, I., Kovalenko, S., and Arisawa, M., 2024, Evaluation of fifteen 5,6-dihydrotetrazolo[1,5-c]quinazolines against *Nakaseomyces glabrata*: integrating in vitro studies, molecular docking, QSAR, and in silico toxicity assessments, *J. Fungi*, 10, 816.
- Arief, I., Pranowo, H.D., Mudasir, M., and Wijaya, K., 2020, QSAR-based design of potent betulinic acid derivatives as HIV maturation inhibitors, *C. Nat. Sci.*, 20, 1–12.
- Arodola, O.A. and Soliman, M.E.S., 2017, Quantum mechanics implementation in drug-design workflows: does it really help?, *Drug Des. Devel. Ther.*, 11, 2551.
- Arwansyah, A., Arif, A.R., Syahputra, G., Sukarti, S., and Kurniawan, I., 2021, Theoretical studies of Thiazolyl-Pyrazoline derivatives as promising drugs against malaria by QSAR modelling combined with molecular docking and molecular dynamics simulation, *Mol. Simul.*, 47, 988–1001.
- Auti, P.S., George, G., and Paul, A.T., 2020, Recent advances in the pharmacological diversification of quinazoline/quinazolinone hybrids, *RSC Adv.*, 10, 41353–41392.
- Azzam, R.A., Elsayed, R.E., and Elgemeie, G.H., 2020, Design, synthesis, and antimicrobial evaluation of a new series of N-sulfonamide 2-pyridones as dual inhibitors of DHPS and DHFR enzymes, *ACS Omega*, 5, 10401–10414.
- Balabadra, S., Kotni, M.K., Manga, V., Allanki, A.D., Prasad, R., and Sijwali, P.S., 2017, Synthesis and evaluation of naphthyl bearing 1,2,3-triazole analogs as antiplasmodial agents, cytotoxicity and docking studies, *Bioorganic Med. Chem.*, 25, 221–232.
- Banerjee, P., Kemmler, E., Dunkel, M., and Preissner, R., 2024, ProTox 3.0: a webserver for the prediction of toxicity of chemicals, *Nucleic Acids Res.*, 52, 513–520.
- Bennink, S., Kiesow, M.J., and Pradel, G., 2016, The development of malaria parasites in the mosquito midgut, *Cell. Microbiol.*, 18, 905–918.
- Bhat, H.R., Pandey, P.K., Ghosh, S.K., and Singh, U.P., 2013, Development of 4-aminoquinoline-1,3,5-triazine conjugates as potent antibacterial agent through facile synthetic route, *Med. Chem. Res.*, 22, 5056–5065.
- Castera, C.D., Azas, N., Verhaeghe, P., Hutter, S., Garrigue, P., Dumtre, A., Mbatchi, L., Laget, M., Remusat, V., Sifredi, F., Rault, S., Rathelot, P., and Vanelle, P., 2011, Targeting the human malaria parasite *Plasmodium falciparum*: In vitro identification of a new antiplasmodial hit in 4-phenoxy-2-trichloromethylquinazoline series, *Eur. J. Med. Chem.*, 46, 4184–4191.
- Chaudhary, K.K. and Mishra, N., 2016, A review on molecular docking: novel tool for drug discovery, *JSM Chem.*, 4, 1029.
- Cheng, F., Li, W., Zhou, Y., Shen, J., Wu, Z., Liu, G., Lee, P.W., and Tang, Y.,

- 2012, AdmetSAR: A comprehensive source and free tool for assessment of chemical ADMET properties, *J. Chem. Inf. Model.*, 52, 3099–3105.
- Cheng, T., Li, Q., Zhou, Z., Wang, Y., and Bryant, S.H., 2012, Structure-based virtual screening for drug discovery: A problem-centric review, *AAPS J.*, 14, 133–141.
- Christian, D.J., Vekariya, R.H., Patel, K.D., Rajani, D.P., Rajani, S.D., Parmar, K., Chhabaria, M.T., and Patel, H.D., 2020, Molecular docking and QSAR study of chalcone and pyrimidine derivatives as potent anti-malarial agents against *Plasmodium falciparum*, *Int. Lett. Chem. Phys. Astron.*, 85, 23–34.
- Chu, C.S. and White, N.J., 2021, The prevention and treatment of *Plasmodium vivax* malaria, *PLoS Med.*, 18, 1003561.
- Cowman, A.F., Healer, J., Marapana, D., and Marsh, K., 2016, Malaria: biology and disease, *Cell*, 167, 610–624.
- Daina, Antoine, Blatter, M.C., Baillie, G.V., Palagi, P.M., Marek, D., Xenarios, I., Schwede, T., Michielin, O., and Zoete, V., 2017, Drug design workshop: a web-based educational tool to introduce computer-aided drug design to the general public, *J. Chem. Educ.*, 94, 335–344.
- Daina, A., Michielin, O., and Zoete, V., 2017, SwissADME: A free web tool to evaluate pharmacokinetics, drug-likeness and medicinal chemistry friendliness of small molecules, *Sci. Rep.*, 7, 1–13.
- Das, A., Sarangi, M., Jangid, K., Kumar, Vijay, Kumar, A., Singh, P.P., Kaur, K., Kumar, Vinod, Chakraborty, S., and Jaitak, V., 2024, Identification of 1,3,4-oxadiazoles as tubulin-targeted anticancer agents: a combined field-based 3D-QSAR, pharmacophore model-based virtual screening, molecular docking, molecular dynamics simulation, and density functional theory calculation approach, *J. Biomol. Struct. Dyn.*, 42, 10323–10341.
- Davies, H.M., Nofal, S.D., McLaughlin, E.J., and Osborne, A.R., 2017, Repetitive sequences in malaria parasite proteins, *FEMS Microbiol. Rev.*, 41, 923–940.
- Dhakal, C.P., 2017, Dealing with outliers and influential points while fitting regression, *J. Inst. Sci. Technol.*, 22, 61–65.
- Dick, A. and Cocklin, S., 2020, Bioisosteric replacement as a tool in anti-HIV drug design, *Pharmaceuticals*, 13, 1–16.
- Du, J., Guo, J., Kang, D., Li, Z., Wang, G., Wu, J., Zhang, Z., Fang, H., Hou, X., Huang, Z., Li, G., Lu, X., Liu, X., Ouyang, L., Rao, L., Zhan, P., Zhang, X., and Zhang, Y., 2020, New techniques and strategies in drug discovery, *Chinese Chem. Lett.*, 31, 1695–1708.
- Dyer, M.E., Tjitra, E., Currie, B.J., and Anstey, N.M., 2000, Failure of the pan-malarial antibody of the ICT Malaria Pf/Pv immunochromatographic test to detect symptomatic *Plasmodium malariae* infection, *Trans. R. Soc. Trop. Med. Hyg.*, 94, 518.
- Egan, W.J., Merz, K.M., and Baldwin, J.J., 2000, Prediction of drug absorption using multivariate statistics, *J. Med. Chem.*, 43, 3867–3877.
- Faghihi, K., Safakish, M., Zebardast, T., Hajimahdi, Z., and Zarghi, A., 2019, Molecular docking and QSAR study of 2-benzoxazolinone, quinazoline and diazocoumarin derivatives as anti-HIV-1 agents, *Iran. J. Pharm. Res.*, 18, 1253–1263.

- Faydy, M.E., Lakhrissi, L., Dahaieh, N., Ounine, K., Tuzun, B., Chahboun, N., Boshala, A., AlObaid, A., Warad, I., Lakhrissi, B., and Zarrouk, A., 2024, Synthesis, biological properties, and molecular docking study of novel 1,2,3-triazole-8-quinolinol hybrids, *ACS Omega*, 9, 25395–25409.
- Ferreira, R.D.F. and Schapira, M., 2017, A systematic analysis of atomic protein-ligand interactions in the PDB, *Medchemcomm*, 8, 1970–1981.
- Free, S.M. and Wilson, J.W., 1964, A Mathematical Contribution to Structure-Activity Studies, *J. Med. Chem.*, 7, 395–399.
- Gao, P., Zhang, Jun, Peng, Q., Zhang, Jie, and Glezakou, V.A., 2020, General protocol for the accurate prediction of molecular ¹³C/¹H NMR chemical shifts via machine learning augmented DFT, *J. Chem. Inf. Model.*, 60, 3746–3754.
- Gellis, A., Primas, N., Hutter, S., Lanzas, G., Remusat, V., Verhaeghe, P., Vanelle, P., and Azas, N., 2016, Looking for new antiplasmodial quinazolines: DMAP-catalyzed synthesis of 4-benzyloxy- and 4-aryloxy-2-trichloromethylquinazolines and their in vitro evaluation toward *Plasmodium falciparum*, *Eur. J. Med. Chem.*, 119, 34–44.
- Ghasemi, J.B. and Shiri, F., 2012, Molecular docking and 3D-QSAR studies of falcipain inhibitors using CoMFA, CoMSIA, and Open3DQSAR, *Med. Chem. Res.*, 21, 2788–2806.
- Ghose, A.K., Viswanadhan, V.N., and Wendoloski, J.J., 1999, A knowledge-based approach in designing combinatorial or medicinal chemistry libraries for drug discovery. 1. A qualitative and quantitative characterization of known drug databases, *J. Comb. Chem.*, 1, 55–68.
- Gramatica, P., 2007, Principles of QSAR models validation: Internal and external, *QSAR Comb. Sci.*, 26, 694–701.
- Grimblat, N. and Sarotti, A.M., 2016, Computational chemistry to the rescue: modern toolboxes for the assignment of complex molecules by GIAO NMR calculations, *Chem. - A Eur. J.*, 22, 12246–12261.
- Gromski, P.S., Henson, A.B., Granda, J.M., and Cronin, L., 2019, How to explore chemical space using algorithms and automation, *Nat. Rev. Chem.*, 3, 119–128.
- Hadni, H., Bakhouch, M., and Elhallaoui, M., 2021, 3D-QSAR, Molecular Docking, DFT and ADMET Studies on Quinazoline Derivatives to Explore Novel DHFR Inhibitors, *J. Biomol. Struct. Dyn.*, 1–15.
- Hadni, H. and Elhallaoui, M., 2020, 2D and 3D-QSAR, molecular docking and ADMET properties: In silico studies of azaaurones as antimalarial agents, *New J. Chem.*, 44, 6553–6565.
- Haloui, R., Daoui, O., Mkhayar, K., Yaqoubi, M. El, Elkhatabi, S., Haoudi, A., Rodi, Y., Ouazzani, F., and Chtita, S., 2022, 3D-QSAR, drug-likeness, ADMET prediction, and molecular docking studies in silico of novel 5-oxo-1-thioxo-4,5-dihydro-1H-thiazolo[3,4-a]quinazoline derivatives as MALT1 protease inhibitors for the treatment of B cell lymphoma, *Chem. Pap.*, 77, 2255–2274.
- Hansch, C., Leo, A., Mekapati, S.B., and Kurup, A., 2004, QSAR and ADME, *Bioorg. Med. Chem.*, 12, 3391–3400.
- Hecht, D. and Fogel, G.B., 2012, Towards predictive structure-based models of

- evolved drug resistance,. In, *IEEE Symposium on Computational Intelligence in Bioinformatics and Computational Biology.*, pp. 120–126.
- Ibrahim, Z.Y., Uzairu, A., Shallangwa, G.A., Abechi, S.E., and Isyaku, S., 2022, Quantitative structure-activity relationship, structure-based design, and ADMET studies of pyrimethamine and cycloguanil analogs inhibitors of Plasmodium falciparum dihydrofolate reductase-thymidylate synthase (PfDHFR-TS)., *Chem. Phys. Impact*, 5, 100132.
- Karan, R., Agarwal, P., Sinha, M., and Mahato, N., 2021, Recent advances on quinazoline derivatives: A potential bioactive scaffold in medicinal chemistry, *ChemEngineering*, 5, 73.
- Kemenkes, R., 2020, Buku saku tatalaksana kasus malaria, Jakarta.
- Khamouli, S., Belaidi, S., Bakhouch, M., Chtita, S., Hashmi, M.A., and Qais, F.A., 2022, QSAR modeling, molecular docking, ADMET prediction and molecular dynamics simulations of some 6-arylquinazolin-4-amine derivatives as DYRK1A inhibitors, *J. Mol. Struct.*, 1258, 132659.
- Khan, P.M. and Roy, K., 2018, Current approaches for choosing feature selection and learning algorithms in quantitative structure–activity relationships (QSAR), *Expert Opin. Drug Discov.*, 13, 1075–1089.
- Krieger, E. and Vriend, G., 2015, New ways to boost molecular dynamics simulations, *J. Comput. Chem.*, 36, 996–1007.
- Kubinyi, H., 1997, QSAR and 3D QSAR in drug design Part 1: methodology, *Drug Discov. Today*, 2, 457–467.
- Kurniawan, Y.S., Yudha, E., Nugraha, G., Fatmasari, N., Pranowo, H.D., Jumina, J., and Sholikhah, E.N., 2024, Molecular docking and molecular dynamic investigations of xanthone-chalcone derivatives against epidermal growth factor receptor for preliminary discovery of novel anticancer agent, *Indones. J. Chem.*, 24, 250–266.
- Leonard, J.T. and Roy, K., 2006, On selection of training and test sets for the development of predictive QSAR models, *QSAR Comb. Sci.*, 25, 235–251.
- Lin, Xiaoqian, Li, X., and Lin, Xubo, 2020, A review on applications of computational methods in drug screening and design, *Molecules*, 25, 1375.
- Lindner, S.E., Miller, J.L., and Kappe, S.H.I., 2012, Malaria parasite pre-erythrocytic infection: preparation meets opportunity, *Cell. Microbiol.*, 14, 316–324.
- Lipinski, C., 2001, Avoiding investment in doomed drugs, is poor solubility an industry wide problem?, *Curr. Drug Discov.*, 17–17.
- Lipinski, C. A., Dominy, B.W., and Feeney, P.J., 1997, Drug delivery reviews experimental and computational approaches to estimate solubility and permeability in drug discovery and development settings, *Adv. Drug Deliv. Rev.*, 23, 3–25.
- Lipinski, Christopher A., Lombardo, F., Dominy, B.W., and Feeney, P.J., 1997, Experimental and computational approaches to estimate solubility and permeability in drug discovery and development settings, *Adv. Drug Deliv. Rev.*, 23, 3–25.
- Long, L., Li, R., and Zhang, J., 2025, Artificial intelligence in retrosynthesis prediction and its applications in medicinal chemistry, *J. Med. Chem.*, 68,

2333–2355.

- Madhavan, T., 2012, A Review of 3D-QSAR in Drug Design, *J. Chosun Nat. Sci.*, 5, 1–5.
- Maltarollo, V.G., Kronenberger, T., Wrenger, C., and Honorio, K.M., 2017, Current trends in quantitative structure-activity relationship validation and applications on drug discovery, *Futur. Sci. OA*, 3, 214.
- Martin, Y.C., 2005, A bioavailability score, *J. Med. Chem.*, 48, 3164–3170.
- Masaoudy, Y.E., Aanouz, I., Moukhliiss, Y., Koubi, Y., Maghat, H., Lakhliifi, T., and Bouachrine, M., 2020, 2D-QSAR study of the antimicrobial activity of a series of 5-(substituted benzaldehyde) thiazolidine-2, 4-dione derivatives against *Staphylococcus aureus* by Multiple Linear Regression method, *J. Mater. Environ. Sci*, 11, 1914–1927.
- Mauri, A. and Bertola, M., 2022, Alvascience : A new software suite for the QSAR workflow applied to the blood – brain barrier permeability, *Int. J. Mol. Sci.*, 23, 12882.
- Mishra, M., Agarwal, S., Dixit, A., Mishra, V.K., Kashaw, V., Agrawal, R.K., and Kashaw, S.K., 2020, Integrated computational investigation to develop molecular design of quinazoline scaffold as promising inhibitors of plasmodium lactate dehydrogenase, *J. Mol. Struct.*, 1207, 127808.
- Mishra, M., Mishra, V.K., Senger, P., Pathak, A.K., and Kashaw, S.K., 2014, Exploring QSAR studies on 4-substituted quinazoline derivatives as antimalarial compounds for the development of predictive models, *Med. Chem. Res.*, 23, 1397–1405.
- Moshawih, S., Goh, H.P., Kifli, N., Darwesh, M.A.E.F., Ardianto, C., Goh, K.W., and Ming, L.C., 2023, Identification and optimization of TDP1 inhibitors from anthraquinone and chalcone derivatives: consensus scoring virtual screening and molecular simulations, *J. Biomol. Struct. Dyn.*, 0, 1–25.
- Mudie, D.M., Amidon, G.L., and Amidon, G.E., 2010, Physiological parameters for oral delivery and in vitro testing, *Mol. Pharm.*, 7, 1388–1405.
- Muegge, I., Heald, S.L., and Brittelli, D., 2001, Simple selection criteria for drug-like chemical matter, *J. Med. Chem.*, 44, 1841–1846.
- Netzeva, T.I., Worth, A.P., Aldenberg, T., Benigni, R., Cronin, M.T.D., Gramatica, P., Jaworska, J.S., Kahn, S., Klopman, G., Marchant, C.A., Myatt, G., Nikolova-Jeliazkova, N., Patlewicz, G.Y., Perkins, R., Roberts, D.W., Schultz, T.W., Stanton, D.T., van de Sandt, J.J.M., Tong, W., et al., 2005, Current status of methods for defining the applicability domain of (quantitative) structure-activity relationships: the report and recommendations of ECVAM workshop 521,2, *Altern. to Lab. Anim.*, 33, 155–173.
- Nzila, A., Rottmann, M., Chitnumsub, P., Kiara, S.M., Kamchonwongpaisan, S., Maneeruttanarungroj, C., Taweethai, S., Yeung, B.K.S., Goh, A., Lakshminarayana, S.B., Zou, B., Wong, J., Ma, N.L., Weaver, M., Keller, T.H., Dartois, V., Wittlin, S., Brun, R., Yuthavong, Y., et al., 2010, Preclinical evaluation of the antifolate QN254, 5-chloro- N'6'-(2,5-dimethoxy-benzyl)-quinazoline-2,4,6-triamine, as an antimalarial drug candidate, *Antimicrob. Agents Chemother.*, 54, 2603–2610.
- Ojha, H., Gahlot, P., Tiwari, A.K., Pathak, M., and Kakkar, R., 2011, Quantitative

- structure activity relationship study of 2,4,6-trisubstituted-s-triazine derivatives as antimalarial inhibitors of Plasmodium falciparum Dihydrofolate Reductase, *Chem. Biol. Drug Des.*, 77, 57–62.
- Pagadala, N.S., Syed, K., and Tuszynski, J., 2017, Software for molecular docking: a review, *Biophys. Rev.*, 9, 91–102.
- Paladugula, D., 2017, A review on computational drug designing and discovery, *Biochem. Mol. Biol. Lett.*, 3, 121.
- Pandey, A.K., Baboo, V., Mishra, V.N., Singh, V.K., and Dwivedi, A., 2021, Comparative study of molecular docking, structural, electronic, vibrational spectra and Fukui function studies of thiadiazole containing Schiff base—a complete density functional study, *Polycycl. Aromat. Compd.*, 42, 13–39.
- Pansari, P., 2021, Computational approaches for drug discovery from medicinal plants in the era of data driven research, *Indian Drugs*, 58, 7–23.
- Patel, H.M., Noolvi, M.N., Sharma, P., Jaiswal, V., Bansal, S., Lohan, S., Kumar, S.S., Abbot, V., Dhiman, S., and Bhardwaj, V., 2014, Quantitative structure-activity relationship (QSAR) studies as strategic approach in drug discovery, *Med. Chem. Res.*, 23, 4991–5007.
- Pathak, P., Naumovich, V., Grishina, M., Shukla, P.K., Verma, A., and Potemkin, V., 2019, Quinazoline based 1,3,5-triazine derivatives as cancer inhibitors by impeding the phosphorylated RET tyrosine kinase pathway: design, synthesis, docking, and QSAR study, *Arch Pharm Chem Life Sci.*, 352, 1–18.
- Peterson, D.S., Walliker, D., and Wellems, T.E., 1988, Evidence that a point mutation in dihydrofolate reductase-thymidylate synthase confers resistance to pyrimethamine in falciparum malaria, *Proc. Natl. Acad. Sci. U. S. A.*, 85, 9114–9118.
- Pires, D.E.V., Blundell, T.L., and Ascher, D.B., 2015, pkCSM: Predicting small-molecule pharmacokinetic and toxicity properties using graph-based signatures, *J. Med. Chem.*, 58, 4066–4072.
- Pointe, S.L. and Weaver, D., 2007, A review of density functional theory quantum mechanics as applied to pharmaceutically relevant systems, *Curr. Comput. Aided-Drug Des.*, 3, 290–296.
- Pore, Y., Kuchekar, B., Bhatia, M., and Ingalea, K., 2009, Quantitative structure-activity relationship (QSAR) studies on 5-cyano, N1, 6-disubstituted, 2-thiouracil derivatives as central nervous system depressants, *Dig. J. Nanomater. Biostructures*, 4, 373 – 382.
- Pulung, M.L., Swasono, R.T., Sholikhah, E.N., Yogaswara, R., Primahana, G., and Raharjo, T.J., 2025, Antiplasmodial and metabolite profiling of Hyrtios sp. sponge extract from Southeast Sulawesi marine using LC-HRMS, molecular docking, pharmacokinetic, drug-likeness, toxicity, and molecular dynamics simulation, *J. Multidiscip. Appl. Nat. Sci.*, 5, 487–508.
- Racz, A., Bajusz, D., and Héberger, K., 2015, Consistency of QSAR models: Correct split of training and test sets, ranking of models and performance parameters, *SAR QSAR Environ. Res.*, 26, 683–700.
- Raju, B., Sapra, B., and Silakari, O., 2023, 3D-QSAR assisted identification of selective CYP1B1 inhibitors: an effective bioisosteric replacement/molecular docking/electrostatic complementarity analysis, *Mol. Divers.*, 27, 2673–2693.

- Rajy, M.E., Fadili, M.E., Mrabti, N.N., Zarougui, S., and Elhallaoui, M., 2022, QSAR, molecular docking, ADMET properties in silico studies for a series of 7-propanamide benzoxaboroles as potent anti-cancer agents, *Chinese J. Anal. Chem.*, 50, 100163.
- Rathod, S.B., Vaibhav, D., and Vaidya, R., 2020, In Silico Drug Discovery: A Review, *World J. Pharm. Res.*, 9, 619–637.
- Rhabori, S.E., Aissouq, A.E., Chtita, S., and Khalil, F., 2023, QSAR, molecular docking and ADMET studies of quinoline, isoquinoline and quinazoline derivatives against Plasmodium falciparum malaria, *Struct. Chem.*, 34, 585–603.
- Riniker, S., 2018, Fixed-charge atomistic force fields for molecular dynamics simulations in the condensed phase: An overview, *J. Chem. Inf. Model.*, 58, 565–578.
- Roy, K., Kar, S., and Das, R.N., 2015, Introduction to 3D-QSAR. In, *Understanding the basics of QSAR for applications in pharmaceutical sciences and risk assessment*. Elsevier, pp. 291–317.
- Ruslin, R., Amelia, R., Yamin, Y., Megantara, S., Wu, C., and Arba, M., 2019, 3D-QSAR, molecular docking, and dynamics simulation of quinazoline-phosphoramidate mustard conjugates as EGFR inhibitor, *J. Appl. Pharm. Sci.*, 9, 89–97.
- Sabatini, A., Vacca, A., and Iotti, S., 2012, Balanced biochemical reactions: a new approach to unify chemical and biochemical thermodynamics, *PLoS One*, 7, 29529.
- Sardarian, A., Douglas, K.T., Read, M., Sims, P.F.G., Hyde, J.E., Chitnumsub, P., Sirawaraporn, R., and Sirawaraporn, W., 2003, Pyrimethamine analogs as strong inhibitors of double and quadruple mutants of dihydrofolate reductase in human malaria parasites, *Org. Biomol. Chem.*, 1, 960–964.
- Sarkar, M., Nath, A., Kumer, A., Mallik, C., Akter, F., Moniruzzaman, M., and Ali, M., 2021, Synthesis, Molecular Docking Screening, ADMET and Dynamics Studies of Synthesized 4-(4-methoxyphenyl)-8-methyl-3,4,5,6,7,8-hexahydroquinazolin-2(1H)-one and Quinazolinone Derivatives, *J. Mol. Struct.*, 1244, 1–10.
- Schaduangrat, N., Lampa, S., Simeon, S., Gleeson, M.P., Spjuth, O., and Nantasenamat, C., 2020, Towards reproducible computational drug discovery, *J. Cheminform.*, 12, 1–30.
- Schaller, D., Pach, S., and Wolber, G., 2019, PyRod: Tracing water molecules in molecular dynamics simulations, *J. Chem. Inf. Model.*, 59, 2818–2829.
- Semenov, V.A. and Krivdin, L.B., 2020, DFT computational schemes for ¹H and ¹³C NMR chemical shifts of natural products, exemplified by strychnine, *Magn. Reson. Chem.*, 58, 56–64.
- Senthamarai, K. and Manoj, K., 2015, Outlier detection in multivariate data, *Appl. Math. Sci.*, 9, 2317–2324.
- Shahzad, S., Qadir, M.A., Ahmed, M., Ahmad, S., Khan, M.J., Gulzar, A., and Muddassar, M., 2020, Folic acid-sulfonamide conjugates as antibacterial agents: Design, synthesis and molecular docking studies, *RSC Adv.*, 10, 42983–42992.

- Shree, P., Mishra, P., Kumar, P., Pandey, H., Giri, R., Chaube, R., Garg, N., and Tripathi, Y.B., 2022, In silico screening of Pueraria tuberosa (PTY-2) for targeting COVID-19 by countering dual targets Mpro and TMPRSS2, *J. Biomol. Struct. Dyn.*, 40, 11611–11624.
- Simm, J., Humbeck, L., Zalewski, A., Sturm, N., Heyndrickx, W., Moreau, Y., Beck, B., and Schuffenhauer, A., 2021, Splitting chemical structure data sets for federated privacy-preserving machine learning, *J. Cheminform.*, 13, 1–14.
- Sirawaraporn, W., 1998, Dihydrofolate reductase and antifolate resistance in malaria, *Drug Resist. Updat.*, 1, 397–406.
- Sirawaraporn, W., Sathitkul, T., Sirawaraporn, R., Yuthavong, Y., and Santi, D. V., 1997, Antifolate-resistant mutants of Plasmodium falciparum dihydrofolate reductase, *Proc. Natl. Acad. Sci.*, 94, 1124–1129.
- Sliwoski, G., Kothiwale, S., Meiler, J., and Lowe, E.W., 2014, Computational methods in drug discovery, *Pharmacol. Rev.*, 66, 334–395.
- Sliwoski, G., Mendenhall, J., and Meiler, J., 2015, Autocorrelation descriptor improvements for QSAR: 2DA _ Sign and 3DA _ Sign, *J. Comput. Aided. Mol. Des.*, 30, 209–217.
- Srivastava, S., 2015, Biological activity of quinazoline: a review, *Int. J. Pharma Sci. Res.*, 6, 1206–1213.
- Supandi, Yeni, and Merdekawati, F., 2018, In silico study of pyrazolylaminoquinazoline toxicity by lazarus, protox, and admet predictor, *J. Appl. Pharm. Sci.*, 8, 119–129.
- Surabhi, S. and Singh, B., 2018, Computer Aided Drug Design: An Overview, *J. Drug Deliv. Ther.*, 8, 504–509.
- Syahri, J., Hidayah, N., Hilma, R., Nurohmah, B.A., and Yuanita, E., 2020, Design of new 2,4-substituted furo [3,2-b] indole derivatives as anticancer compounds using quantitative structure-activity relationship (QSAR) and molecular docking, *Molekul*, 15, 9–17.
- Syahri, J., Nasution, H., Nurohmah, B.A., Purwono, B., Yuanita, E., Zakaria, N.H., and Hassan, N.I., 2020, Design, synthesis and biological evaluation of aminoalkylated chalcones as antimalarial agent, *Sains Malaysiana*, 49, 2667–2677.
- Syahri, J., Yuanita, E., Nurohmah, B.A., Wathon, M.H., Syafri, R., Armunanto, R., and Purwono, B., 2017, Xanthone as antimalarial: QSAR analysis, synthesis, molecular docking and in-vitro antimalarial evaluation, *Orient. J. Chem.*, 33, 29–40.
- Sydow, D., Morger, A., Driller, M., and Volkamer, A., 2019, TeachOpenCadd: A teaching platform for computer-aided drug design using open source packages and data, *J. Cheminform.*, 11, 1–7.
- Sydow, D., Rodríguez-Guerra, J., Kimber, T.B., Schaller, D., Taylor, C.J., Chen, Y., Leja, M., Misra, S., Wichmann, M., Ariamajd, A., and Volkamer, A., 2021, TeachOpenCADD 2021: Open Source and FAIR Python Pipelines to Assist in Structural Bioinformatics and Cheminformatics Research,.
- Tabti, K., Sbai, A., Maghat, H., Bouachrine, M., and Lakhlifi, T., 2020, 2D and 3D-QSAR/CoMSIA comparative study on a series of thiazole derivatives as SDHI inhibitors, *Maghrebian J. Pure Appl. Sci.*, 6, 73–90.

- Tmej, C., Chiba, P., Huber, M., Richter, E., Hitzler, M., Schaper, K.J., and Ecker, G., 1998, A combined Hansch/Free-Wilson approach as predictive tool in QSAR studies on propafenone-type modulators of multidrug resistance, *Int. J. Pharm. Med. Chem.*, 331, 233–240.
- Tropsha, A., 2010, Best practices for QSAR model development, validation, and exploitation, *Mol. Inform.*, 29, 476–488.
- Veber, D.F., Johnson, S.R., Cheng, H.Y., Smith, B.R., Ward, K.W., and Kopple, K.D., 2002, Molecular properties that influence the oral bioavailability of drug candidates, *J. Med. Chem.*, 45, 2615–2623.
- Venianakis, T., Oikonomaki, C., Siskos, M.G., Primikyri, A., and Gerothanassis, I.P., 2021, DFT calculations of ¹H NMR chemical shifts of geometric isomers of conjugated linolenic acids, hexadecatrienyl pheromones, and model triene-containing compounds: Structures in solution and revision of NMR assignments, *Molecules*, 26, 1–19.
- Vivo, M.D., Masetti, M., Bottegoni, G., and Cavalli, A., 2016, Role of molecular dynamics and related methods in drug discovery, *J. Med. Chem.*, 59, 4035–4061.
- Wang, J.L., Cheng, L.P., Wang, T.C., Deng, W., and Wu, F.H., 2017, Molecular modeling study of CP-690550 derivatives as JAK3 kinase inhibitors through combined 3D-QSAR, molecular docking, and dynamics simulation techniques, *J. Mol. Graph. Model.*, 72, 178–186.
- Wang, T., Yuan, X. song, Wu, M. Bin, Lin, J.P., and Yang, L.R., 2017, The advancement of multidimensional QSAR for novel drug discovery - where are we headed?, *Expert Opin. Drug Discov.*, 12, 769–784.
- Wold, S., Esbensen, K., and Geladi, P., 1987, Principal component analysis, *Chemom. Intell. Lab. Syst.*, 2, 37–52.
- Wunsch, F.M., Wünsch, B., Bernal, F.A., and Schmidt, T.J., 2021, Quantitative structure-activity relationships of natural product inspired, aminoalkyl-substituted-1-benzopyrans as novel antiplasmodial agents, *Molecules*, 26, 5249.
- Yeni and Rachmania, R.A., 2023, Toxicity of anti-inflammatory substances in *Hemigraphis alternata* leaves: In silico study using ProTox-II, *J. Sains dan Kesehat.*, 5, 810–815.
- Yogaswara, R., Pulung, M.L., Yuliani, S.H., and Istyastono, E.P., 2020, Docking-guided 3D-QSAR studies of 4-aminoquinoline-1, 3, 5-triazines as inhibitors for *Plasmodium falciparum* dihydrofolate reductase, *Indones. J. Chem.*, 20, 1455–1460.
- Yu, X., Gao, L., Jia, L., Yamamoto, Y., and Bao, M., 2018, Synthesis of quinazolin-4(3 H)-ones via the reaction of 2-halobenzamides with nitriles, *J. Org. Chem.*, 83, 10352–10358.
- Yuan, J., Yu, S., Gao, S., Gan, Y., Zhang, Y., Zhang, T., Wang, Y., Yang, L., Shi, J., and Yao, W., 2016, Predicting the biological activities of triazole derivatives as SGLT2 inhibitors using multilayer perceptron neural network, support vector machine, and projection pursuit regression models, *Chemom. Intell. Lab. Syst.*, 156, 166–173.
- Yuthavong, Y., Yuvaniyama, J., Chitnumsub, P., Vanichtanankul, J.,

- Chusacultanachai, S., Tarnchompoo, B., Vilaivan, T., and Kamchonwongpaisan, S., 2005, Malarial (*Plasmodium falciparum*) dihydrofolate reductase-thymidylate synthase: Structural basis for antifolate resistance and development of effective inhibitors, *Parasitology*, 130, 249–259.
- Yuvaniyama, J., Chitnumsub, P., Kamchonwongpaisan, S., Vanichtanankul, J., Sirawaraporn, W., Taylor, P., Walkinshaw, M.D., and Yuthavong, Y., 2003, Insights into antifolate resistance from malarial DHFR-TS structures, *Nat. Struct. Mol. Biol.*, 10, 357–365.
- Zheng, W. and Tropsha, A., 1999, Novel variable selection quantitative structure–property relationship approach based on the k-Nearest-Neighbor principle, *J. Chem. Inf. Comput. Sci.*, 40, 185–194.