

REFERENCES

- Abdel-Kader, S., Sartelli, M., & Abu-Zidan, F. M. (2019). Complicated intra-abdominal infections: a prospective validation study of the WSES Sepsis Severity Score. *Singapore Medical Journal*, 60(6), 317–321.
<https://doi.org/10.11622/smedj.2018120>
- Bone, R. C., Balk, R. A., Cerra, F. B., Dellinger, R. P., Fein, A. M., Knaus, W. A., Schein, R. M., & Sibbald, W. J. (1992). Definitions for sepsis and organ failure and guidelines for the use of innovative therapies in sepsis. The ACCP/SCCM Consensus Conference Committee. American College of Chest Physicians/Society of Critical Care Medicine. *Chest*, 101(6), 1644–1655.
<https://doi.org/10.1378/chest.101.6.1644>
- Brinkmann, V., Reichard, U., Goosmann, C., Fauler, B., Uhlemann, Y., Weiss, D. S., Weinrauch, Y., & Zychlinsky, A. (2004). Neutrophil extracellular traps kill bacteria. *Science (New York, N.Y.)*, 303(5663), 1532–1535.
<https://doi.org/10.1126/science.1092385>
- Christou, N. V., Barie, P. S., Dellinger, E. P., Waymack, J. P., & Stone, H. H. (1993). Surgical Infection Society intra-abdominal infection study: prospective evaluation of management techniques and outcome. *Arch Surg*, 128, 193–199.
- Clements, T. W., Tolonen, M., Ball, C. G., & Kirkpatrick, A. W. (2021). Secondary peritonitis and intra-abdominal sepsis: An increasingly global disease in search

of better systemic therapies. *Scandinavian Journal of Surgery: SJS: Official Organ for the Finnish Surgical Society and the Scandinavian Surgical Society*, 110(2), 139–149. <https://doi.org/10.1177/1457496920984078>

Cox, M. C., Brakenridge, S. C., Stortz, J. A., Hawkins, R. B., Darden, D. B., Ghita, G. L., Mohr, A. M., Moldawer, L. L., Efron, P. A., & Moore, F. A. (2020). Abdominal sepsis patients have a high incidence of chronic critical illness with dismal long-term outcomes. *American Journal of Surgery*, 220(6), 1467–1474. <https://doi.org/10.1016/j.amjsurg.2020.07.016>

Dellinger, E. P., Oreskovich, M. R., Wertz, M. J., Hamasaki, V., & Lennard, E. S. (1984). Risk of infection following laparotomy for penetrating abdominal injury. *Archives of Surgery (Chicago, Ill.: 1960)*, 119(1), 20–27. <https://doi.org/10.1001/archsurg.1984.01390130010002>

Enderson, L. W., & Nolph, K. D. (1969). Altered permeability of the peritoneal membrane after using hypertonic peritoneal dialysis fluid. *J Clin Invest*, 48(6), 992–1001.

Faix, J. D. (2013). Biomarkers of sepsis. *Critical Reviews in Clinical Laboratory Sciences*, 50(1), 23–36. <https://doi.org/10.3109/10408363.2013.764490>

Goto, T., Hara, K., Hashimoto, K., Soeno, S., Shirakawa, T., Sonoo, T., & Nakamura, K. (2020). Validation of chief complaints, medical history, medications, and physician diagnoses structured with an integrated emergency department information system in Japan: the Next Stage ER system. *Acute Medicine & Surgery*, 7(1), e554. <https://doi.org/10.1002/ams2.554>

- Granholm, A., Møller, M. H., Krag, M., Perner, A., & Hjortrup, P. B. (2016). Predictive performance of the Simplified Acute Physiology Score (SAPS) II and the initial Sequential Organ Failure Assessment (SOFA) score in acutely ill intensive care patients: Post-hoc analyses of the SUP-ICU inception cohort study. *PLoS One*, *11*(12), e0168948. <https://doi.org/10.1371/journal.pone.0168948>
- Hall, J. C., Heel, K. A., Papadimitriou, J. M., & Platell, C. (1998). The pathobiology of peritonitis. *Gastroenterology*, *114*(1), 185–196. [https://doi.org/10.1016/s0016-5085\(98\)70646-8](https://doi.org/10.1016/s0016-5085(98)70646-8)
- Heemken, R., Gandawidjaja, L., & Hau, T. (1997). Peritonitis: pathophysiology and local defense mechanisms. *Hepato-Gastroenterology*, *44*(16), 927–936. <https://www.ncbi.nlm.nih.gov/pubmed/9261580>
- Horan, T. C., Gaynes, R. P., Martone, W. J., Jarvis, W. R., & Emori, T. G. (1992). CDC definitions of nosocomial surgical site infections, 1992: A modification of CDC definitions of surgical wound infections. *Infection Control and Hospital Epidemiology: The Official Journal of the Society of Hospital Epidemiologists of America*, *13*(10), 606–608. <https://doi.org/10.2307/30148464>
- Iranya, R. N., Mbiine, R., Semulimi, A. W., Nasige, J., Makumbi, T., & Galukande, M. (2022). Comparison of the PIPAS severity score tool and the QSOFA criteria for predicting in-hospital mortality of peritonitis in a tertiary hospital in Uganda: a prospective cohort study. *BMC Surgery*, *22*(1), 291. <https://doi.org/10.1186/s12893-022-01743-4>

- Kievlan, D. R., Zhang, L. A., Chang, C.-C. H., Angus, D. C., & Seymour, C. W. (2018). Evaluation of repeated quick sepsis-related Organ Failure Assessment measurements among patients with suspected infection. *Critical Care Medicine*, 46(12), 1906–1913. <https://doi.org/10.1097/ccm.0000000000003360>
- Kumar, M., & S., C. C. (2018). SOFA scoring system in assessing prognosis of critically ill surgical and trauma patients: a prospective study. *International Surgery Journal*, 5(7), 2528. <https://doi.org/10.18203/2349-2902.isj20182767>
- Laih, C.-Y., Hsiao, P.-J., Hsieh, P.-F., Wang, Y.-D., Lai, C.-M., Yang, C.-T., Lin, C.-H., & Huang, C.-P. (2022). QSOFA and SOFA scores are valuable tools for predicting postoperative sepsis resulting from ureteroscopic lithotripsy (URSL). *Medicine*, 101(50), e31765. <https://doi.org/10.1097/MD.00000000000031765>
- Lalisang, T. J. M., Usman, N., Hendrawidjaya, I., Handaya, A. Y., Nasution, S., Saunar, R. Y., Loho, T., Karuniawati, A., Moenadjat, Y., & Widyahening, I. S. (2019). Clinical practice guidelines in complicated intra-abdominal infection 2018: An Indonesian perspective. *Surgical Infections*, 20(1), 83–90. <https://doi.org/10.1089/sur.2018.120>
- Luo, X., Li, L., Ou, S., Zeng, Z., & Chen, Z. (2022). Risk factors for mortality in abdominal infection patients in ICU: A retrospective study from 2011 to 2018. *Frontiers in Medicine*, 9, 839284. <https://doi.org/10.3389/fmed.2022.839284>

- Makama, J. G., & Garba, E. S. (2013). Abscess rate of patients with penetrating abdominal injury in Zaria. *Health*, 05(04), 769–773. <https://doi.org/10.4236/health.2013.54102>
- Malangoni, M. A., & Inui, T. (2006). Peritonitis - the Western experience. *World Journal of Emergency Surgery*, 1(1), 25. <https://doi.org/10.1186/1749-7922-1-25>
- Marshall, J. C. (2004). Intra-abdominal infections. *Microbes and Infection*, 6(11), 1015–1025. <https://doi.org/10.1016/j.micinf.2004.05.017>
- Minne, L., Abu-Hanna, A., & de Jonge, E. (2008). Evaluation of SOFA-based models for predicting mortality in the ICU: A systematic review. *Critical Care (London, England)*, 12(6), R161. <https://doi.org/10.1186/cc7160>
- Nugraha, G. L., Septarendra, D., & Lesmana, T. (2022). Comparative study of SOFA, WSESSSS, and CPIRO scoring systems as mortality predictors in a patient with complicated intra-abdominal infection. *Bali Medical Journal*, 11(3), 1397–1403. <https://doi.org/10.15562/bmj.v11i3.3523>
- Pan, J., Zhu, Q., Zhang, X., Xu, J., Pan, L., Mao, X., & Wu, X. (2021). Factors influencing the prognosis of patients with intra-abdominal infection and its value in assessing prognosis. *Infection and Drug Resistance*, 14, 3425–3432. <https://doi.org/10.2147/IDR.S325386>
- Sartelli, M., Abu-Zidan, F. M., Catena, F., Griffiths, E. A., Di Saverio, S., Coimbra, R., Ordoñez, C. A., Leppaniemi, A., Fraga, G. P., Coccolini, F., Agresta, F., Abbas, A., Abdel Kader, S., Agboola, J., Amhed, A., Ajibade, A., Akkucuk, S.,

- Alharthi, B., Anyfantakis, D., ... Ansaloni, L. (2015). Global validation of the WSES Sepsis Severity Score for patients with complicated intra-abdominal infections: a prospective multicentre study (WISS Study). *World Journal of Emergency Surgery*, 10(1), 61. <https://doi.org/10.1186/s13017-015-0055-0>
- Sartelli, M., Catena, F., Ansaloni, L., Coccolini, F., Corbella, D., Moore, E. E., Malangoni, M., Velmahos, G., Coimbra, R., Koike, K., Leppaniemi, A., Biffi, W., Balogh, Z., Bendinelli, C., Gupta, S., Kluger, Y., Agresta, F., Di Saverio, S., Tugnoli, G., ... Barnabé, R. (2014). Complicated intra-abdominal infections worldwide: the definitive data of the CIAOW Study. *World Journal of Emergency Surgery*, 9(1), 37. <https://doi.org/10.1186/1749-7922-9-37>
- Sartelli, M., Catena, F., Ansaloni, L., Moore, E., Malangoni, M., Velmahos, G., Coimbra, R., Koike, K., Leppaniemi, A., Biffi, W., Balogh, Z., Bendinelli, C., Gupta, S., Kluger, Y., Agresta, F., Di Saverio, S., Tugnoli, G., Jovine, E., Ordonez, C., ... Ishii, W. (2013). Complicated intra-abdominal infections in a worldwide context: an observational prospective study (CIAOW Study). *World Journal of Emergency Surgery*, 8(1), 1. <https://doi.org/10.1186/1749-7922-8-1>
- Singer, M., Deutschman, C. S., Seymour, C. W., Shankar-Hari, M., Annane, D., Bauer, M., Bellomo, R., Bernard, G. R., Chiche, J.-D., Coopersmith, C. M., Hotchkiss, R. S., Levy, M. M., Marshall, J. C., Martin, G. S., Opal, S. M., Rubenfeld, G. D., van der Poll, T., Vincent, J.-L., & Angus, D. C. (2016). The third international consensus definitions for sepsis and septic shock (sepsis-3).

JAMA: The Journal of the American Medical Association, 315(8), 801–810.

<https://doi.org/10.1001/jama.2016.0287>

Solomkin, J. S., Mazuski, J. E., Bradley, J. S., Rodvold, K. A., Goldstein, E. J. C., Baron, E. J., O’Neill, P. J., Chow, A. W., Dellinger, E. P., Eachempati, S. R., Gorbach, S., Hilfiker, M., May, A. K., Nathens, A. B., Sawyer, R. G., & Bartlett, J. G. (2010). Diagnosis and management of complicated intra-abdominal infection in adults and children: guidelines by the Surgical Infection Society and the Infectious Diseases Society of America. *Surgical Infections*, 11(1), 79–109. <https://doi.org/10.1089/sur.2009.9930>

Srzić, I., Neseck Adam, V., & Tunjić Pejak, D. (2022). Sepsis definition: What’s new in the treatment guidelines. *Acta Clinica Croatica*, 61(Suppl 1), 67–72. <https://doi.org/10.20471/acc.2022.61.s1.11>

Vincent, J. L., de Mendonça, A., Cantraine, F., Moreno, R., Takala, J., Suter, P. M., Sprung, C. L., Colardyn, F., & Blecher, S. (1998). Use of the SOFA score to assess the incidence of organ dysfunction/failure in intensive care units: results of a multicenter, prospective study. Working group on “sepsis-related problems” of the European Society of Intensive Care Medicine. *Critical Care Medicine*, 26(11), 1793–1800. <https://doi.org/10.1097/00003246-199811000-00016>

Vincent, J. L., Moreno, R., Takala, J., Willatts, S., De Mendonça, A., Bruining, H., Reinhart, C. K., Suter, P. M., & Thijs, L. G. (1996). The SOFA (Sepsis-related Organ Failure Assessment) score to describe organ dysfunction/failure. On


behalf of the Working Group on Sepsis-Related Problems of the European Society of Intensive Care Medicine. *Intensive Care Medicine*, 22(7), 707–710.
<https://doi.org/10.1007/bf01709751>

Wacha, H., Hau, T., Dittmer, R., & Ohmann, C. (1999). Risk factors associated with intraabdominal infections: a prospective multicenter study. Peritonitis Study Group. *Langenbeck s Archives of Surgery*, 384(1), 24–32.
<https://doi.org/10.1007/s004230050169>

Yasin, N. A., Ali, A. M., Ahmed, M. A., & Keleşoğlu, S. İ. (2022). Epidemiological characteristics, antimicrobial susceptibility profile, and surgical outcome of complicated intra-abdominal infection at a tertiary hospital in Somalia. *International Journal of General Medicine*, 15, 7427–7434.
<https://doi.org/10.2147/IJGM.S379711>

Attachment

Patient Intake Form

 UNIVERSITAS GADJAH MADA FACULTY OF MEDICINE PUBLIC HEALTH AND NURSING	<table border="1" style="width: 100%; height: 20px;"> <tr> <td style="width: 10%;"></td> <td style="width: 10%;"></td> <td style="width: 10%;"></td> <td style="width: 10%;"></td> <td style="width: 10%;"></td> <td style="width: 10%;"></td> <td style="width: 10%;"></td> <td style="width: 10%;"></td> <td style="width: 10%;"></td> <td style="width: 10%;"></td> </tr> </table> Medical Record Number										

Data Collection Sheet		
1. Demographic Data		
1.1	Age	<input type="checkbox"/> >18 years old <input type="checkbox"/> <18 years old
1.2	Gender	<input type="checkbox"/> Male <input type="checkbox"/> Female
1.3	Date of hospital Admission	
1.4	Date of hospital discharge	
2. qSOFA Criteria		
2.1	Respiratory rate \geq 22/min	<input type="checkbox"/> YES (1 POINT) <input type="checkbox"/> NO
2.2	Change in mental status	<input type="checkbox"/> YES (1 POINT) <input type="checkbox"/> NO
2.3	Systolic blood pressure \leq 100 mmHg	<input type="checkbox"/> YES (1 POINT) <input type="checkbox"/> NO
3. WSESSSS Criteria of Risk Factors		
3.1	Age > 70 Years old	<input type="checkbox"/> YES (2 POINTS) <input type="checkbox"/> NO
3.2	Immunosuppression	<input type="checkbox"/> YES (3 POINTS) <input type="checkbox"/> NO
Setting of acquisition		
3.3	Healthcare-associated infection	<input type="checkbox"/> YES (2 POINTS) <input type="checkbox"/> NO
Clinical condition at admission		
3.4	Severe sepsis	<input type="checkbox"/> YES (3 POINTS) <input type="checkbox"/> NO
3.5	Septic shock	<input type="checkbox"/> YES (5 POINTS) <input type="checkbox"/> NO

Origin of cIAI		
3.6	Colonic non-diverticular perforation peritonitis	<input type="checkbox"/> YES (2 POINTS) <input type="checkbox"/> NO
3.7	Diveritcular diffuse peritonitis	<input type="checkbox"/> YES (2 POINTS) <input type="checkbox"/> NO
3.8	Postoperative diffuse peritonitis	<input type="checkbox"/> YES (2 POINTS) <input type="checkbox"/> NO
3.9	Small bowel perforation peritonitis	<input type="checkbox"/> YES (3 POINTS) <input type="checkbox"/> NO
Delay in source control		
3.10	Delayed initial intervention >24 hour	<input type="checkbox"/> YES (3 POINTS) <input type="checkbox"/> NO

Notes

- Data is taken from the medical records if patients at RSUP dr.Sardjito from 2020 - 2023
- This sheet is used to record the relationship of qSOFA and WSES Scoring System as Length of Stay and Mortality Predictors in a Patient With Non Trauma Complicated Intra-Abdominal Infection In Dr Sardjito General Hospital in years 2020-2023

Attachment 1. Assessment Paper