

DAFTAR PUSTAKA

- Adetunji, A. I., du Clou, H., Walford, S. N., & Taylor, J. R. N. (2016). Complementary effects of cell wall degrading enzymes together with lactic acid fermentation on cassava tuber cell wall breakdown. *Industrial Crops and Products*, *90*, 110–117. <https://doi.org/10.1016/j.indcrop.2016.06.028>
- Amarullah, Indradewa, Yudono, dan Sunarminto. (2016). Evaluasi kualitas dan hasil tiga varietas ubi kayu. Prosiding Seminar Hasil Penelitian Aneka Kacang dan Umbi, Balitkabi, Malang.
- Bisswanger, H. (2014). Enzyme assays. *Perspectives in Science*, *1*(1–6), 41–55. <https://doi.org/10.1016/j.pisc.2014.02.005>
- Blagbrough, I. S., Bayoumi, S. A. L., Rowan, M. G., & Beeching, J. R. (2010). Cassava: An appraisal of its phytochemistry and its biotechnological prospects. *Phytochemistry*, *71*(17–18), 1940–1951. <https://doi.org/10.1016/j.phytochem.2010.09.001>
- BPS (2016). *Produksi tanaman pangan ATAP tahun 2015*. Badan Pusat Statistik, Jakarta, Indonesia.
- Bradbury, J.H. (2006). Simple wetting method to reduce cyanogens content of cassava flour. *Journal of Food Composition and Analysis* *19*:388-393.
- Breuninger, W.F., Piyachomkwan, K., dan Sriroth, K. (2009). Tapioca/cassava starch: production and use. *Dalam: BeMiller, J. dan Whistler, R (ed.). Starch chemistry and technology*, hal 541-568. 3rd edn. Academic Press, New York.
- Burns, A. E., Bradbury, J. H., Cavagnaro, T. R., & Gleadow, R. M. (2012a). Total cyanide content of cassava food products in Australia. *Journal of Food Composition and Analysis*, *25*(1), 79–82. <https://doi.org/10.1016/j.jfca.2011.06.005>
- Burns, A. E., Gleadow, R. M., Zacarias, A. M., Cuambe, C. E., Miller, R. E., & Cavagnaro, T. R. (2012b). Variations in the chemical composition of cassava (*Manihot esculenta* Crantz) leaves and roots as affected by genotypic and environmental variation. *Journal of Agricultural and Food Chemistry*, *60*(19), 4946–4956. <https://doi.org/10.1021/jf2047288>
- Caffall, K. H., & Mohnen, D. (2009). The structure, function, and biosynthesis of plant cell wall pectic polysaccharides. *Carbohydrate Research*, *344*(14), 1879–1900. <https://doi.org/10.1016/j.carres.2009.05.021>
- Divya-Nair, M. P., Padmaja, G., & Moorthy, S. N. (2011). Biodegradation of cassava starch factory residue using a combination of cellulases, xylanases and hemicellulases. *Biomass and Bioenergy*, *35*(3), 1211–1218. <https://doi.org/10.1016/j.biombioe.2010.12.009>

- Domínguez, E.R, Vázquez-Luna, A., Rodríguez-Landa, J. F., & Díaz-Sobac, R. (2013). Neurotoxic effect of linamarin in rats associated with cassava (*Manihot esculenta* Crantz) consumption. *Food and Chemical Toxicology*, 59, 230–235. <https://doi.org/10.1016/j.fct.2013.06.004>
- Duvernay, W. H., Chinn, M. S., & Yench, G. C. (2013). Hydrolysis and fermentation of sweetpotatoes for production of fermentable sugars and ethanol. *Industrial Crops and Products*, 42(1), 527–537. <https://doi.org/10.1016/j.indcrop.2012.06.028>
- Dzogbefia, V., Ofosu, G., & Oldham, J. (2008). Evaluation of locally produced *Saccharomyces cerevisiae* pectinase enzyme for industrial extraction of starch from cassava in Ghana. *Sci Res Essays*, 3(August), 365–369. Retrieved from <http://www.academicjournals.org/sre/PDF/pdf2008/Aug>
- El-Sharkawy, M. A. (2004). Cassava biology and physiology. *Plant Molecular Biology*, 56(4), 481–501. <https://doi.org/10.1007/s11103-005-2270-7>
- Emmanuel, O. A., Clement, A., Agnes, S. B., Chiwona-Karltun, L., & Drinah, B. N. (2012). Chemical composition and cyanogenic potential of traditional and high yielding CMD resistant cassava (*Manihot esculenta* crantz) varieties. *International Food Research Journal*, 19(1), 175–181.
- Esen, A. (2003). -glucosidase. In *Handbook of Food Enzymology* (pp. 774–786).
- Essers, A.J.A., Bennik, M.H.J. dan Nour, M.J.R. (1995). Mechanisms of increased linamarin degradation during solid-substrate fermentation of cassava. *World Journal of Microbiology and Biotechnology* 11:266-270.
- FAO/WHO, 1995. Codex standard for edible cassava flour, codex standard 176-1989. Food and Agriculture Organisation and World Health Organisation of the United Nations, Rome, Italy.
- Food and Agriculture Organization (FAO) (2017). FAOSTAT Agricultural Database <http://www.fao.org/faostat/en/#search/cassava> [23 Februari 2017].
- Favaro, S. P., Beléia, A., da Silva Fonseca Junior, N., & Waldron, K. W. (2008). The roles of cell wall polymers and intracellular components in the thermal softening of cassava roots. *Food Chemistry*, 108(1), 220–227. <https://doi.org/10.1016/j.foodchem.2007.10.070>
- Ghildyal, N. P., Ramakrishna, M., & Lonsane, B. K. (1989). Comparative Economics of the Production of High Fructose Syrup from Cassava Chips and Cassava Starch. *Starch Stärke*, 41(2), 64–68. <https://doi.org/10.1002/star.19890410207>
- Gohel, H. R., Contractor, C. N., Ghosh, S. K., & Braganza, V. J. (2014). A comparative study of various staining techniques for determination of extra cellular cellulase activity on Carboxy Methyl Cellulose (CMC) agar plates. *International Journal of Current Microbiology and Applied Sciences*, 3(5), 261–266.

- Gueguen, Y., Chemardin, P., Labrot, P., Arnaud, a, & Galzy, P. (1997). Purification and characterization of an intracellular α -glucosidase from a new strain of *Leuconostoc mesenteroides* isolated from cassava. *Journal of Applied Microbiology*, 82(Okafor 1977), 469–476. <https://doi.org/DOI.10.1046/j.1365-2672.1997.00136.x>
- Haque, M.R., & Bradbury, J.H (2002). Total cyanide determination of plants and foods using the picrate and acid hydrolysis methods. *Food Chemistry*, 77(1), 107–114. [https://doi.org/10.1016/S0308-8146\(01\)00313-2](https://doi.org/10.1016/S0308-8146(01)00313-2)
- Ikediodi, C. O., & Onyike, E. (1982). Linamarase activity and detoxification of cassava (*Manihot esculenta*) during fermentation for gari production. *Agricultural and Biological Chemistry*, 46(6), 1667–1669. <https://doi.org/10.1080/00021369.1982.10865302>
- Jekayinfa, S. O., & Olajide, J. O. (2007). Analysis of energy usage in the production of three selected cassava-based foods in Nigeria. *Journal of Food Engineering*, 82(2), 217–226. <https://doi.org/10.1016/j.jfoodeng.2007.02.003>
- Johnson, R., Padmaja, G., & Moorthy, S. N. (2009). Comparative production of glucose and high fructose syrup from cassava and sweet potato roots by direct conversion techniques. *Innovative Food Science and Emerging Technologies*, 10(4), 616–620. <https://doi.org/10.1016/j.ifset.2009.04.001>
- Jorgensen, K., Morant, A. V., Morant, M., Jensen, N. B., Olsen, C. E., Kannangara, R., Motawia, M.S., Moller, B.L., Bak, S. (2011). Biosynthesis of the Cyanogenic Glucosides Linamarin and Lotaustralin in Cassava: Isolation, Biochemical Characterization, and Expression Pattern of CYP71E7, the Oxime-Metabolizing Cytochrome P450 Enzyme. *Plant Physiology*, 155(1), 282–292. <https://doi.org/10.1104/pp.110.164053>
- Juturu, V., & Wu, J. C. (2014). Microbial cellulases: Engineering, production and applications. *Renewable and Sustainable Energy Reviews*, 33, 188–203. <https://doi.org/10.1016/j.rser.2014.01.077>
- Kementrian Perindustrian, (2017). http://www.kemenperin.go.id/statistik/query_komoditi.php?komoditi=glucose+syrup [19 Agustus 2017].
- McMahon, J. M., White, W. L. B., & Sayre, R. T. (1995). Cyanogenesis in cassava (*Manihot esculenta* Crantz). *J. Exp. Botany*, 46(288), 731–741.
- Mkpong, O. E., Yan, H., Chism, G., & Sayre, R. T. (1990). Purification, characterization, and localization of linamarase in cassava. *Plant Physiology*, 93, 176–181. <https://doi.org/10.1104/pp.93.1.176>
- Montagnac, J. A., Davis, C. R., & Tanumihardjo, S. A. (2009). Processing techniques to reduce toxicity and antinutrients of Cassava for use as a staple food. *Comprehensive Reviews in Food Science and Food Safety*, 8(1), 17–27. <https://doi.org/10.1111/j.1541-4337.2008.00064.x>

- Montagnac, J. a, Davis, C. R., & Tanumihardjo, S. a. (2009). Nutritional Value of Cassava for Use as a Staple Food and Recent Advances for Improvement. *Comprehensive Reviews in Food Science and Food Safety*, 8.
- Mosier, N., Wyman, C., Dale, B., Elander, R., Lee, Y. Y., Holtzapfle, M., & Ladisch, M. (2005). Features of promising technologies for pretreatment of lignocellulosic biomass. *Bioresource Technology*, 96(6), 673–686. <https://doi.org/10.1016/j.biortech.2004.06.025>
- Murugan, K., Sekar, K., & Sohaibani, S. A.-. (2012). Detoxification of cyanides in cassava flour by linamarase of *Bacillus subtilis* KM05 isolated from cassava peel, 11(28), 7232–7237. <https://doi.org/10.5897/AJB11.831>
- Nambisan, B. (1999). Cassava latex as a source of linamarase for determination of linamarin. *Journal of Agricultural and Food Chemistry*, 47(2), 372–373. <https://doi.org/10.1021/jf980768r>
- Nambisan, B. (2011). Strategies for elimination of cyanogens from cassava for reducing toxicity and improving food safety. *Food and Chemical Toxicology*, 49(3), 690–693. <https://doi.org/10.1016/j.fct.2010.10.035>
- Nambisan, B., & Sundaresan, S. (1994). Distribution of linamarin and its metabolising enzymes in cassava tissues. *Journal of the Science of Food and Agriculture*, 66(4), 503–507. <https://doi.org/10.1002/jsfa.2740660413>
- Nwokoro, O., & Anya, F. O. (2011). Linamarase Enzyme from *Lactobacillus delbrueckii* NRRL -763: Purification and some properties of a -glucosidase. *Journal of the Mexican Chemical Society*, 55(4), 246–250.
- Oguntoyinbo, F. A., & Dodd, C. E. R. (2010). Bacterial dynamics during the spontaneous fermentation of cassava dough in gari production. *Food Control*, 21(3), 306–312. <https://doi.org/10.1016/j.foodcont.2009.06.010>
- Palmer, T. (1991). *Understanding enzymes*. 3rd edition, Ellis Horwood Ltd., Great Britain.
- Petruccioli, M., Brimer, L., Cicalini, A. R., & Federici, F. (1999). The linamarase of *Mucor circinelloides* LU M40 and its detoxifying activity on cassava. *Journal of Applied Microbiology*, 86(2), 302–310. <https://doi.org/10.1046/j.1365-2672.1999.00655.x>
- Ribeiro, J.M.C.R. (1994). *Nutritive value of treated straw*. In : Tisserand J.-L. (ed.). *Les pailles dans l'alimentation des ruminants en zone méditerranéenne*. Zaragoza: CIHEAM, hal 79-95.
- Rose, J.K.C. (2003). *The plant cell wall*. Blackwell Publishing, CRC Press, USA.
- Ryu, D. D. Y., & Mandels, M. (1980). Cellulases: biosynthesis and applications. *Microbiology and Technology*, 2, 91–102.

- Salvador, L. D., Sukanuma, T., Kitahara, K., Tanoue, H., & Ichiki, M. (2000). Monosaccharide composition of sweetpotato fiber and cell wall polysaccharides from sweetpotato, cassava, and potato analyzed by the high-performance anion exchange chromatography with pulsed amperometric detection method. *Journal of Agricultural and Food Chemistry*, 48(8), 3448–3454. <https://doi.org/10.1021/jf991089z>
- Santana, M. A., Vásquez, V., Matehus, J., & Aldao, R. R. (2002). Linamarase Expression in Cassava Cultivars with Roots of Low-and High-Cyanide Content. *Source: Plant Physiology*, 129(4), 1686–1694. <https://doi.org/10.1104/pp.000927>
- Silveira, M. H. L., Aguiar, R. S., Siika-aho, M., & Ramos, L. P. (2014). Assessment of the enzymatic hydrolysis profile of cellulosic substrates based on reducing sugar release. *Bioresource Technology*, 151, 392–396. <https://doi.org/10.1016/j.biortech.2013.09.135>
- Sornyotha, S., Kyu, K. L., & Ratanakhanokchai, K. (2010). An efficient treatment for detoxification process of cassava starch by plant cell wall-degrading enzymes. *Journal of Bioscience and Bioengineering*, 109(1), 9–14. <https://doi.org/10.1016/j.jbiosc.2009.06.021>
- Sriroth, K., Piyachomkwan, K., Wanlapatit, S., & Oates, C. G. (2000). Cassava starch technology: the Thai experience. *Starch - Stärke*, 52(12), 439–449. [https://doi.org/10.1002/1521-379X\(200012\)52:12<439::AID-STAR439>3.0.CO;2-E](https://doi.org/10.1002/1521-379X(200012)52:12<439::AID-STAR439>3.0.CO;2-E)
- Sundari, T. (2010). Petunjuk teknis pengenalan varietas unggul dan teknik budidaya ubi kayu (Materi pelatihan agribisnis bagi KMPH). Balai Penelitian Kacang Kacangan dan Umbi Umbian, Malang.
- Tenkanen, M., Niku-Paavola, M., Linder, M., dan Viikari, L. (2003). Cellulases in food processing. *Dalam: Whitaker, J.R., Vorgan, A.G.J., dan Wong, D.W.S (ed.). Handbook of Food Enzymology*, hal 754-773. Marcel Dekker Inc., New York.
- Tylleskär, T., Rosling, H., Banea, M., Bikangi, N., Cooke, R. D., & Poulter, N. H. (1992). Cassava cyanogens and konzo, an upper motoneuron disease found in Africa. *The Lancet*, 339(8787), 208–211. [https://doi.org/10.1016/0140-6736\(92\)90006-O](https://doi.org/10.1016/0140-6736(92)90006-O)
- Utomo, A.R.P. (2013). Pengaruh pengupasan singkong dan penambahan -glukanase pada bubur singkong untuk produksi sirup glukosa secara enzimatik. Tesis Program Studi Ilmu dan Teknologi Pangan, Program Pascasarjana, UGM, Yogyakarta.
- Van Dyk, J. S., & Pletschke, B. I. (2012). A review of lignocellulose bioconversion using enzymatic hydrolysis and synergistic cooperation between enzymes-Factors affecting enzymes, conversion and synergy. *Biotechnology Advances*, 30(6), 1458–1480. <https://doi.org/10.1016/j.biotechadv.2012.03.002>

- Vetter, J. (2000). Plant cyanogenic glycosides. *Toxicon*, 38(1), 11–36. [https://doi.org/10.1016/S0041-0101\(99\)00128-2](https://doi.org/10.1016/S0041-0101(99)00128-2)
- Westby, A. (2002). Cassava utilization, storage and small-scale processing. *Cassava: Biology, Production and Utilization*, 1, 281–300. <https://doi.org/10.1264/jsme2.ME11201>
- White, W., Arias-Garzon, D., McMahon, J., & Sayre, R. (1998). Cyanogenesis in cassava. The role of hydroxynitrile lyase in root cyanide production. *Plant Physiology*, 116(4), 1219–1225. <https://doi.org/10.1104/pp.116.4.1219>
- Whitaker, J. R. (1972). *Principles of enzymology for the food science*. New York: Marcel Dekker, Inc.
- Yeoh, H.H., Tan, T.K. dan Loh, C.M. (1995). Sources of fungal linamarase. *World Journal of Microbiology and Technology* 11:678-680.
- Yeoh, H.H. dan Sun, F. (2001). Assesing cyanogens content in cassava-based food using the enzyme-dipstick method. *Food and Chemical Toxicology* 39(7):649-653.