



DAFTAR PUSTAKA

- Anderson, J. B., Horsley, M., Pratt, J., King, E., Iyer, S. B., Varadarajan, K., Lannon, C., Schidlow, D. N., Williams, R., Slicker, J. 2012. Variation in growth of infants with a single ventricle. *Journal of Pediatrics*, 161(1), pp.16–21.e3.
- Ansong-Assoku, B., Adnan, M., Daley, S.F., Ankola, P.A. 2025. Neonatal jaundice. StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing. Tersedia di: <https://www.ncbi.nlm.nih.gov/books/NBK532930/> [Diakses 3 Mei 2025].
- Aradhana, A., Panda, S.K., Jena, P., Singh, R., Kumar, V., Mehta, S., Sharma, D. & Gupta, P. 2025. Intravenous fluid supplementation in management of severe unconjugated hyperbilirubinemia in preterm neonates: a randomized controlled trial. *European Journal of Pediatrics*, 184, 383.
- Asaye, S., Bekele, M., Getachew, A., Fufa, D., Adugna, T., Tadasa, E. 2023. Hyperbilirubinemia and associated factors among neonates admitted to the neonatal care unit in Jimma Medical Center. *Clinical Medicine Insights: Pediatrics*, 17, pp. 1–10.
- Asefa, G.G., Gebrewahid, T.G., Nuguse, H., Gebremichael, M.W., Birhane, M., Zereabruk, K., Zemicheal, T.M., Hailay, A., Aberha Abrha, W., Aregawi Hadera, S., Gebreegzabiher Hailu, A., Hagos Beyene, B., Ayele Dagnazgi, E., Gebretsadkan Tekulu, F. & Welay, F.T. 2020. Determinants of neonatal jaundice among neonates admitted to neonatal intensive care unit in public general hospitals of Central Zone, Tigray, Northern Ethiopia, 2019: a case-control study. *BioMed Research International*, 2020, Article ID 4743974.
- Auliasari, N.A. 2019. Faktor risiko kejadian ikterus neonatorum di RSUD Dr. Soetomo Surabaya. Skripsi tidak dipublikasikan, Universitas Airlangga, Surabaya. Tersedia di: <https://repository.unair.ac.id/82244> [Diakses 3 Juni 2025].
- Bahadir, A., Okur, M., Kocamaz, H., Dönmez, M. Bozkaya, D., 2018. A comparison of single and repeated phototherapy in neonates: Weight loss and additional fluid requirements. *Journal of Clinical Neonatology*, 7(3), pp.154–157.
- Bansal, A., Jain, S., Parmar, V.R. Chawla, D. 2010. Bilirubin rebound after intensive phototherapy for neonatal jaundice. *Indian Pediatrics*, 47(7), pp.607–609.
- Bell, E.F., Acarregui, M.J. and Oh, W. (2020) Fluid and electrolyte management in the newborn. In: Polin, R.A., Abman, S.H., Rowitch, D.H. and Benitz, W.E. (eds.) *Fetal and Neonatal Physiology*. 5th edn. Philadelphia: Elsevier, pp. 392–403.



- Bhutani, V.K., Wong, R.J., Turkewitz, D., Rauch, D.A. 2024. Phototherapy to prevent severe neonatal hyperbilirubinemia in the newborn infant ≥ 35 weeks of gestation: technical report. *Pediatrics*, 154(3), e2024068026
- Bhutani, V.K., Zipursky, A., Blencowe, H., Khanna, R., Sgro, M., Ebbesen, F., Bell, J., Mori, R., Slusher, T.M., Fahmy, N., Paul, V.K., Du, L., Okolo, A.A., de Almeida, M.F., Olusanya, B.O., Kumar, P., Cousens, S. & Lawn, J.E. 2013. Neonatal hyperbilirubinemia and Rhesus disease of the newborn: incidence and impairment estimates for 2010 at regional and global levels. *Pediatric Research*, 74(Suppl 1), pp. 86–100.
- Boonyarittipong, P., Kriangburapa, W. Booranavanich, K., 2008. Effectiveness of double-surface intensive phototherapy versus single-surface intensive phototherapy for neonatal hyperbilirubinemia. *Journal of the Medical Association of Thailand*, 91(1), pp.50–55.
- Chou, C.Y., Lin, H.C., Tsao, P.N., Hsieh, W.S. & Chen, C.Y. 2024. Clinical outcomes of repeated phototherapy in neonates: A retrospective review. *Journal of Neonatal Medicine*.
- Chow, S.-C., Shao, J., Wang, H. 2008. *Sample Size Calculations in Clinical Research*. 2nd edn. Boca Raton, FL: Chapman & Hall/CRC.
- Darmady, E.M., Barr, D.G.D. 2020. Renal function and sodium handling in the newborn: hormonal and physiological adaptations. *Pediatric Nephrology*, 35(6): 987–995.
- Demirel, H.N., Ozumut, S.S., Ovalı, H.F. 2024. Continuous versus intermittent phototherapy in treatment of neonatal jaundice: a randomized controlled trial. *European Journal of Pediatrics*, 183, pp. 3389–3396.
- Diala, U.M., Usman, F., Appiah, D., Olorunfemi, O., Afolayan, J.M., Lawal, M.A., Okeke, I.O., Ibrahim, B., Mohammed, A., Bello, A., Yusuf, K., Salami, A., Sadiq, U. & Adejumo, O. 2023. Global prevalence of severe neonatal jaundice among hospital admissions: A systematic review and meta-analysis. *Journal of Clinical Medicine*, 12(11), p.3738.
- DiTomasso, D. Paiva, A.L. 2015. Weighing the facts: a systematic review of expected patterns of weight loss in full-term, breastfed infants. *Journal of Human Lactation*, 31(4): 1–12.
- Elhawary, I.M., El-Banna, A.A., Abdelrahman, S.M., El-Sayed, H.M. El-Masry, S.A. 2018. Incidence and risk factors of post-phototherapy neonatal rebound hyperbilirubinemia. *World Journal of Pediatrics*, 14(4), pp.350–356.



- El-Mazary, A.M., Abdel-Latif, M.M. El-Kashishy, R.A., 2020. Risk factors for significant weight loss in jaundiced neonates receiving phototherapy. *Egyptian Pediatric Association Gazette*, 68(1), pp.1–5.
- Elshazzly, M., Anekar, A.A., Shumway, K.R. Slusher, T.M. 2025. *Physiology, newborn*. Treasure Island (FL): StatPearls Publishing. Tersedia di: <https://www.ncbi.nlm.nih.gov/books/NBK499951/> [Diakses 3 Mei 2025].
- Flaherman, V. J., Schaefer, E. W., Kuzniewicz, M. W., Li, S. X., Walsh, E. M., Paul, I. M. 2015. Early weight loss nomograms for exclusively breastfed newborns. *Pediatrics*, 135(1), e16–e23.
- Flenady, V.J. dan Woodgate, P.G. 2003. Radiant warmers versus incubators for regulating body temperature in newborn infants. *Cochrane Database of Systematic Reviews*, 4, CD000435.
- Fonseca, M.J., Severo, M., Barros, H., Santos, A.C. 2014. Determinants of weight changes during the first 96 hours of life in full-term newborns. *Birth*, 41(2): 160–168.
- Girija, G., Reddy, K.R., Ravichander, B. 2021. The efficacy of intermittent versus continuous phototherapy in the duration of phototherapy required in the given bilirubin level. *International Journal of Paediatrics and Geriatrics*, 4(1), pp. 18–21.
- Gomella, T.L., Cunningham, M.D., Eyal, F.G. and Zenk, K.E., 2020. *Gomella's Neonatology: Management, Procedures, On-Call Problems, Diseases, and Drugs*. 8th ed. New York: McGraw-Hill Education.
- Gounaris, A. Drakatos, A., Economou, E., Kounnis, V. & Karpathios, T. 1998. Gut hormone levels in neonates under phototherapy. *Early Human Development*, 51(1), pp.57–60.
- Grünhagen, D.J., de Boer, M.G.J., de Beaufort, A.J., 2002. Transepidermal water loss during halogen spotlight phototherapy in preterm infants. *Pediatric Research*, 51(3), pp. 402–406.
- Hansen, T.W.R., 2021. Narrative review of the epidemiology of neonatal jaundice. *Pediatric Medicine*, 4, p.18.
- Haseli, A., Mohammadi, M., Sabzehei, M.K., Gholami, M. & Heidarzadeh, M. 2025. Postnatal weight loss and exclusive breastfeeding in newborn. *BMC Pediatrics*, 25(366).
- Joseph, A. Samant, H., 2025. *Jaundice*. Treasure Island (FL): StatPearls Publishing. Tersedia di: <https://www.ncbi.nlm.nih.gov/books/NBK544252/> [Diakses 3 Mei 2025].



- Kadavakollu, N., Gopal, K.V. 2023. Electrolyte changes in neonates receiving phototherapy. *International Journal of Pediatrics and Neonatology*.
- Kato, S. Nakamura, H., Saito, M., Yamada, K., Okuyama, K. & Ibara, S. 2019. Standardization of phototherapy for neonatal hyperbilirubinemia using multiple-wavelength irradiance integration. *Pediatrics & Neonatology*.
- Kemper, A.R. Newman, T.B., Slaughter, J.L., Maisels, M.J., Watchko, J.F., Downs, S.M. 2022. American Academy of Pediatrics Subcommittee on Hyperbilirubinemia Clinical practice guideline revision: Management of hyperbilirubinemia in the newborn infant 35 or more weeks of gestation. *Pediatrics*, 150(3), e2022058859.
- Kovacic, K., Flaherman, V.J., Kuzniewicz, M.W., Walsh, E.M. & Wickremasinghe, A.C. 2020. Supporting breastfeeding in infants hospitalized for jaundice. *Hospital Pediatrics*, 10(6), pp.502–508.
- Kumar, P., Chawla, D. Deorari, A., 2022. Light-emitting diode phototherapy for unconjugated hyperbilirubinaemia in neonates. *Cochrane Database of Systematic Reviews*, (3), CD007969.
- Lin, Q., Chen, Y., Chen, J., Zhang, L., Wang, Y. & Huang, H. 2022. Risk factors for neonatal hyperbilirubinemia: A systematic review and meta-analysis. *Translational Pediatrics*, 11(6), pp.1001–1009. <https://doi.org/10.21037/tp-22-229>.
- Lindqvist, D., Hansson, M., Thomas, M., Hulzebos, C.V., Vitek, L., Blokzijl, A., Van Berkel, M. 2025. Bilirubin photoisomers in neonatal jaundice. *International Journal of Molecular Sciences*, 26(21): 10791.
- Lopez, M.G., Cadenas, E.G. Cadenas, L.G., 2018. Weight decrease in full-term newborns in the first 48 hours post natal. *Revista Chilena de Pediatría*, 89(3), pp.325–331.
- Maayan-Metzger, A., Yosipovitch, G., Hadad, E. Sirota, L. 2001. Transepidermal water loss and skin hydration in preterm infants during phototherapy. *American Journal of Perinatology*, 18(7), pp.393–396.
- Maisels, M.J. McDonagh, A.F., 2008. Phototherapy for neonatal jaundice. *New England Journal of Medicine*, 358(9), pp.920–928.
- Mani, M. Kaur, A., Kaur, P., Kler, N. 2022. Effects of different types of phototherapy units on neonatal jaundice: A cross-sectional study. *Journal of Neonatal Nursing*, 28(6), pp.347–351.
- Medoff-Cooper, B., Irving, S.Y., Hanlon, A.L., Saenz, L., Stallings, V.A., Ravishankar, C. 2016. Weight change in infants with a functionally

univentricular heart: From surgical intervention to hospital discharge. *Cardiology in the Young*, 26(4), pp.1–8.

Morris, B.H., Oh, W., Tyson, J.E., Stevenson, D.K., Phelps, D.L., O’Shea, T.M., McDavid, G.E., Perritt, R.L., Van Meurs, K.P., Vohr, B.R., Lemons, J.A., Bauer, C.R., Ehrenkranz, R.A. 2008. Aggressive vs. conservative phototherapy for infants with extremely low birth weight. *New England Journal of Medicine*, 359(18), pp.1885–1896.

Naveed, A., Syed, F., Hussain, M., Ahmed, R., Khan, S., Iqbal, Z. (2024). Effect of fluid supplementation on the duration of phototherapy in neonates with non-hemolytic hyperbilirubinemia: a randomized controlled trial. *History of Medicine Journal*, 5(9), pp. 12–17.

Ng, P.C., Lam, H.S., Chui, K.M., Wong, R.P.O., Leung, T.F., Li, K., Wong, E., Fok, T.F., Chan, I.H.S. 2021. Diagnostic markers for neonatal sepsis. *Current Opinion in Pediatrics*, 33(2), pp.219–226.

Nizam, M.A., Khan, M.R., Chowdhury, M.A.K., Sarker, M.A.S., Islam, M.S., Rahman, M.M. 2020. Efficacy of double versus single phototherapy in treatment of neonatal jaundice: A meta-analysis. *European Journal of Pediatrics*, 179(6), pp.865–874.

Noel-Weiss, J., Courant, G., Woodend, A.K. 2008. Physiological weight loss in the breastfed neonate: a systematic review. *Open Medicine*, 2(4): e99–e110.

Novoa, J.M., González, A., Jiménez, R., Fernández, M., García, L. & Pérez, M. 2023. LED versus non-LED phototherapy devices for neonatal hyperbilirubinemia: A systematic review and meta-analysis. *Pediatric Research*, 93(2), pp.288–295.

Okulu, E., 2024. Neonatal jaundice: Recommendations for follow-up and treatment. *Global Pediatrics*, 7(3), p.100131.

Parlawar, S., Agarwal, A., Dutta, K.P., Joshi, U. 2024. To study the effect of phototherapy on serum calcium of term neonates with hyperbilirubinemia. *International Journal of Pharmaceutical and Clinical Research*, 16(12), pp. 969–973.

Polit, D.F., Beck, C.T. 2021. *Nursing Research: Generating and Assessing Evidence for Nursing Practice*. 11th ed. Philadelphia: Wolters Kluwer.

Regnault, N., Botton, J., Forhan, A., Hankard, R., Thiebaugeorges, O., Hillier, T.A., Charles, M.A., Heude, B. 2011. Determinants of early weight loss in term infants: a population-based study. *Pediatric Research*, 70(3): 281–288.

Sabzehei, M.K., Basiri, B., Shokouhi, M., Gholami, M. & Heidarzadeh, M. 2021. Single-surface intensive phototherapy or double-surface intensive

- phototherapy in neonatal non-hemolytic hyperbilirubinemia: A comparison of effectiveness and complications. *Medical Journal of the Islamic Republic of Iran*, 35, p.192.
- Sarici, S.U., Yurdakök, M., Serdar, M.A., Alpay, F. & Erdem, G. 2024. Duration of a standard phototherapy course in jaundiced newborns with no neurotoxicity risk. *Journal of Tropical Pediatrics*, 70(6), fmae037.
- Scrafford, C.G., Mullany, L.C., Katz, J., Khatry, S.K., LeClerq, S.C., Darmstadt, G.L. & Tielsch, J.M. 2013. Incidence of and risk factors for neonatal jaundice among newborns in southern Nepal. *Tropical Medicine and International Health*, 18(11), pp.1317–1328.
- Seidman, D.S., Moise, J., Ergaz, Z., Laor, A., Vreman, H.J. & Stevenson, D.K. 2003. A prospective randomized controlled study of phototherapy using blue and blue-green light-emitting devices, and conventional halogen-quartz phototherapy. *Journal of Perinatology*, 23, pp.123–127.
- Shin, T., Kim, H.J., Lee, J.H., Park, M.S. & Kim, S.H. 2025. Effect of phototherapy on blood pressure, heart rate, and body temperature in early preterm infants with gestational age <32 weeks. *Perinatology*, 36(1).
- Siddappa, A.M., Prekker, F.L., Slusher, T.M., 2020. Improving effectiveness of phototherapy in an academic center: A quality improvement project. *Global Pediatric Health*, 7, 2333794X20969275.
- Singh, A., Koritala, T., Jialal, I., 2025. *Unconjugated hyperbilirubinemia*. Treasure Island (FL): StatPearls Publishing. Tersedia di: <https://www.ncbi.nlm.nih.gov/books/NBK549796/> [Diakses 3 Mei 2025].
- Tan, W., Lee, J., Wong, H.B. & Chan, S.H. 1998. Phototherapy duration and weight loss in neonates. *Journal of Paediatrics*.
- Treesirichod, A., Nuntnarumit, P., Chutinet, A. & Chareonsiriwatana, W. 2022. Incidence of hyperthermia after conventional phototherapy in term infants. *Journal of Neonatal Nursing*.
- Vandborg, P.K., Hansen, B.M., Greisen, G. & Ebbesen, F. 2012. Dose-response relationship of phototherapy for hyperbilirubinemia. *Pediatrics*, 130(2), pp.e352–e357.
- Waite, W.M., Taylor, J.A., 2016. Phototherapy for the treatment of neonatal jaundice and breastfeeding duration and exclusivity. *Breastfeeding Medicine*, 11, pp.180–185.
- Wang, J., Guo, G., Li, A., Cai, W.-Q., Wang, X., 2021. Challenges of phototherapy for neonatal hyperbilirubinemia (Review). *Experimental and Therapeutic Medicine*, 21, p. 231.



- Wu, P.Y., Wong, W.H., Hodgman, J.E. & Lindquist, R. 1974. Effect of phototherapy in preterm infants on growth in the neonatal period. *Journal of Pediatrics*, 85(4), pp.563–566.
- Wynn, J.L., Wong, H.R., 2010. Pathophysiology and treatment of septic shock in neonates. *Clinics in Perinatology*, 37(2), pp.439–479.
- Yassin, O., Hassan, M., El-Gendy, F.M., El-Shimi, O.S. & El-Farrash, R.A. 2021. Impact of LED phototherapy on incubator temperature and neonatal stress. *Journal of Perinatal Medicine*, 49(5), pp.587–593.
- Yurdakok, M. 2015. Phototherapy in the newborn: What's new? *Journal of Pediatric and Neonatal Individualized Medicine*, 4(2), e040255.
- Zhao, D., Li, Y., Wu, X., Zhang, L., Huang, J. & Li, H. 2021. Effect of early supplemental formula intervention on body weight loss and hyperbilirubinemia at 72 hours in term neonates. *Frontiers in Pediatrics*, 9: 625536.