

DAFTAR PUSTAKA

- Abdelmohsen, S.M., Soliman, A., Fathy, M., Takrouney, M.H., Ahmed, A.A., El-Din Mostafa Madany, M., Abou-Taleb, A., *et al.* (2025). Upper Egyptian outcome of Kasai portoenterostomy in biliary atresia: A cross-sectional retrospective study, *Pediatrics and Neonatology* [Preprint]. <https://doi.org/10.1016/j.pedneo.2025.05.013>.
- Adrin, O.E., Widyastuti, Y. & Sari, D. (2024). Risk Factors of Early Complications in Pediatric Biliary Atresia After Kasai Procedure in Dr. Sardjito Hospital, *Malaysian Journal of Medicine and Health Sciences*, 20, pp. 11–16. <https://doi.org/10.47836/mjmhs.20.s6.3>.
- Azarow, K., James Phillips, M., Sandier, A.D., Hagerstrand, Inga & Superina Toronto, R.A. (1997). Biliary Atresia: Should All Patients Undergo a Portoenterostomy?, *Journal of Pediatric Surgery*, 32(2), pp. 168–4.
- Bakula, A. & Dadalski, M. (2015). AST-to-Platelet Ratio Index (APRI) as Marker in Liver Disease, in *Biomarkers in Liver Disease*.
- Balistreri, W.F., Bezerra, J.A. & Ryckman, F.C. (2014). Biliary atresia and other disorders of the extrahepatic bile ducts, in *Liver Disease in Children, Fourth Edition*. Cambridge University Press, pp. 155–176. <https://doi.org/10.1017/CBO9781139012102.012>.
- Bessho, K. (2015). Complications and Quality of Life in Long-Term Survivors of Biliary Atresia with Their Native Livers, *Journal of Pediatrics*, 167(6), pp. 1202–1206. <https://doi.org/10.1016/j.jpeds.2015.08.041>.
- Brits, E., Brown, S., Botes, L., Sempa, J.B. & Pienaar, M. (2025). Aspartate Aminotransferase-to-platelet Ratio Index (APRI) as Biomarker for Liver Damage in Biliary Atresia (BA): A Meta-analysis, *Journal of Pediatric Surgery*. W.B. Saunders. <https://doi.org/10.1016/j.jpedsurg.2025.162234>.
- Cavallo, L., Kovar, E.M., Aqul, A., McLoughlin, L., Mittal, N.K., Rodriguez-Baez, N., Shneider, B.L., *et al.* (2022). The Epidemiology of Biliary Atresia: Exploring the Role of Developmental Factors on Birth Prevalence, *The Journal of pediatrics*, 246, p. 89. <https://doi.org/10.1016/J.JPEDI.2022.03.038>.
- Chen, S.Y., Lin, C.C., Tsan, Y.T., Chan, W.C., Wang, J. Der, Chou, Y.J. & Lin, C.H. (2018). Number of cholangitis episodes as a prognostic marker to predict timing of liver transplantation in biliary atresia patients after Kasai portoenterostomy, *BMC Pediatrics*, 18(1). <https://doi.org/10.1186/s12887-018-1074-2>.

- Chiba, T. & Kasai, M. (1975). *An Attempt to Determine Surgical Indication for Biliary Atresia by Laboratory Examination, Tohoku J. exp. Med.*
- Chiu, C.-Y., Chen, P.-H., Chan, C.-F., Chang, M.-H. & Wu, T.-C. (2013). Biliary Atresia in Preterm Infants in Taiwan: A Nationwide Survey, *The Journal of Pediatrics*, 163(1), pp. 100-103.e1. <https://doi.org/10.1016/j.jpeds.2012.12.085>.
- Chung, P.H.Y., Chan, E.K.W., Yeung, F., Chan, A.C.Y., Mou, J.W.C., Lee, K.H., Hung, J.W.S., *et al.* (2021). Life long follow up and management strategies of patients living with native livers after Kasai portoenterostomy, *Scientific Reports*, 11(1). <https://doi.org/10.1038/s41598-021-90860-w>.
- Chusilp, S., Sookpotarom, P., Tepmalai, K., Rajatapiti, P., Chongsrisawat, V., Poovorawan, Y. & Vejchapipat, P. (2016). Prognostic values of serum bilirubin at 7th day post-Kasai for survival with native livers in patients with biliary atresia, *Pediatric Surgery International*, 32(10), pp. 927–931. <https://doi.org/10.1007/s00383-016-3951-9>.
- Clark, T.G., Bradburn, M.J., Love, S.B. & Altman, D.G. (2003). Survival analysis part IV: Further concepts and methods in survival analysis, *British Journal of Cancer*, pp. 781–786. <https://doi.org/10.1038/sj.bjc.6601117>.
- Davenport, M., Ong, E., Sharif, K., Alizai, N., McClean, P., Hadzic, N. & Kelly, D.A. (2011). Biliary atresia in England and Wales: Results of centralization and new benchmark, *Journal of Pediatric Surgery*, 46(9), pp. 1689–1694. <https://doi.org/10.1016/j.jpedsurg.2011.04.013>.
- Emre, S., Benibol, Y., Hakalmaz, A.E., Karakurt, B., Kalyoncu Ucar, A., Kepil, N., Beşer, Ö.F., *et al.* (2025). Three decades of progress: evolution of outcomes and prognostic indicators in biliary atresia management, *BMC Pediatrics*, 25(1). <https://doi.org/10.1186/s12887-025-05848-6>.
- Evans, H.M., Asher, M.I., Cameron-Christie, S., Farthing, S., McCall, J., Robertson, S.P., Wong, H., *et al.* (2018). Ethnic Disparity in the Incidence and Outcome of Biliary Atresia in New Zealand, *Journal of Pediatric Gastroenterology and Nutrition*, 66(2), pp. 218–221. <https://doi.org/10.1097/MPG.0000000000001781>.
- Fang, H., Li, Z., Xian, R., Yin, Y., Wang, J., Guo, H., Dai, X., *et al.* (2023). Early life growth and developmental trajectory in children with biliary atresia undergoing primary liver transplantation, *Frontiers in Pediatrics*, 11. <https://doi.org/10.3389/fped.2023.1198360>.
- Farahdina, F., Prihaningtyas, R.A., Setyo-boedi, B., Sera, A.C. & Arief, S. (2024). Seeking Liver Fibrosis in Biliary Atresia using De Ritis Ratio, *Biomolecular and*

Health Science Journal, 7(2), pp. 96–100.
https://doi.org/10.4103/bhsj.bhsj_10_24.

Gad, E.H., Kamel, Y., Salem, T.A.H., Ali, M.A.H. & Sallam, A.N. (2021). Short- and long-term outcomes after Kasai operation for type III biliary atresia: Twenty years of experience in a single tertiary Egyptian center-A retrospective cohort study, *Annals of Medicine and Surgery*, 62, pp. 302–314.
<https://doi.org/10.1016/j.amsu.2021.01.052>.

Garcia, A. V., Cowles, R.A., Kato, T. & Hardy, M.A. (2012). Morio Kasai: A remarkable impact beyond the Kasai procedure, in *Journal of Pediatric Surgery*, pp. 1023–1027. <https://doi.org/10.1016/j.jpedsurg.2012.01.065>.

Ge, L., Zhan, J., Gao, W., Zhao, S., Xu, X. & Dou, R. (2020). Relevant factors for early liver transplantation after Kasai portoenterostomy, *BMC Pediatrics*, 20(1).
<https://doi.org/10.1186/s12887-020-02355-8>.

Girard, M., Jannot, A.S., Besnard, M., Jacquemin, E. & Henrion-Caude, A. (2012). Biliary atresia: Does ethnicity matter?, *Journal of Hepatology*, 57(3), pp. 700–701. <https://doi.org/10.1016/j.jhep.2012.03.011>.

Gunadi, Sirait, D.N., Budiarti, L.R., Paramita, V.M.W., Fauzi, A.R., Ryantono, F., Afandy, D., *et al.* (2020). Histopathological findings for prediction of liver cirrhosis and survival in biliary atresia patients after Kasai procedure, *Diagnostic Pathology*, 15(1). <https://doi.org/10.1186/s13000-020-00996-y>.

He, L., Ip, D.K.M., Tam, G., Lui, V.C.H., Tam, P.K.H. & Chung, P.H.Y. (2021). Biomarkers for the diagnosis and post-Kasai portoenterostomy prognosis of biliary atresia: a systematic review and meta-analysis, *Scientific Reports*, 11(1).
<https://doi.org/10.1038/s41598-021-91072-y>.

Hoshino, E., Muto, Y., Sakai, K., Shimohata, N., Urayama, K.Y. & Suzuki, M. (2023). Age at surgery and native liver survival in biliary atresia: a systematic review and meta-analysis, *European Journal of Pediatrics*, 182(6), pp. 2693–2704. <https://doi.org/10.1007/s00431-023-04925-1>.

Huang, C.Y., Chang, M.H., Chen, H.L., Ni, Y.H., Hsu, H.Y. & Wu, J.F. (2020). Bilirubin level 1 week after hepatportoenterostomy predicts native liver survival in biliary atresia, *Pediatric Research*, 87(4), pp. 730–734.
<https://doi.org/10.1038/s41390-019-0610-6>.

Hukkinen, M., Fischler, B., Liliemark, U., Gatzinsky, V., Ekvall, N., Almaas, R., Elmi, H., *et al.* (2025). Multicenter Study on Early Predictors of Biliary Atresia Outcomes, *Journal of Pediatric Surgery*, 60(9).
<https://doi.org/10.1016/j.jpedsurg.2025.162404>.

- Ihn, K., Na, Y., Ho, I.G., Lee, D., Koh, H. & Han, S.J. (2019). A periodic comparison of the survival and prognostic factors of biliary atresia after Kasai portoenterostomy: a single-center study in Korea, *Pediatric Surgery International*, 35(3), pp. 285–292. <https://doi.org/10.1007/s00383-018-04434-5>.
- Jahangirnia, A., Oltean, I., Nasr, Y., Islam, N., Weir, A., de Nanassy, J., Nasr, A., *et al.* (2022). Peri-Operative Liver Fibrosis and Native Liver Survival in Pediatric Patients with Biliary Atresia: A Systematic Review and Meta-Analysis, *Pediatric Gastroenterology, Hepatology and Nutrition*. Korean Society of Pediatric Gastroenterology, Hepatology and Nutrition, pp. 353–375. <https://doi.org/10.5223/pghn.2022.25.5.353>.
- Jain, A.K., Anand, R., Lerret, S., Yanni, G., Chen, J.Y., Mohammad, S., Doyle, M., *et al.* (2021). Outcomes following liver transplantation in young infants: Data from the SPLIT registry, *American Journal of Transplantation*, 21(3), pp. 1113–1127. <https://doi.org/10.1111/ajt.16236>.
- Jayarathna, M.J.S., Dassanayake, B.K., Dorji, T., Lucero-Prisno, D.E., Samarasinghe, S., Pinto, V. & Lamawansa, M.D. (2024). Challenges of liver transplantation programs in low- and middle-income countries: An experience from Sri Lanka, *Public Health Challenges*. John Wiley and Sons Inc. <https://doi.org/10.1002/puh2.162>.
- Kakos, C.D., Ziogas, I.A., Alexopoulos, S.P. & Tsoulfas, G. (2021). Management of biliary atresia: To transplant or not to transplant, *World Journal of Transplantation*. Baishideng Publishing Group Co, pp. 400–409. <https://doi.org/10.5500/WJT.V11.I9.400>.
- Kelly, D.A. & Davenport, M. (2007). Current management of biliary atresia, *Archives of Disease in Childhood*, pp. 1132–1135. <https://doi.org/10.1136/adc.2006.101451>.
- Khanna, K., Bhatnagar, V., Agarwala, S., Srinivas, M. & Gupta, S. (2018). Ratio of preoperative and postoperative serum bilirubin levels predicts early outcome following biliary atresia surgery, *Journal of Indian Association of Pediatric Surgeons*, 23(2), pp. 81–86. https://doi.org/10.4103/jiaps.JIAPS_192_17.
- Kusumawati, N.R., Ritonga, R.S., Kevin, C., Sulaiman, S., Siahaan, S.S. & Pratiwi, J. (2022). Demographic Characteristics of Children with Biliary Atresia in dr. Kariadi General Hospital, Semarang, *Pediatric Gastroenterology, Hepatology, and Nutrition*, 1. <https://doi.org/10.58427/apg>.
- Lala, V., Zubair, M. & Affiliations, D.A.M. (2023). *Liver Function Tests*. Internet: StatPearls.

- Lee, K.J., Kim, J.W., Moon, J.S. & Ko, J.S. (2017). Epidemiology of Biliary Atresia in Korea, *Journal of Korean Medical Science*, 32(4), pp. 656–660. <https://doi.org/10.3346/JKMS.2017.32.4.656>.
- Li, B., Jia, Y. & Ren, H. (2025). Value of preoperative APRI and FIB-4 in assessing short-term prognosis after Kasai portoenterostomy, *World Journal of Pediatric Surgery*, 8(1). <https://doi.org/10.1136/wjps-2025-001002>.
- Lianti, H., Kurnia, N., Rinaldhy, K., Aji, A.S., Ismet, M.F. & Amaliah, R. (2020). Analysis of knowledge regarding Biliary Atresia among healthcare providers and laypersons in East Jakarta after educational intervention, *ASEAN Journal of Community Engagement*, 4(1), p. 13. <https://doi.org/10.7454/ajce.v4i1.1053>.
- Liu, J.Q., Chen, W.J., Zhou, M.J., Li, W.F., Tang, J. & Zhou, Q.C. (2021). A Nomogram Predicting the Prognosis of Children With Biliary Atresia After Hepatoportoenterostomy, *Frontiers in Pediatrics*, 9. <https://doi.org/10.3389/fped.2021.641318>.
- Liu, S., Yang, Q., Ji, Q., Wang, Z., Sun, R. & Zhan, J. (2024). Effect of Kasai procedure on liver transplantation in children with biliary atresia: a systematic review and updated meta-analysis, *Translational Pediatrics*, 13(1), pp. 10–31. <https://doi.org/10.21037/tp-23-504>.
- Lurie, Y., Webb, M., Cytter-Kuint, R., Shteingart, S. & Lederkremer, G.Z. (2015). Non-invasive diagnosis of liver fibrosis and cirrhosis, *World Journal of Gastroenterology*, 21(41), pp. 11567–11583. <https://doi.org/10.3748/wjg.v21.i41.11567>.
- Machino, K., Mimori, K., Ogata, S., Minami, Y., Shimizu, H., Yamashita, M. & Tanaka, H. (2024). Pre-operative Serum Albumin Predicts Native Liver Survival in Biliary Atresia, *African journal of paediatric surgery : AJPS*, 21(4), pp. 232–235. https://doi.org/10.4103/ajps.ajps_158_22.
- Mack, C.L. (2015). What Causes Biliary Atresia? Unique Aspects of the Neonatal Immune System Provide Clues to Disease Pathogenesis, *CMGH*, 1(3), pp. 267–274. <https://doi.org/10.1016/J.JCMGH.2015.04.001/ASSET/6AB47DBE-F72A-4341-A478-BD3387A0F210/MAIN.ASSETS/GR1.JPG>.
- Mo, Y.H., Chen, H.L., Hsu, W.M., Chang, C.H. & Peng, S.S.F. (2021). A noninvasive index to predict liver cirrhosis in biliary atresia, *Pediatric Radiology*, 51(2), pp. 257–264. <https://doi.org/10.1007/s00247-020-04823-w>.
- Mojumder, Md.R.K., Hossain, A.Z., Huda, S.M.S., Arifuzzaman, A.S.M., Mohanta, D.C., Rahman, Md.M., Rahman, Md.S., *et al.* (2024). Relationship between Preoperative Aspartate Aminotransferase-To-Platelet Ratio Index (APRI) and

- Gamma Glutamyl Transpeptidase (GGT) With Early Postoperative Jaundice Clearance and Native Liver Survival in Biliary Atresia, *SAS Journal of Surgery*, 12(11), pp. 1267–1276. <https://doi.org/10.36347/sasjs.2024.v10i11.017>.
- Nakajima, H., Koga, H., Okawada, M., Nakamura, H., Lane, G.J. & Yamataka, A. (2018). Does time taken to achieve jaundice-clearance influence survival of the native liver in post-Kasai biliary atresia?, *World Journal of Pediatrics*, 14(2), pp. 191–196. <https://doi.org/10.1007/s12519-018-0139-5>.
- Neto, B., Borges-Dias, M., Trindade, E., Estevaõ-Costa, J. & Campos, J.M. (2018). Biliary Atresia-Clinical Series, *GE Portuguese Journal of Gastroenterology*, 25(2), pp. 68–73. <https://doi.org/10.1159/000480708>.
- Ng, V.L., Sorensen, L.G., Alonso, E.M., Fredericks, E.M., Ye, W., Moore, J., Karpen, S.J., *et al.* (2018). Neurodevelopmental Outcome of Young Children with Biliary Atresia and Native Liver: Results from the ChiLDReN Study, *Journal of Pediatrics*, 196, pp. 139-147.e3. <https://doi.org/10.1016/j.jpeds.2017.12.048>.
- Nio, M., Ohi, R., Miyano, T., Saeki, M., Shiraki, K. & Tanaka, K. (2003). Five- and 10-year survival rates after surgery for biliary atresia: A report from the Japanese Biliary Atresia Registry, *Journal of Pediatric Surgery*, 38(7), pp. 997–1000. [https://doi.org/10.1016/S0022-3468\(03\)00178-7](https://doi.org/10.1016/S0022-3468(03)00178-7).
- Noor, H.Z. & Makhmudi, A. (2020). The impact of serum total bilirubin, alanine transaminase and gamma-glutamyl transferase on survival of biliary atresia patients following Kasai procedure, *Med J Malaysia*, 75.
- Nyholm, I., Hukkinen, M. & Pakarinen, M.P. (2024). Predicting and managing liver fibrosis in biliary atresia, *Seminars in Pediatric Surgery*. W.B. Saunders. <https://doi.org/10.1016/j.sempedsurg.2025.151473>.
- Oswari, H., Rahayatri, T.H. & Soedibyo, S. (2020). Pediatric Living Donor Liver Transplant in Indonesia's National Referral Hospital, *Transplantation*, 104(7), pp. 1305–1307. <https://doi.org/10.1097/TP.00000000000003154>.
- Prawirohartono, E.P. (2022). *Memahami Penelitian Epidemiologi Klinis secara Mudah*. 23rd ed. Yogyakarta: Gadjah Mada University Press.
- Ramos-Gonzalez, G., Elisofon, S., Dee, E.C., Staffa, S.J., Medford, S., Lillehei, C. & Kim, H.B. (2019). Predictors of Need for Liver Transplantation in Children Undergoing Hepatoportoenterostomy for Biliary Atresia, *Journal of Pediatric Surgery*, 54(6), pp. 1127–1131. <https://doi.org/10.1016/j.jpedsurg.2019.02.051>.

- Rhu, J., Jung, S.M., Choe, Y.H., Seo, J.M. & Lee, S.K. (2012). PELD score and age as a prognostic index of biliary atresia patients undergoing Kasai portoenterostomy, *Pediatric Surgery International*, 28(4), pp. 385–391. <https://doi.org/10.1007/s00383-012-3060-3>.
- Sastroasmoro, S., Firmansyah, A., Said, M., Akib, A.P. & Harimurti, K. (2014). *Analisis Kesintasan (Survival Analysis): Dasar-dasar Metodologi Penelitian Klinis*. 5th ed. Yogyakarta: Sagung Seto.
- Schreiber, R.A., Harpavat, S., Hulscher, J.B.F., Wildhaber, B.E., Schreiber, R.A., Harpavat, S., Hulscher, J.B.F., *et al.* (2022). Biliary Atresia in 2021: Epidemiology, Screening and Public Policy, *Journal of Clinical Medicine* 2022, Vol. 11, 11(4). <https://doi.org/10.3390/JCM11040999>.
- Seah Lee, W. & Fan Chai, P. (2010). Clinical Features Differentiating Biliary Atresia from Other Causes of Neonatal Cholestasis, *Annals Academy of Medicine*, 39(8), pp. 648–654.
- Shah, I., Magdum, N. & Shetty, N.S. (2020). Correlation of APRI Index with Metavir Index in Children with Biliary Atresia, *Tropical Gastroenterology*, 41(4), pp. 168–173.
- Shneider, B.L., Brown, M.B., Haber, B., Whittington, P.F., Schwarz, K., Squires, R., Bezerra, J., *et al.* (2006). A multicenter study of the outcome of biliary atresia in the United States, 1997 to 2000, *Journal of Pediatrics*, 148(4). <https://doi.org/10.1016/j.jpeds.2005.12.054>.
- Shneider, B.L., Magee, J.C., Karpen, S.J., Rand, E.B., Narkewicz, M.R., Bass, L.M., Schwarz, K., *et al.* (2016). Total Serum Bilirubin within 3 Months of Hepatoportoenterostomy Predicts Short-Term Outcomes in Biliary Atresia, *Journal of Pediatrics*, 170, pp. 211-217.e2. <https://doi.org/10.1016/j.jpeds.2015.11.058>.
- Shneider, B.L. & Mazariegos, G. V. (2007). Biliary atresia: A transplant perspective, *Liver Transplantation*, 13(11), pp. 1482–1495. <https://doi.org/10.1002/lt.21303>.
- Sinha, C.K. & Davenport, M. (2008). Biliary Atresia, *J Indian Assoc Pediatr Surg*, 13(2), p. 49. Available at: <http://www.jiaps.com>.
- Soeters, P.B., Wolfe, R.R. & Shenkin, A. (2019). Hypoalbuminemia: Pathogenesis and Clinical Significance, *Journal of Parenteral and Enteral Nutrition*. John Wiley and Sons Inc, pp. 181–193. <https://doi.org/10.1002/jpen.1451>.

- Stetson, A., Bondoc, A. & Tiao, G. (2024). Revision Kasai portoenterostomy: A review of indications and outcomes, *Seminars in Pediatric Surgery*. W.B. Saunders. <https://doi.org/10.1016/j.sempedsurg.2025.151476>.
- Sun, S., Zheng, S., Lu, X., Chen, G., Ma, Y., Chen, L. & Dong, K. (2018). Clinical and pathological features of patients with biliary atresia who survived for more than 5 years with native liver, *Pediatric Surgery International*, 34(4), pp. 381–386. <https://doi.org/10.1007/s00383-018-4231-7>.
- Sundaram, S.S., Mack, C.L., Feldman, A.G. & Sokol, R.J. (2017). Biliary atresia: Indications and timing of liver transplantation and optimization of pretransplant care, *Liver Transplantation*. John Wiley and Sons Ltd, pp. 96–109. <https://doi.org/10.1002/lt.24640>.
- Suominen, J.S., Lampela, H., Heikkilä, P., Lohi, J., Jalanko, H. & Pakarinen, M.P. (2015). APRI predicts native liver survival by reflecting portal fibrogenesis and hepatic neovascularization at the time of portoenterostomy in biliary atresia, *Journal of Pediatric Surgery*, 50(9), pp. 1528–1531. <https://doi.org/10.1016/j.jpedsurg.2014.11.046>.
- Suthantirakumar, R.L. & Gupte, G.L. (2025). Timing and Indications for Liver Transplantation for Children with Chronic Liver Disease, *Children*. Multidisciplinary Digital Publishing Institute (MDPI). <https://doi.org/10.3390/children12040449>.
- Tambucci, R., de Magnée, C., Szabo, M., Channaoui, A., Pire, A., de Meester de Betzenbroeck, V., Scheers, I., *et al.* (2021). Sequential Treatment of Biliary Atresia With Kasai Hepatopertoenterostomy and Liver Transplantation: Benefits, Risks, and Outcome in 393 Children, *Frontiers in Pediatrics*, 9. <https://doi.org/10.3389/fped.2021.697581>.
- Terui, K., Okubo, R., Sakurai, T., Sasaki, H., Kudo, W. & Kawaguchi, Y. (2025). Prediction of Native Liver Survival in Patients With Biliary Atresia 20 Years After the Kasai Procedure, *Journal of Pediatric Surgery*, 60(3). <https://doi.org/https://doi.org/10.1016/j.jpedsurg.2025.162158>.
- Tessier, M.E.M., Harpavat, S., Shepherd, R.W., Hiremath, G.S., Brandt, M.L., Fisher, A. & Goss, J.A. (2014). Beyond the Pediatric end-stage liver disease system: Solutions for infants with biliary atresia requiring liver transplant, *World Journal of Gastroenterology*. WJG Press, pp. 11062–11068. <https://doi.org/10.3748/wjg.v20.i32.11062>.
- Wada, H., Muraji, T., Yokoi, A., Okamoto, T., Sato, S., Takamizawa, S., Tsugawa, J., *et al.* (2007). Insignificant seasonal and geographical variation in incidence

- of biliary atresia in Japan: a regional survey of over 20 years, *Journal of Pediatric Surgery*, 42(12), pp. 2090–2092.
<https://doi.org/10.1016/j.jpedsurg.2007.08.035>.
- Wang, X., Bai, F., Pang, H. & George, S.L. (2019). Bias-adjusted Kaplan–Meier survival curves for marginal treatment effect in observational studies, *Journal of Biopharmaceutical Statistics*, 29(4), pp. 592–605.
<https://doi.org/10.1080/10543406.2019.1633659>.
- Wang, Z., Chen, Y., Peng, C., Pang, W., Zhang, T., Wu, D., Shen, Q., *et al.* (2019). Five-year native liver survival analysis in biliary atresia from a single large Chinese center: The death/liver transplantation hazard change and the importance of rapid early clearance of jaundice, *Journal of Pediatric Surgery*, 54(8), pp. 1680–1685. <https://doi.org/10.1016/j.jpedsurg.2018.09.025>.
- Xu, X., Wang, X., Ding, M., Zhao, Y., Zhao, Li, Zhao, Linsheng, Li, M., *et al.* (2023). Development and post-Kasai procedure prognostic relevance of histological features for biliary atresia, *BMC Pediatrics*, 23(1). <https://doi.org/10.1186/s12887-023-04413-3>.
- Yang, C., Hy, J., Tan, B., Xiong, Q., Wang, R., Hu, J., Le, Y., *et al.* (2025). Progression Characteristics of Biliary Atresia during The Waiting Period for Liver Transplantation: a Single-Center Observational Study, *Front. Med.* [Preprint]. <https://doi.org/https://doi.org/10.3389/fmed.2025.1679733>.
- Yang, L.Y., Fu, J., Peng, X.F., Pang, S.Y., Gao, K.K., Chen, Z.R., He, L.J., *et al.* (2015). Validation of aspartate aminotransferase to platelet ratio for diagnosis of liver fibrosis and prediction of postoperative prognosis in infants with biliary atresia, *World Journal of Gastroenterology*, 21(19), pp. 5893–5900. <https://doi.org/10.3748/wjg.v21.i19.5893>.