

## DAFTAR PUSTAKA

- Admon, A. J., Bohnert, A. S. B., Cooke, C. R., & Taylor, S. P. (2022). Beyond Confounding: Identifying Selection Bias in Observational Pulmonary and Critical Care Research. *Annals of the American Thoracic Society*, 19(7), 1084–1089. <https://doi.org/10.1513/AnnalsATS.202110-1188PS>
- Aduen, J., Bernstein, W. K., Miller, J., Kerzner, R., Bhatiani, A., Davison, L., & Chernow, B. (1995). Relationship between blood lactate concentrations and ionized calcium, glucose, and acid-base status in critically ill and noncritically ill patients: *Critical Care Medicine*, 23(2), 246–252. <https://doi.org/10.1097/00003246-199502000-00008>
- Agrafiotis, M., Keklikoglou, I., Papoti, S., Diminikos, G., Diplaris, K., & Michaelidis, V. (2014). Effect of the Independent Acid Base Variables on Anion Gap Variation in Cardiac Surgical Patients: A Stewart-Figge Approach. *The Scientific World Journal*, 2014, 1–6. <https://doi.org/10.1155/2014/907521>
- Agrafiotis, M., Mpliamplias, D., Papathanassiou, M., Ampatzidou, F., & Drossos, G. (2018). Comparison of a new simplified acid–base tool to the original Stewart–Figge approach: A study on cardiac surgical patients. *Journal of Anesthesia*, 32(4), 499–505. <https://doi.org/10.1007/s00540-018-2503-y>
- Akamatsu, T., Inata, Y., Tachibana, K., Hatachi, T., & Takeuchi, M. (2017). Elevated Central Venous to Arterial Co 2 Difference Is Not Associated With Poor Clinical Outcomes After Cardiac Surgery With Cardiopulmonary Bypass in Children. *Pediatric Critical Care Medicine*, 18(9), 859–862. <https://doi.org/10.1097/pcc.0000000000001229>
- Al Duhailib, Z., Hegazy, A. F., Lalli, R., Fiorini, K., Priestap, F., Iansavichene, A., & Slessarev, M. (2020). The Use of Central Venous to Arterial Carbon Dioxide Tension Gap for Outcome Prediction in Critically Ill Patients: A Systematic Review and Meta-Analysis\*. *Critical Care Medicine*, 48(12), 1855–1861. <https://doi.org/10.1097/ccm.0000000000004578>
- Algarni, K. D. (2020). The effect of hyperlactatemia timing on the outcomes after cardiac surgery. *The Cardiothoracic Surgeon*, 28(1), 18. <https://doi.org/10.1186/s43057-020-00029-w>
- Allen, M. (2011). Lactate and acid base as a hemodynamic monitor and markers of cellular perfusion: *Pediatric Critical Care Medicine*, 12, S43–S49. <https://doi.org/10.1097/PCC.0b013e3182211aed>
- Altun, H. I., Altun, G., Altas, O. F., & Aran, G. (2023). Prognostic Significance of the Strong Ion Gap in Patients in Medical and Surgical Intensive Care Units. *Cureus*, 15(10), e47964. <https://doi.org/10.7759/cureus.47964>
- Åmark, K., Berggren, H., Björk, K., Ekroth, A., Ekroth, R., Nilsson, K., & Sunnegårdh, J. (2005). Blood Cardioplegia Provides Superior Protection in Infant Cardiac Surgery. *The Annals of Thoracic Surgery*, 80(3), 989–994. <https://doi.org/10.1016/j.athoracsur.2005.03.095>

- Bakker, J., Nijsten, M. W., & Jansen, T. C. (2013). Clinical use of lactate monitoring in critically ill patients. *Annals of Intensive Care*, 3(1), 12. <https://doi.org/10.1186/2110-5820-3-12>
- Bakker, J., Vincent, J.-L., Gris, P., Leon, M., Coffernils, M., & Kahn, R. J. (1992). Venous-arterial Carbon Dioxide Gradient in Human Septic Shock. *Chest*, 101(2), 509–515. <https://doi.org/10.1378/chest.101.2.509>
- Balasubramanian, N., Havens, P. L., & Hoffman, G. M. (1999). Unmeasured anions identified by the Fencl-Stewart method predict mortality better than base excess, anion gap, and lactate in patients in the pediatric intensive care unit: *Critical Care Medicine*, 27(8), 1577–1581. <https://doi.org/10.1097/00003246-199908000-00030>
- Balzer, F., Sander, M., Simon, M., Spies, C., Habicher, M., Treskatsch, S., Mezger, V., Schirmer, U., Heringlake, M., Wernecke, K.-D., Grubitzsch, H., & von Heymann, C. (2015). High central venous saturation after cardiac surgery is associated with increased organ failure and long-term mortality: An observational cross-sectional study. *Critical Care (London, England)*, 19(1), 168. <https://doi.org/10.1186/s13054-015-0889-6>
- Baraldi, A. N., & Enders, C. K. (2010). An introduction to modern missing data analyses. *Journal of School Psychology*, 48(1), 5–37. <https://doi.org/10.1016/j.jsp.2009.10.001>
- Bartels, K., Shaw, A. D., Fox, A., Thiel, R. H., & Howard-Quijano, K. (Eds.). (2025). *Hensley's practical approach to cardiothoracic anaesthesia* (Seventh edition). Wolters Kluwer Health.
- Belza, C., Szentkúti, P., Horváth-Puhó, E., Ray, J. G., Nelson, K. E., Grandi, S. M., Brown, H. K., Sørensen, H. T., & Cohen, E. (2024). Use of Latent Class Analysis to Predict Intensive Care Unit Admission and Mortality in Children with a Major Congenital Anomaly. *The Journal of Pediatrics*, 270, 114013. <https://doi.org/10.1016/j.jpeds.2024.114013>
- Brill, S. A., Stewart, T. R., Brundage, S. I., & Schreiber, M. A. (2002). Base Deficit Does Not Predict Mortality When Secondary to Hyperchloremic Acidosis: *Shock*, 17(6), 459–462. <https://doi.org/10.1097/00024382-200206000-00003>
- Buuren, S. V., & Groothuis-Oudshoorn, K. (2011). **mice**: Multivariate Imputation by Chained Equations in R. *Journal of Statistical Software*, 45(3). <https://doi.org/10.18637/jss.v045.i03>
- Castañuela-Sánchez, V., García-Benítez, L., Hernández-Suárez, A., Díaz-García, L., Zamora-Arámburo, M., Sánchez-Cervantes, A., & Palacios-Macedo-Quenot, A. (2023). Central venous-to-arterial CO<sub>2</sub> difference as a biomarker of outcome in children who underwent surgery for congenital heart disease. *Boletín Médico del Hospital Infantil de México*, 80(4). <https://doi.org/10.24875/bmhim.23000066>
- Chen, T., Pan, T., Luo, X., & Wang, D. (2019). High Central Venous-to-Arterial CO<sub>2</sub> Difference is Associated With Poor Outcomes in Patients After Cardiac Surgery: A Propensity Score Analysis. *Shock*, 52(6), 583–589. <https://doi.org/10.1097/shk.0000000000001324>

- Cheung, P.-Y., Chui, N., Joffe, A. R., Rebeyka, I. M., & Robertson, C. M. T. (2005). Postoperative lactate concentrations predict the outcome of infants aged 6 weeks or less after intracardiac surgery: A cohort follow-up to 18 months. *The Journal of Thoracic and Cardiovascular Surgery*, *130*(3), 837–843. <https://doi.org/10.1016/j.jtcvs.2005.04.029>
- Cusack, R., Rhodes, A., Lochhead, P., Jordan, B., Perry, S., Ball, J., Grounds, R., & Bennett, E. (2002). The strong ion gap does not have prognostic value in critically ill patients in a mixed medical/surgical adult ICU. *Intensive Care Medicine*, *28*(7), 864–869. <https://doi.org/10.1007/s00134-002-1318-2>
- Dubin, A., Ferrara, G., Kanoore Edul, V. S., Martins, E., Canales, H. S., Canullán, C., Murias, G., Pozo, M. O., & Estenssoro, E. (2017). Venoarterial PCO<sub>2</sub>-to-arteriovenous oxygen content difference ratio is a poor surrogate for anaerobic metabolism in hemodilution: An experimental study. *Annals of Intensive Care*, *7*(1), 65. <https://doi.org/10.1186/s13613-017-0288-z>
- Dubin, A., Loudet, C. I., Hurtado, F. J., Pozo, M. O., Comande, D., Gibbons, L., Cairoli, F. R., & Bardach, A. (2022). Comparação da proporção entre pressão venosa central menos arterial de dióxido de carbono e conteúdo de oxigênio arterial menos venoso central e níveis de lactato como preditores de mortalidade em pacientes críticos: Uma revisão sistemática e metanálise. *Revista Brasileira de Terapia Intensiva*, *34*(2). <https://doi.org/10.5935/0103-507X.20220026-pt>
- Dungan, K. M., Braithwaite, S. S., & Preiser, J.-C. (2009). Stress hyperglycaemia. *The Lancet*, *373*(9677), 1798–1807. [https://doi.org/10.1016/S0140-6736\(09\)60553-5](https://doi.org/10.1016/S0140-6736(09)60553-5)
- Durward, A., Tibby, S. M., Skellett, S., Austin, C., Anderson, D., & Murdoch, I. A. (2005). The strong ion gap predicts mortality in children following cardiopulmonary bypass surgery. *Pediatric Critical Care Medicine*, *6*(3), 281–285. <https://doi.org/10.1097/01.PCC.0000163979.33774.89>
- Enders, C., & Bandalos, D. (2001). The Relative Performance of Full Information Maximum Likelihood Estimation for Missing Data in Structural Equation Models. *Structural Equation Modeling: A Multidisciplinary Journal*, *8*(3), 430–457. [https://doi.org/10.1207/S15328007SEM0803\\_5](https://doi.org/10.1207/S15328007SEM0803_5)
- Fencl, V., Jabor, A., Kazda, A., & Figge, J. (2000). Diagnosis of Metabolic Acid–Base Disturbances in Critically Ill Patients. *American Journal of Respiratory and Critical Care Medicine*, *162*(6), 2246–2251. <https://doi.org/10.1164/ajrccm.162.6.9904099>
- Finch, W. H., & Bronk, K. C. (2011a). Conducting Confirmatory Latent Class Analysis Using M *plus*. *Structural Equation Modeling: A Multidisciplinary Journal*, *18*(1), 132–151. <https://doi.org/10.1080/10705511.2011.532732>
- Finch, W. H., & Bronk, K. C. (2011b). Conducting Confirmatory Latent Class Analysis Using M *plus*. *Structural Equation Modeling: A Multidisciplinary Journal*, *18*(1), 132–151. <https://doi.org/10.1080/10705511.2011.532732>
- Gavelli, F., Teboul, J.-L., & Monnet, X. (2019). How can CO<sub>2</sub>-derived indices guide resuscitation in critically ill patients? *Journal of Thoracic Disease*, *11*(S11), S1528–S1537. <https://doi.org/10.21037/jtd.2019.07.10>

- Gudicha, D. W., Tekle, F. B., & Vermunt, J. K. (2016). Power and Sample Size Computation for Wald Tests in Latent Class Models. *Journal of Classification*, 33(1), 30–51. <https://doi.org/10.1007/s00357-016-9199-1>
- Guinot, P.-G., Badoux, L., Bernard, E., Abou-Arab, O., Lorne, E., & Dupont, H. (2017). Central Venous-to-Arterial Carbon Dioxide Partial Pressure Difference in Patients Undergoing Cardiac Surgery is Not Related to Postoperative Outcomes. *Journal of Cardiothoracic and Vascular Anesthesia*, 31(4), 1190–1196. <https://doi.org/10.1053/j.jvca.2017.02.015>
- Gunnerson, K. J., Srisawat, N., & Kellum, J. A. (2010). Is there a difference between strong ion gap in healthy volunteers and intensive care unit patients? *Journal of Critical Care*, 25(3), 520–524. <https://doi.org/10.1016/j.jcrc.2009.11.001>
- Gunnerson, K., Saul, M., He, S., & Kellum, J. (2006). Lactate versus non-lactate metabolic acidosis: A retrospective outcome evaluation of critically ill patients. *Critical Care*, 10(1), R22. <https://doi.org/10.1186/cc3987>
- Habicher, M., Von Heymann, C., Spies, C. D., Wernecke, K.-D., & Sander, M. (2015). Central Venous-Arterial pCO<sub>2</sub> Difference Identifies Microcirculatory Hypoperfusion in Cardiac Surgical Patients With Normal Central Venous Oxygen Saturation: A Retrospective Analysis. *Journal of Cardiothoracic and Vascular Anesthesia*, 29(3), 646–655. <https://doi.org/10.1053/j.jvca.2014.09.006>
- Hajjar, L. A., Almeida, J. P., Fukushima, J. T., Rhodes, A., Vincent, J.-L., Osawa, E. A., & Galas, F. R. B. G. (2013). High lactate levels are predictors of major complications after cardiac surgery. *The Journal of Thoracic and Cardiovascular Surgery*, 146(2), 455–460. <https://doi.org/10.1016/j.jtcvs.2013.02.003>
- Hatherill, M. (2005). Hyperchloraemic metabolic acidosis following open cardiac surgery. *Archives of Disease in Childhood*, 90(12), 1288–1292. <https://doi.org/10.1136/adc.2005.078006>
- Hayhoe, M., Bellomo, R., Liu, G., McNicol, L., & Buxton, B. (1999). The aetiology and pathogenesis of cardiopulmonary bypass-associated metabolic acidosis using polygeline pump prime. *Intensive Care Medicine*, 25(7), 680–685. <https://doi.org/10.1007/s001340050930>
- He, Q., Tan, Z., Chen, D., Cai, S., & Zhou, L. (2023). Association between intraoperative hyperglycemia/hyperlactatemia and acute kidney injury following on-pump cardiac surgery: A retrospective cohort study. *Frontiers in Cardiovascular Medicine*, 10, 1218127. <https://doi.org/10.3389/fcvm.2023.1218127>
- Henson, J. M., Reise, S. P., & Kim, K. H. (2007). Detecting Mixtures From Structural Model Differences Using Latent Variable Mixture Modeling: A Comparison of Relative Model Fit Statistics. *Structural Equation Modeling: A Multidisciplinary Journal*, 14(2), 202–226. <https://doi.org/10.1080/10705510709336744>
- Ho, K. M., Lan, N. S. H., Williams, T. A., Harahsheh, Y., Chapman, A. R., Dobb, G. J., & Magder, S. (2016). A comparison of prognostic significance of strong ion gap (SIG) with other acid-base markers in the critically ill: A

- cohort study. *Journal of Intensive Care*, 4(1), 43.  
<https://doi.org/10.1186/s40560-016-0166-z>
- Hosein, R. B. M., Morris, K. P., Brawn, W. J., & Barron, D. J. (2008). Use of tissue microdialysis to investigate hyperlactataemia following paediatric cardiac surgery. *Interactive Cardiovascular and Thoracic Surgery*, 7(3), 384–388. <https://doi.org/10.1510/icvts.2007.166264>
- Huette, P., Beyls, C., Mallat, J., Martineau, L., Besserve, P., Haye, G., Guilbart, M., Dupont, H., Guinot, P.-G., Diouf, M., Mahjoub, Y., & Abou-Arab, O. (2021). Central venous-to-arterial CO<sub>2</sub> difference is a poor tool to predict adverse outcomes after cardiac surgery: A retrospective study. *Canadian Journal of Anesthesia/Journal Canadien d'anesthésie*, 68(4), 467–476. <https://doi.org/10.1007/s12630-020-01881-4>
- Ince, C. (2015). Hemodynamic coherence and the rationale for monitoring the microcirculation. *Critical Care*, 19(S3), S8. <https://doi.org/10.1186/cc14726>
- Insom, G., Marinari, E., Scolari, A. F., Garisto, C., Vitale, V., Di Chiara, L., & Ricci, Z. (2021). Venous-arterial CO<sub>2</sub> difference and cardiac index in children after cardiac surgery. *Cardiology in the Young*, 31(4), 597–601. <https://doi.org/10.1017/s1047951120004357>
- Kaakinen, T. I., Ikäläinen, T., Erkinaro, T. M., Karhu, J. M., Liisanantti, J. H., Ohtonen, P. P., & Ala-Kokko, T. I. (2022). Association of low mixed venous oxygen saturations during early ICU stay with increased 30-day and 1-year mortality after cardiac surgery: A single-center retrospective study. *BMC Anesthesiology*, 22(1), 322. <https://doi.org/10.1186/s12871-022-01862-8>
- Klee, P., Rimensberger, P. C., & Karam, O. (2020). Association Between Lactates, Blood Glucose, and Systemic Oxygen Delivery in Children After Cardiopulmonary Bypass. *Frontiers in Pediatrics*, 8, 332. <https://doi.org/10.3389/fped.2020.00332>
- Kobayashi, Y., Peng, Y.-C., Yu, E., Bush, B., Jung, Y.-H., Murphy, Z., Goeddel, L., Whitman, G., Venkataraman, A., & Brown, C. H. (2023). Prediction of lactate concentrations after cardiac surgery using machine learning and deep learning approaches. *Frontiers in Medicine*, 10, 1165912. <https://doi.org/10.3389/fmed.2023.1165912>
- Kogan, A., Preisman, S., Bar, A., Sternik, L., Lavee, J., Malachy, A., Spiegelstein, D., Berkenstadt, H., & Raanani, E. (2012). The impact of hyperlactatemia on postoperative outcome after adult cardiac surgery. *Journal of Anesthesia*, 26(2), 174–178. <https://doi.org/10.1007/s00540-011-1287-0>
- Koning, N. J., Simon, L. E., Asfar, P., Baufreton, C., & Boer, C. (2014). Systemic microvascular shunting through hyperdynamic capillaries after acute physiological disturbances following cardiopulmonary bypass. *American Journal of Physiology-Heart and Circulatory Physiology*, 307(7), H967–H975. <https://doi.org/10.1152/ajpheart.00397.2014>
- Kurniawaty, J., Setianto, B. Y., Widyastuti, Y., Supomo, S., Boom, C. E., & Ancilla, C. (2022). Validation for EuroSCORE II in the Indonesian cardiac surgical population: A retrospective, multicenter study. *Expert Review of Cardiovascular Therapy*, 20(6), 491–496. <https://doi.org/10.1080/14779072.2022.2078703>

- Laine, G. A., Hu, B. Y., Wang, S., Thomas Solis, R., & Reul, G. J. (2013). Isolated High Lactate or Low Central Venous Oxygen Saturation After Cardiac Surgery and Association With Outcome. *Journal of Cardiothoracic and Vascular Anesthesia*, 27(6), 1271–1276. <https://doi.org/10.1053/j.jvca.2013.02.031>
- Lanza, S. T., & Rhoades, B. L. (2013). Latent Class Analysis: An Alternative Perspective on Subgroup Analysis in Prevention and Treatment. *Prevention Science*, 14(2), 157–168. <https://doi.org/10.1007/s11121-011-0201-1>
- Li, J., Tian, Y., Wang, L., Chen, J., Chen, X., Huang, H., & Li, Y. (2022). Postoperative anion gap associates with short- and long-term mortality after cardiac surgery: A large-scale cohort study. *Frontiers in Cardiovascular Medicine*, 9, 1024484. <https://doi.org/10.3389/fcvm.2022.1024484>
- Liao, Y.-C., Chang, C.-C., Chen, C.-Y., Liu, C.-C., Liao, C.-C., Shih, Y.-R. V., & Lin, C.-S. (2023). Preoperative renal insufficiency predicts postoperative adverse outcomes in a mixed surgical population: A retrospective matched cohort study using the NSQIP database. *International Journal of Surgery*, 109(4), 752–759. <https://doi.org/10.1097/JS9.0000000000000278>
- Lo, Y. (2001). Testing the number of components in a normal mixture. *Biometrika*, 88(3), 767–778. <https://doi.org/10.1093/biomet/88.3.767>
- Maciel, A. T., & Park, M. (2007). UNMEASURED ANIONS ACCOUNT FOR MOST OF THE METABOLIC ACIDOSIS IN PATIENTS WITH HYPERLACTATEMIA. *Clinics*, 62(1), 55–62. <https://doi.org/10.1590/S1807-59322007000100009>
- Maillet, J.-M., Le Besnerais, P., Cantoni, M., Nataf, P., Ruffenach, A., Lessana, A., & Brodaty, D. (2003). Frequency, Risk Factors, and Outcome of Hyperlactatemia After Cardiac Surgery. *Chest*, 123(5), 1361–1366. <https://doi.org/10.1378/chest.123.5.1361>
- Mallat, J., Lemyze, M., Meddour, M., Pepy, F., Gasan, G., Barrailler, S., Durville, E., Temime, J., Vangrunderbeek, N., Tronchon, L., Vallet, B., & Thevenin, D. (2016). Ratios of central venous-to-arterial carbon dioxide content or tension to arteriovenous oxygen content are better markers of global anaerobic metabolism than lactate in septic shock patients. *Annals of Intensive Care*, 6(1), 10. <https://doi.org/10.1186/s13613-016-0110-3>
- Marik, P. E., & Bellomo, R. (2013). Stress hyperglycemia: An essential survival response! *Critical Care*, 17(2), 305. <https://doi.org/10.1186/cc12514>
- Martin, M. J., FitzSullivan, E., Salim, A., Brown, C. V. R., Demetriades, D., & Long, W. (2006). Discordance between lactate and base deficit in the surgical intensive care unit: Which one do you trust? *The American Journal of Surgery*, 191(5), 625–630. <https://doi.org/10.1016/j.amjsurg.2006.02.014>
- Matsumoto, S., Omiya, H., Fujinaka, W., & Morimatsu, H. (2022). Association between intraoperative hyperglycemia and postoperative end-organ dysfunctions after cardiac surgery: A retrospective observational study. *Journal of Anesthesia*, 36(2), 174–184. <https://doi.org/10.1007/s00540-021-03024-5>
- Matteucci, M., Ferrarese, S., Cantore, C., Cappabianca, G., Massimi, G., Mantovani, V., Rossi, M. B., & Beghi, C. (2020). Hyperlactatemia during

cardiopulmonary bypass: Risk factors and impact on surgical results with a focus on the long-term outcome. *Perfusion*, 35(8), 756–762.  
<https://doi.org/10.1177/0267659120907440>

- Mecher, C. E., Rackow, E. C., Astiz, M. E., & Weil, M. H. (1990). Venous hypercarbia associated with severe sepsis and systemic hypoperfusion. *Critical Care Medicine*, 18(6), 585–589. <https://doi.org/10.1097/00003246-199006000-00001>
- Miao, Q., Wu, D. J., Chen, X., Xu, M., Sun, L., Guo, Z., He, B., & Wu, J. (2021). Target blood pressure management during cardiopulmonary bypass improves lactate levels after cardiac surgery: A randomized controlled trial. *BMC Anesthesiology*, 21(1), 309. <https://doi.org/10.1186/s12871-021-01537-w>
- Minton, J., & Sidebotham, D. A. (2017). Hyperlactatemia and Cardiac Surgery. *The Journal of Extra-Corporeal Technology*, 49(1), 7–15.
- Miranda, C. D. A., Meletti, J. F. A., Lima, L. H. N., & Marchi, E. (2020). Correlação entre saturação venosa central de oxigênio perioperatória e mortalidade em cirurgia cardíaca: Estudo prospectivo observacional. *Brazilian Journal of Anesthesiology*, 70(5), 484–490. <https://doi.org/10.1016/j.bjan.2020.04.014>
- Moons, K. G. M., de Groot, J. A. H., Bouwmeester, W., Vergouwe, Y., Mallett, S., Altman, D. G., Reitsma, J. B., & Collins, G. S. (2014). Critical appraisal and data extraction for systematic reviews of prediction modelling studies: The CHARMS checklist. *PLoS Medicine*, 11(10), e1001744. <https://doi.org/10.1371/journal.pmed.1001744>
- Morgan, T. J., & Hall, J. A. (1999). Hyperlactaemia without acidosis—An investigation using an in vitro model. *Critical Care and Resuscitation: Journal of the Australasian Academy of Critical Care Medicine*, 1(4), 354–359.
- Murray, D., Grant, D., Murali, N., & Butt, W. (2007). Unmeasured anions in children after cardiac surgery. *The Journal of Thoracic and Cardiovascular Surgery*, 133(1), 235–240. <https://doi.org/10.1016/j.jtcvs.2006.09.017>
- Murray, D. M., Olhsson, V., & Fraser, J. I. (2004). Defining acidosis in postoperative cardiac patients using Stewart's method of strong ion difference\*. *Pediatric Critical Care Medicine*, 5(3), 240–245. <https://doi.org/10.1097/01.pcc.0000112367.50051.3b>
- Nahas, R. W. (2024). *Introduction to Regression Methods for Public Health Using R* (1st ed.). Chapman and Hall/CRC. <https://doi.org/10.1201/9781003263197>
- Naik, R., George, G., Karuppiah, S., & Philip, M. (2016). Hyperlactatemia in patients undergoing adult cardiac surgery under cardiopulmonary bypass: Causative factors and its effect on surgical outcome. *Annals of Cardiac Anaesthesia*, 19(4), 668. <https://doi.org/10.4103/0971-9784.191579>
- Nashef, S. A. M., Roques, F., Sharples, L. D., Nilsson, J., Smith, C., Goldstone, A. R., & Lockowandt, U. (2012). EuroSCORE II. *European Journal of Cardio-Thoracic Surgery*, 41(4), 734–745. <https://doi.org/10.1093/ejcts/ezs043>

- Nogueira, P. M., Mendonça-Filho, H. T., Campos, L. A., Gomes, R. V., Felipe, A. R., Fernandes, M. A., Villela-Nogueira, C. A., & Rocco, J. R. (2010). Central Venous Saturation: A Prognostic Tool in Cardiac Surgery patients. *Journal of Intensive Care Medicine*, 25(2), 111–116. <https://doi.org/10.1177/0885066609355398>
- Nylund, K. L., Asparouhov, T., & Muthén, B. O. (2007). Deciding on the Number of Classes in Latent Class Analysis and Growth Mixture Modeling: A Monte Carlo Simulation Study. *Structural Equation Modeling: A Multidisciplinary Journal*, 14(4), 535–569. <https://doi.org/10.1080/10705510701575396>
- O'Brien, S. M., Feng, L., He, X., Xian, Y., Jacobs, J. P., Badhwar, V., Kurlansky, P. A., Furnary, A. P., Cleveland, J. C., Lobdell, K. W., Vassileva, C., Wyler Von Ballmoos, M. C., Thourani, V. H., Rankin, J. S., Edgerton, J. R., D'Agostino, R. S., Desai, N. D., Edwards, F. H., & Shahian, D. M. (2018). The Society of Thoracic Surgeons 2018 Adult Cardiac Surgery Risk Models: Part 2—Statistical Methods and Results. *The Annals of Thoracic Surgery*, 105(5), 1419–1428. <https://doi.org/10.1016/j.athoracsur.2018.03.003>
- O'Connor, E., & Fraser, J. F. (2012). The Interpretation of Perioperative Lactate Abnormalities in Patients Undergoing Cardiac Surgery. *Anaesthesia and Intensive Care*, 40(4), 598–603. <https://doi.org/10.1177/0310057X1204000404>
- Palermo, R. A., Palac, H. L., Wald, E. L., Wainwright, M. S., Costello, J. M., Eltayeb, O. M., Backer, C. L., & Epting, C. L. (2015). Metabolic Uncoupling Following Cardiopulmonary Bypass: Elevated Lactate and Glucose after CPB. *Congenital Heart Disease*, 10(6), E250–E257. <https://doi.org/10.1111/chd.12285>
- Perz, S., Uhlig, T., Kohl, M., Bredle, D. L., Reinhart, K., Bauer, M., & Kortgen, A. (2011). Low and “supranormal” central venous oxygen saturation and markers of tissue hypoxia in cardiac surgery patients: A prospective observational study. *Intensive Care Medicine*, 37(1), 52–59. <https://doi.org/10.1007/s00134-010-1980-8>
- Ranucci, M., De Toffol, B., Isgro, G., Romitti, F., Conti, D., & Vicentini, M. (2006). Hyperlactatemia during cardiopulmonary bypass: Determinants and impact on postoperative outcome. *Critical Care*, 10(6), R167. <https://doi.org/10.1186/cc5113>
- Raper, R. F., Cameron, G., Walker, D., & Bowey, C. J. (1997). Type B lactic acidosis following cardiopulmonary bypass: *Critical Care Medicine*, 25(1), 46–51. <https://doi.org/10.1097/00003246-199701000-00011>
- Renew, J. R., Barbara, D. W., Hyder, J. A., Dearani, J. A., Rivera, M., & Pulido, J. N. (2016). Frequency and outcomes of severe hyperlactatemia after elective cardiac surgery. *The Journal of Thoracic and Cardiovascular Surgery*, 151(3), 825–830. <https://doi.org/10.1016/j.jtcvs.2015.10.063>
- Rhodes, L. A., Erwin, W. C., Borasino, S., Cleveland, D. C., & Alten, J. A. (2017). Central Venous to Arterial Co 2 Difference After Cardiac Surgery in Infants and Neonates\*. *Pediatric Critical Care Medicine*, 18(3), 228–233. <https://doi.org/10.1097/pcc.0000000000001085>

- Rivers, E., Nguyen, B., Havstad, S., Ressler, J., Muzzin, A., Knoblich, B., Peterson, E., & Tomlanovich, M. (2001). Early Goal-Directed Therapy in the Treatment of Severe Sepsis and Septic Shock. *New England Journal of Medicine*, 345(19), 1368–1377. <https://doi.org/10.1056/NEJMoa010307>
- Robertson, J., & Kaptein, M. (Eds.). (2016). *Modern Statistical Methods for HCI*. Springer International Publishing. <https://doi.org/10.1007/978-3-319-26633-6>
- Robin, E., Costecalde, M., Lebuffe, G., & Vallet, B. (2006). Clinical relevance of data from the pulmonary artery catheter. *Critical Care*, 10(Suppl 3), S3. <https://doi.org/10.1186/cc4830>
- Rocktaeschel, J., Morimatsu, H., Uchino, S., & Bellomo, R. (2003). Unmeasured anions in critically ill patients: Can they predict mortality?\*: *Critical Care Medicine*, 31(8), 2131–2136. <https://doi.org/10.1097/01.CCM.0000079819.27515.8E>
- Seghrouchni, A., Atmani, N., Moutakiallah, Y., Belmekki, A., El Bekkali, Y., & Houssa, M. A. (2022). Does severe hyperlactatemia during cardiopulmonary bypass predict a worse outcome? *Annals of Medicine & Surgery*, 73. <https://doi.org/10.1016/j.amsu.2021.103198>
- Sigwadhi, L. N., Tamuzi, J. L., Zemlin, A. E., Chapanduka, Z. C., Allwood, B. W., Koegelenberg, C. F., Irusen, E. M., Lalla, U., Ngah, V. D., Yalew, A., Savieri, P., Fwemba, I., Jalavu, T. P., Erasmus, R. T., Matsha, T. E., Zumla, A., & Nyasulu, P. S. (2022). Latent class analysis: An innovative approach for identification of clinical and laboratory markers of disease severity among COVID-19 patients admitted to the intensive care unit. *IJID Regions*, 5, 154–162. <https://doi.org/10.1016/j.ijregi.2022.10.004>
- Sinha, P., Calfee, C. S., & Delucchi, K. L. (2021). Practitioner’s Guide to Latent Class Analysis: Methodological Considerations and Common Pitfalls. *Critical Care Medicine*, 49(1), e63–e79. <https://doi.org/10.1097/CCM.0000000000004710>
- Siuba, M. T., Bhardwaj, A., Kirincich, J., Perez, O., Flanagan, P., Lane, J., Toth, D., Paul, D., Lehr, C., Duggal, A., & Tonelli, A. R. (2023). Does veno-arterial carbon dioxide gradient provide an adequate estimation of cardiac index in pulmonary hypertension? *European Heart Journal. Acute Cardiovascular Care*, 12(1), 38–47. <https://doi.org/10.1093/ehjacc/zuac139>
- Sterba, S. K. (2016). Cautions on the Use of Multiple Imputation When Selecting Between Latent Categorical versus Continuous Models for Psychological Constructs. *Journal of Clinical Child & Adolescent Psychology*, 45(2), 167–175. <https://doi.org/10.1080/15374416.2014.958839>
- Surgical and Clinical Outcome REsearch (SCORE) Group, Ranucci, M., Isgrò, G., Carlucci, C., De La Torre, T., Enginoli, S., & Frigiola, A. (2010). Central venous oxygen saturation and blood lactate levels during cardiopulmonary bypass are associated with outcome after pediatric cardiac surgery. *Critical Care*, 14(4), R149. <https://doi.org/10.1186/cc9217>
- Teboul, J.-L., & Scheeren, T. (2017). Understanding the Haldane effect. *Intensive Care Medicine*, 43(1), 91–93. <https://doi.org/10.1007/s00134-016-4261-3>

- Textoris, J., Fouché, L., Wiramus, S., Antonini, F., Tho, S., Martin, C., & Leone, M. (2011). High central venous oxygen saturation in the latter stages of septic shock is associated with increased mortality. *Critical Care*, *15*(4), R176. <https://doi.org/10.1186/cc10325>
- Tönnies, T., Kahl, S., & Kuss, O. (2022). Collider bias in observational studies: Consequences for medical research. Part 30 of a series on evaluation of scientific publications. *Deutsches Ärzteblatt International*. <https://doi.org/10.3238/arztebl.m2022.0076>
- Vallet, B., Teboul, J.-L., Cain, S., & Curtis, S. (2000). Venoarterial CO<sub>2</sub> difference during regional ischemic or hypoxic hypoxia. *Journal of Applied Physiology*, *89*(4), 1317–1321. <https://doi.org/10.1152/jappl.2000.89.4.1317>
- Van Der Linden, P., Rausin, I., Deltell, A., Bekrar, Y., Gilbert, E., Bakker, J., & Vincent, J.-L. (1995). Detection of Tissue Hypoxia by Arteriovenous Gradient for PCO<sub>2</sub> and pH in Anesthetized Dogs During Progressive Hemorrhage. *Anesthesia & Analgesia*, *80*(2), 269–275. <https://doi.org/10.1097/00000539-199502000-00012>
- Wendel Garcia, P. D., Caccioppola, A., Coppola, S., Pozzi, T., Ciabattoni, A., Cenci, S., & Chiumello, D. (2021). Latent class analysis to predict intensive care outcomes in Acute Respiratory Distress Syndrome: A proposal of two pulmonary phenotypes. *Critical Care*, *25*(1), 154. <https://doi.org/10.1186/s13054-021-03578-6>
- White, I. R., Royston, P., & Wood, A. M. (2011). Multiple imputation using chained equations: Issues and guidance for practice. *Statistics in Medicine*, *30*(4), 377–399. <https://doi.org/10.1002/sim.4067>
- Zakkar, M., Guida, G., Suleiman, M.-S., & Angelini, G. D. (2015). Cardiopulmonary Bypass and Oxidative Stress. *Oxidative Medicine and Cellular Longevity*, *2015*, 1–8. <https://doi.org/10.1155/2015/189863>
- Zante, B., & Kluge, S. (2020). Central Venous-Arterial CO<sub>2</sub> Difference in Cardiac Surgery Patients—A Parameter in Relationship to Cardiac Output and Altered Microcirculatory Blood Flow. *Shock*, *54*(2), 273–274. <https://doi.org/10.1097/shk.0000000000001525>
- Zante, B., Reichenspurner, H., Kubik, M., Schefold, J. C., & Kluge, S. (2019). Increased admission central venous-arterial CO<sub>2</sub> difference predicts ICU-mortality in adult cardiac surgery patients. *Heart & Lung*, *48*(5), 421–427. <https://doi.org/10.1016/j.hrtlng.2019.05.015>