

DAFTAR PUSTAKA

- Al-Salem, A.H. (2007) "Congenital intrinsic duodenal obstruction: A review of 35 cases," *Annals of Saudi Medicine*, 27(4), pp. 289–292.
- Applebaum, H. and Sydorak, R. (2012) "Duodenal Atresia and Stenosis-Annular Pancreas," in A.G. Coran et al. (eds) *Pediatric surgery*. 7th edn. Philadelphia, PA: Elsevier Mosby, pp. 1051–1058.
- Battersby C, Statnikov Y, Santhakumaran S, Gray D, Modi N. The effect of birth weight on neonatal mortality: systematic review and meta-analysis. *Arch Dis Child Fetal Neonatal Ed*. 2018;103(3):F275–F281
- Bethell, G., Long, A., Knight, M., & Hall, N. (2023). Factors associated with outcomes in congenital duodenal obstruction: population-based study. *BJS*, 110(9), 1053–1056. <https://doi.org/https://doi.org/10.1093/bjs/znad040>
- Cairo SB, Tabak BD, Berman L, Berkelhamer SK, Yu G, Rothstein DH. Mortality after emergency abdominal operations in premature infants. *J Pediatr Surg* 2018; <https://doi.org/10.1016/j.jpedsurg.2018.01.009>.
- Chandrasekaran, S., & Asokaraju, A. (2017). Clinical profile and predictors of outcome in congenital duodenal obstruction. *International Surgery Journal*, 4(8), 2605–2611. <https://doi.org/10.18203/2349-2902.isj20173396>
- Chiarenza SF, Bucci V, Conighi ML, Zolpi E, Costa L, Fasoli L, Bleve C. Duodenal atresia: open versus MIS repair analysis of our experience over the last 12 years. *BioMed Res Int* 2017; 2017:1e7. <https://doi.org/10.1155/2017/4585360>.
- Choudhry, M.S. et al. (2009) "Duodenal atresia: Associated anomalies, prenatal diagnosis and outcome," *Pediatric Surgery International*, 25(8), pp. 727–730.
- Coran, A.G., Adzick, N.S., Krummel, T.M. and Laberge, J.M., eds., (2012). *Coran's Pediatric Surgery* 7th ed. Philadelphia: Elsevier
- Corebima, Brigitta I. R. V., Sulistijono E, Widanto, Choe, Cheah Fook (2023). Factors associated with infection and mortality in neonates following abdominal surgery. *Paediatr Indones*, Vol.63, No. 01 (Suppl.) (2022). p.S8-S13; DOI: <https://doi.org/10.14238/pi63.01.2022.S8-S13>.
- Dahlan, M. (2015). *Analisis Survival: Dasar-dasar Teori & Aplikasi Dengan Program SPSS*. Jakarta: Epidemiologi Indonesia.
- Deguchi K, Tazuke Y, Matsuura R, Nomura M, Yamanak H, Soh H, Yodena A. Factors associated with adverse outcomes following duodenal atresia surgery

in neonates: a retrospective study. *Cureus* 2022;17.
<https://doi.org/10.7759/cureus.22349>.

Desdwianto D, Hariastawa A, Matulatan F. Prognostic Factors Affecting the Mortality of Neonates with Duodenal Obstruction at RSUD Dr. Soetomo. *J. Med. Chem. Sci.* 2024, 7(4) 579-589. DOI:10.26655/JMCHEMSCI.2024.4.2

Dindo, et al. (2004). "Classification of Surgical Complications a New Proposal with Evaluation in a Cohort of 6336 Patients and Results of a Survey", *Annals of Surgery* 240(2): p205-213, August 2004. DOI: 10.1097/01.sla.0000133083.54934.ae

Escobar, M. A., Ladd, A. P., Grosfeld, J. L., West, K. W., Rescorla, F. J., & Scherer, L. R. (2004). Duodenal atresia and stenosis: Long-term follow-up over 30 years. *Journal of Pediatric Surgery*, 39(6), 867–871. DOI: 10.1016/j.jpedsurg.2004.02.025

Estiarla and Agustriani, N. (2021) "Prognostic factors for mortality in patients with congenital duodenal obstruction at dr. Moewardi Hospital Surakarta," *Precision Medical Sciences*, 10(1), pp. 26–32.

Filan PM, Hunt RW, Anderson PJ, Doyle LW, Inder TE. Neurologic outcomes in very preterm infants undergoing surgery. *J Pediatr* 2012; <https://doi.org/10.1016/j.jpeds.2011.09.009>.

Guyton, A.C. & Hall, J.E., (2020). *Guyton and Hall Textbook of Medical Physiology*. 13th Ed. Philadelphia (PA): Elsevier, Inc.

Holcom, A. & Aschraft, B., (2020). *Pediatric Surgery: Principles and Practice*, ed. by J. Smith, 2nd ed. New York: Springer.

Kammoun, H., Hammami, H., Slim, I., & Ben Rhouma, R. (2023). Predictors of postoperative mortality among neonates after major-risk surgery: A one-year experience from a Tunisian hospital. *Journal of Neonatal Surgery*, 12(1), 10. <https://www.mendeley.com/catalogue/af08548e-057e-3489-9a4e-569c464d2bc4/>

Kimura, K. et al. (1990) "Diamond-shaped anastomosis for duodenal atresia: An experience with 44 patients over 15 years," *Journal of Pediatric Surgery*, 25(9), pp. 977–979.

Kumar, P., kumar, chiranjiv, pandey, prince raj, & Sarin, Y. K. (2020). Congenital Duodenal Obstruction in Neonates: Over 13 Years' Experience from a Single Centre. *Journal of Neonatal Surgery*, 5(4), 50. <https://doi.org/10.52783/jns.v5.313>

- Lin YJ, Chen JC, Peng CC, et al. Outcomes in infants with congenital anomalies undergoing major surgery: a population-based study. *Pediatr Surg Int.* 2020;36(10):1187–94. doi:10.1007/s00383-020-04730-7.
- Liu, X., Shen, Y., Wang, H., Ge, Q., Fei, A., & Pan, S. (2016). Prognostic Significance of Neutrophil-to-Lymphocyte Ratio in Patients with Sepsis: A Prospective Observational Study. *Mediators of Inflammation*, 2016. <https://doi.org/10.1155/2016/8191254>
- Makkadafi, M., Fauzi, A. R., Wandita, S., Makhmudi, A., & Gunadi. (2021). Outcomes and survival of infants with congenital duodenal obstruction following Kimura procedure with post-anastomosis jejunostomy feeding tube. *BMC Gastroenterology*, 21(1). <https://doi.org/10.1186/s12876-021-01679-8>
- Munandar, et al. 2023. Prognostic factors following Kimura surgery in cases of congenital duodenum obstruction toward survival. thesis. Repository Universitas Gadjah Mada.
- Ni, J., Wang, H., Li, Y., Shu, Y., & Liu, Y. (2019). Neutrophil to lymphocyte ratio (NLR) as a prognostic marker for in-hospital mortality of patients with sepsis: A secondary analysis based on a single-center, retrospective, cohort study. *Medicine*, 98(46), e18029. <https://doi.org/10.1097/MD.00000000000018029>
- Ogle, S.B., Nichol, P.F. and Ostlie, D.J. (2020) “Duodenal and Intestinal Atresia and Stenosis,” in G.W. Holcomb et al. (eds) *Holcomb and Ashcraft's pediatric surgery*. 7th edn. Edinburgh: Elsevier, pp. 489–506.
- Prathana, S., Adipurwadi, Y.C. and Agustriani, N. (2021) Rasio Neutrofil Limfosit Pra Pembedahan sebagai Faktor Prognostik Kesintasan Pasca Pembedahan pada Obstruksi Duodenum Kongenital. thesis. Repository Universitas Gadjah Mada.
- Puri, P., O'Donnell, A., O'Connor, A., McDermott, M., Heffernan, A., & Quinn, F. (2019). A pilot study on neonatal surgical mortality: A multivariable analysis of predictors of mortality in a resource-limited setting. *Journal of Neonatal Surgery*, 8(1), 6. <https://pmc.ncbi.nlm.nih.gov/articles/PMC6322181/>
- Qamer, S., Arshad, M., Imran, M., Niaz, S., & Khan, A. (2023). Clinical outcomes and predictors of survival in congenital duodenal obstruction: A retrospective study. *Journal of Pediatric Surgery*, 58(9), 1805–1811. <https://doi.org/10.1016/j.jpedsurg.2023.03.012>
- Raval MV, Dillon PW, Bruny JL, et al. Defining the association between operative time and outcomes in children's surgery. *JAMA Pediatr.* 2017;171(7):e171325. doi:10.1001/jamapediatrics.2017.1325

- Sastiono, S., & Rahmaania, J. C. (2019). The relationship between sepsis and timing achievement of peristaltic function in congenital duodenal obstruction. *The New Ropanasuri Journal of Surgery*, 4(1), 24–26. <https://doi.org/10.7454/nrjs.v4i1.73>
- Sastroasmoro, Sudigdo (2014). *Dasar-Dasar Metodologi Penelitian Klinis*. Edisi kelima. Jakarta: Sagung Seto
- Sharma, P., Pawar, N., Gupta, A.R. (2022). Congenital intrinsic duodenal obstruction: Clinical study and predictors of neonatal surgical mortality. *Journal of Neonatal Surgery*, 9(2), Article 1034. <https://doi.org/10.21699/jns.v9i2.1034>
- Spitz, L. and Coran, A.G., (2013). *Operative Pediatric Surgery*. 7th ed. Philadelphia: CRC Press
- St-Louis, E., Sharif, S., Flageole, H., & Bouchard, S. (2018). Neonatal duodenal obstruction: Predictors of outcome and long-term results. *Journal of Pediatric Surgery*, 53(9), 1808–1813. <https://doi.org/10.1016/j.jpedsurg.2017.10.066>
- Westerdijk, K., Simons, K. S., Zegers, M., Wever, P. C., Pickkers, P., & de Jager C. P. C. (2019). The value of the neutrophil-lymphocyte count ratio in the diagnosis of sepsis in patients admitted to the Intensive Care Unit: A retrospective cohort study. *PLoS ONE*, 14(2), 1–13 <https://doi.org/10.1371/journal.pone.0212861>
- Wilar, R. (2019). Diagnostic value of eosinopenia and neutrophil to lymphocyte ratio on early onset neonatal sepsis. *Korean Journal of Pediatrics*, 62(6)217–223. <https://doi.org/10.3345/kjp.2018.06723>
- Wilar, R. Koesmarsono, B. Gunawan, S. (2023). The combination of neutrophil to lymphocyte ratio and platelet to lymphocyte ratio improves accuracy of neonatal sepsis diagnosis. *Pediatrica Indonesiana*, Vol.63, No.4(2023). p.213-8; DOI: <https://doi.org/10.14238/pi63.4.2023.213-8>
- Zani, A., Eaton, S., Hoellwarth, M. E., et al. (2017). European paediatric surgeons' association survey on the management of congenital duodenal obstruction. *European Journal of Pediatric Surgery*, 27(01), 81–86.