

REFERENCES

- Agard, A. *et al.* (2022) ‘Clinical comparison and agreement of PCR, antigen, and viral culture for the diagnosis of COVID-19: Clinical Agreement Between Diagnostics for COVID-19.’, *Journal of clinical virology plus*, 2(3). Available at: <https://doi.org/10.1016/j.jcvp.2022.100099>.
- Casado, M.S., M. Quintana Díaz, Palacios, D., V. Hortigüela, Schulke, C.M., J. García, A. Canabal, Pedrero, M.J.P., Ramos, A.V. and M.A. Arrese (2012). Relationship between the alveolar-arterial oxygen gradient and PaO₂/FiO₂—Introducing peep into the model. *Medicina Intensiva (English Edition)*, 36(5), pp.329–334. doi:<https://doi.org/10.1016/j.medine.2012.07.001>.
- Casella, M. *et al.* (2023) *Features, Evaluation, and Treatment of Coronavirus (COVID-19)*, *StatPearls*. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/33572857>.
- Dhont, S. *et al.* (2020) ‘The pathophysiology of “happy” hypoxemia in COVID-19’, *Respiratory Research*, 21(1). doi:10.1186/s12931-020-01462-5.
- Frat, J.-P. *et al.* (2015) ‘High-Flow Oxygen through Nasal Cannula in Acute Hypoxemic Respiratory Failure’, *New England Journal of Medicine*, 372(23), pp. 2185–2196. Available at: <https://doi.org/10.1056/NEJMoa1503326>.
- Gupta, B., Jain, G., Agarwal, A., Chandrakar, S. and Gupta, N. (2021). Arterial Blood Gas as a Predictor of Mortality in COVID Pneumonia Patients Initiated on Noninvasive Mechanical Ventilation: A Retrospective Analysis. *Indian Journal of Critical Care Medicine*, [online] 25(8), pp.866–871. doi: <https://doi.org/10.5005/jp-journals-10071-23917>.
- Hanci, P. *et al.* (2023) ‘ROX Index Dynamics According to High-flow Nasal Cannula Success in Intensive Care Unit Patients with COVID-19-Related Acute Respiratory Failure.’, *Balkan medical journal*, 40(2), pp. 111–116. Available at: <https://doi.org/10.4274/balkanmedj.galenos.2022.2022-6-31>.
- Huppert, L.A., Matthay, M.A. and Ware, L.B. (2019) ‘Pathogenesis of Acute Respiratory Distress Syndrome.’, *Seminars in respiratory and critical care medicine*, 40(1). Available at: <https://doi.org/10.1055/s-0039-1683996>.
- Li, C. *et al.* (2021) ‘Overview of the pathogenesis of COVID-19 (Review).’, *Experimental and therapeutic medicine*, 22(3). Available at: <https://doi.org/10.3892/etm.2021.10444>.

- Li, C. *et al.* (2021) ‘Overview of the pathogenesis of COVID-19 (Review).’, *Experimental and therapeutic medicine*, 22(3). Available at: <https://doi.org/10.3892/etm.2021.10444>.
- Nishiyama, K. *et al.* (2023) ‘The ROX index (Index combining the respiratory rate with oxygenation) is a prognostic factor for acute respiratory distress syndrome.’, *PloS one*, 18(2). Available at: <https://doi.org/10.1371/journal.pone.0282241>.
- Nishiyama, K. *et al.* (2023) ‘The ROX index (Index combining the respiratory rate with oxygenation) is a prognostic factor for acute respiratory distress syndrome.’, *PloS one*, 18(2). Available at: <https://doi.org/10.1371/journal.pone.0282241>.
- Nugraha, P.K. *et al.* (2022) ‘ROX Index as a Predictor of Intubation Risk and 28-Day Mortality in COVID-19 Patients with Acute Respiratory Distress Syndrome’, *International Journal Of Scientific Advances*, 3(1). Available at: <https://doi.org/10.51542/ijscia.v3i1.4>.
- Odeyemi, Y., Moraes, A.G. De and Gajic, O. (2020) ‘What factors predispose patients to acute respiratory distress syndrome?’, in *Evidence-Based Practice of Critical Care*. Elsevier. Available at: <https://doi.org/10.1016/B978-0-323-64068-8.00024-9>.
- Pipitone, G. *et al.* (2022) ‘Alveolar–arterial gradient is an early marker to predict severe pneumonia in COVID-19 patients’, *Infectious Disease Reports*, 14(3), pp. 470–478. doi:10.3390/idr140300 mmHg50.
- Polydora, E. *et al.* (2022) ‘Predictors of high-flow oxygen therapy failure in COVID-19-related severe hypoxemic respiratory failure.’, *Journal of thoracic disease*, 14(4). Available at: <https://doi.org/10.21037/jtd-21-1373>.
- Poopipatpab, S. *et al.* (2023) ‘The use of respiratory rate-oxygenation index to predict failure of high-flow nasal cannula in patients with coronavirus disease 2019-associated acute respiratory distress syndrome: A retrospective study.’, *PloS one*, 18(6). Available at: <https://doi.org/10.1371/journal.pone.0287432>.
- Poopipatpab, S. *et al.* (2023) ‘The use of Respiratory Rate-oxygenation index to predict failure of high-flow nasal cannula in patients with coronavirus disease 2019-associated acute respiratory distress syndrome: A retrospective study’, *PLOS ONE*, 18(6). doi:10.1371/journal.pone.0287432.
- Sweeney, R. Mac and McAuley, D.F. (2016) ‘Acute respiratory distress syndrome’, *The Lancet*, 388(10058), pp. 2416–2430. Available at: [https://doi.org/10.1016/S0140-6736\(16\)00578-X](https://doi.org/10.1016/S0140-6736(16)00578-X).

- Wirakusuma, I.B. *et al.* (2022) 'Correlation between alveolar-arterial oxygen tension difference gradient baseline with spirometry test in severe coronavirus disease 2019 (COVID-19)', *International journal of health sciences*, pp. 2578–2588. doi:10.53730/ijhs.v6ns9.12986.
- Yadam, S., Bihler, E. and Balaan, M. (2016) 'Acute Respiratory Distress Syndrome.', *Critical care nursing quarterly*, 39(2). Available at: <https://doi.org/10.1097/CNQ.000000000000111>.
- Yau, C.E. *et al.* (2023) 'Performance of the ROX index in predicting high-flow nasal cannula failure in COVID-19 patients: A systematic review and meta-analysis', *Critical Care*, 27(1). doi:10.1186/s13054-023-04567-7.