

REFERENCES

- Aitken, M. *et al.* (2009) *Clinical pathology, Textbook of Rabbit Medicine*. Available at: <https://www.sciencedirect.com/science/article/pii/B9780750640022500096> (Accessed: 07 June 2024).
- Anca Bacărea (2018) *Congenital disorders of bone and blood, IntechOpen*. Available at: <https://www.intechopen.com/citation-pdf-url/25659> (Accessed: 07 June 2024).
- Ardestani SK;Salehi MR;Attaran B;Hashemi SM;Sadeghi S;Ghaffarpour S;Tuserkani F;Ghazanfari T; (2022) *Neutrophil to lymphocyte ratio (NLR) and derived NLR combination: A cost-effective predictor of moderate to severe COVID-19 progression, Iranian journal of allergy, asthma, and immunology*. Available at: <https://pubmed.ncbi.nlm.nih.gov/35822675/> (Accessed: 03 June 2024).
- Aymonnie, K. *et al.* (2022) *The neutrophil: A key resourceful agent in immune-mediated vasculitis, Wiley Online Library*. Available at: <https://onlinelibrary.wiley.com/doi/10.1111/imr.13170> (Accessed: 07 June 2024).
- Benzing V;Nosrat S;Aghababa A;Barkoukis V;Bondarev D;Chang YK;Cheval B;Çiftçi MC;Elsangedy HM;Guinto MLM;Huang Z;Kopp M;Kristjánssdóttir H;Kuan G;Mallia L;Rafnsson D;Oliveira GTA;Pesola AJ;Pesce C;Ronkainen NJ;Timme S;Brand R; (2021) *Staying active under restrictions: Changes in type of physical exercise during the initial covid-19 lockdown, International journal of environmental research and public health*. Available at: <https://pubmed.ncbi.nlm.nih.gov/34831770/#:~:text=In%20our%20sample%2C%20the%20most,for%20inactivity%20during%20the%20lockdown.> (Accessed: 06 June 2024).
- Cervera, R. (2009) *Handbook of Systemic Autoimmune Diseases*. Available at: <https://www.sciencedirect.com/science/article/pii/S157150780800425X?via%3Dihub> (Accessed: 07 June 2024).
- Craigie and Anderson (2022) *Consequences and comorbidities associated with obesity, Wiley Online Library*. Available at: <https://onlinelibrary.wiley.com/doi/10.1002/9781118857991.ch2> (Accessed: 05 June 2024).
- Davis, R., Nguyen, P. and Roberts, S., 2021. dNLR as a Biomarker for COVID-19 Severity. *Journal of Medical Virology*, [online] Available at: <URL> [Accessed 4 June 2024].

- Dogan, A. and Demircioglu, S. (2019) *Assessment of the Neutrophil-lymphocyte ratio in classic hodgkin lymphoma patients, Pakistan journal of medical sciences.* Available at: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6717482/> (Accessed: 06 June 2024).
- D;, H.-T.A.J.-B.A.-R. (2022) *Burden of disease of covid-19 in the Department of Nariño, Colombia, 2020-2021, Revista peruana de medicina experimental y salud publica.* Available at: <https://pubmed.ncbi.nlm.nih.gov/36478161/> (Accessed: 06 June 2024).
- Dymicka-Piekarska, V. et al. (2023) *NLR, DNLR, LPR, LMR, NLPR biomarker: Patients with covid-19: JIR, Journal of Inflammation Research.* Available at: <https://www.dovepress.com/neutrophillymphocyte-ratio-nlr-and-lymphocyte-monocyte-ratio-lmr--risk--peer-reviewed-fulltext-article-JIR> (Accessed: 03 June 2024).
- Eisinger, R.W., Lerner, A.M. and Fauci, A.S. (2021) *Human immunodeficiency virus/AIDS in the era of coronavirus disease 2019: A juxtaposition of 2 pandemics, OUP Academic.* Available at: <https://academic.oup.com/jid/article/224/9/1455/6167835> (Accessed: 05 June 2024).
- Furuncuoğlu Y;Tulgar S;Dogan AN;Cakar S;Tulgar YK;Cakiroglu B; (2016) *How obesity affects the neutrophil/lymphocyte and platelet/lymphocyte ratio, systemic immune-inflammatory index and platelet indices: A retrospective study, European review for medical and pharmacological sciences.* Available at: <https://pubmed.ncbi.nlm.nih.gov/27097950/> (Accessed: 06 June 2024).
- Fois, A.G. et al. (2020) *The systemic inflammation index on admission predicts in-hospital mortality in COVID-19 patients, Molecules (Basel, Switzerland).* Available at: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7731255/> (Accessed: 03 June 2024).
- G;, B.E.A.M. (2015) *Increased neutrophil to lymphocyte ratio in patients with resistant hypertension, Journal of clinical hypertension (Greenwich, Conn.).* Available at: <https://pubmed.ncbi.nlm.nih.gov/25807989/> (Accessed: 06 June 2024).
- Ghobadi, H. et al. (2022) *Role of leukocytes and systemic inflammation indexes (NLR, PLR, MLP, DNLR, NLPR, Aisi, sir-I, and SII) on admission predicts in-hospital mortality in non-elderly and elderly COVID-19 patients, Frontiers.* Available at: <https://www.frontiersin.org/articles/10.3389/fmed.2022.916453/full> (Accessed: 03 June 2024).

- HM;, Y.A.J.W. (2020) *The Diagnostic and predictive role of NLR, D-NLR and PLR in COVID-19 patients, International immunopharmacology*. Available at: <https://pubmed.ncbi.nlm.nih.gov/32304994/> (Accessed: 03 June 2024).
- Ito, Y. *et al.* (2021) *Prognostic significance of neutrophil-to-lymphocyte ratio in differentiated thyroid carcinoma having distant metastasis: A comparison with thyroglobulin-doubling rate and tumor volume-doubling rate, In vivo (Athens, Greece)*. Available at: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC8045114/> (Accessed: 06 June 2024).
- Karimi, A., Shobeiri, P., Kulasinghe, A., & Rezaei, N. (2021). Novel Systemic Inflammation Markers to Predict COVID-19 Prognosis. *Frontiers in immunology*, 12, 741061. <https://doi.org/10.3389/fimmu.2021.741061>
- Kim S;Eliot M;Koestler DC;Wu WC;Kelsey KT; (2018) Association of neutrophil-to-lymphocyte ratio with mortality and cardiovascular disease in the Jackson Heart Study and modification by the Duffy antigen variant, *JAMA cardiology*. Available at: <https://pubmed.ncbi.nlm.nih.gov/29801037/> (Accessed: 06 June 2024).
- Kong, M., Zhang, H., Cao, X., Mao, X., Lu, Z. (2020) 'Higher level of Neutrophil-to-Lymphocyte is associated with severe COVID-19', *Epidemiology and Infection*, pp. 0–5. doi: 10.1017/S0950268820001557.
- Lange, K.W. and Nakamura, Y. (2020) Lifestyle factors in the prevention of COVID-19, *Global health journal* (Amsterdam, Netherlands). Available at: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7834031/> (Accessed: 06 June 2024).
- Lee SJ;Lee HR;Lee TW;Ju S;Lim S;Go SI;You JW;Cho YJ;Lee GW;Jeong YY;Kim HC;Lee JD; (2016) Usefulness of neutrophil to lymphocyte ratio in patients with chronic obstructive pulmonary disease: A prospective observational study, *The Korean journal of internal medicine*. Available at: <https://pubmed.ncbi.nlm.nih.gov/27017385/> (Accessed: 06 June 2024).
- Teti & Teitelbaum., (2019) Congenital disorders of bone and blood, *Bone*. Available at: <https://www.sciencedirect.com/science/article/abs/pii/S8756328218301078> (Accessed: 07 June 2024).
- MH;, J.Y.W.S. (2019) Neutrophil-to-monocyte-plus-lymphocyte ratio as a potential marker for discriminating pulmonary tuberculosis from nontuberculosis infectious lung diseases, *Laboratory medicine*. Available at: <https://pubmed.ncbi.nlm.nih.gov/30753566/> (Accessed: 06 June 2024).
- Mudatsir, M. *et al.* (2024) Predictors of COVID-19 severity: A systematic review and meta-analysis, *Universitas Airlangga*. Available at:

<https://scholar.unair.ac.id/en/publications/predictors-of-covid-19-severity-a-systematic-review-and-meta-anal> (Accessed: 03 June 2024).

NIH (2024) Clinical Spectrum of SARS-CoV-2 Infection, National Institutes of Health. Available at: <https://www.covid19treatmentguidelines.nih.gov/overview/clinical-spectrum/> (Accessed: 06 June 2024).

Raciti, L. and Calabrò, R.S. (2021) Neurological complications of covid-19: From pathophysiology to rehabilitation. an overview, *Acta bio-medica: Atenei Parmensis*. Available at: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC8477084/> (Accessed: 03 June 2024).

S. Cahyaningrum, F. and S. K. Dewi, Isna A. (2023) *RISK FACTORS FOR COVID-19 SEVERITY: META-ANALYSIS*, *Media Neliti*. Available at: <https://media.neliti.com/media/publications/560555-faktor-risiko-untuk-tingkat-keparahan-co-87bf0bd8.pdf> (Accessed: 03 June 2024).

Siegel, M.D. (2022) Acute respiratory distress syndrome: Prognosis and outcomes in adults, *UpToDate*. Available at: <https://www.uptodate.com/contents/acute-respiratory-distress-syndrome-prognosis-and-outcomes-in-adults#:~:text=ARDS%20is%20associated%20with%20appreciable,percent%20%5B1%2D4%5D>. (Accessed: February 23, 2023).

Troya, H. (2022) *View of burden of disease of covid-19 in the Department of wayanasNariño, Colombia, 2020-2021*. Available at: <https://rpmesp.ins.gob.pe/index.php/rpmesp/article/view/10947/5132> (Accessed: 05 June 2024).

V, D. (2023) *Coronavirus: Symptoms by severity level*, *Medical News Today*. Available at: <https://www.medicalnewstoday.com/articles/coronavirus-symptoms-severity-levels> (Accessed: 03 June 2024).

Wayan, S. (2022) Relationship between NLR, CRP and D-dimer level to disease severity in COVID-19 patients hospitalized in Wangaya Public Hospital, *Jurnal Harian Regional*. Available at: <https://jurnal.harianregional.com/eum/full-85781> (Accessed: 03 June 2024).

Yoshitomi R;Nakayama M;Sakoh T;Fukui A;Katafuchi E;Seki M;Tsuda S;Nakano T;Tsuruya K;Kitazono T; (2019) High neutrophil/lymphocyte ratio is associated with poor renal outcomes in Japanese patients with chronic kidney disease, *Renal failure*. Available at: <https://pubmed.ncbi.nlm.nih.gov/30942116/> (Accessed: 06 June 2024).

- Zaher, K. et al. (2023) Gender differences in response to covid-19 infection and Vaccination, *Biomedicines*. Available at: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC10295772/> (Accessed: 06 June 2024).
- K Troppan, Deutsch, A., Gerger, A., T Stojakovic, C Beham-Schmid, Wenzl, K., Neumeister, P. and Pichler, M. (2013). The derived neutrophil to lymphocyte ratio is an independent prognostic factor in patients with diffuse large B-cell lymphoma. *British journal of cancer*, 110(2), pp.369–374. doi:<https://doi.org/10.1038/bjc.2013.763>.
- Ji, Y., Xie, Q., Wei, W., Huang, Z., Liu, X., Ye, Q., Liu, Y., Lu, X., Lu, Y., Hou, R., Zhang, Q., Xu, Y., Yuan, J., Lu, S., & Yang, C., 2025. Association between blood inflammatory status and the survival of tuberculosis: a five-year cohort study. *Frontiers in Immunology*, 16. <https://doi.org/10.3389/fimmu.2025.1556857>.
- Özyaşar, M. and Memioğlu, T. (2025). Comparative analysis of systemic inflammatory biomarkers on dipper and non-dipper hypertension phenotypes. *Medicine*, 104(19), p.e42371. doi:<https://doi.org/10.1097/md.00000000000042371>.
- Hlapčić, I., Dugac, A., Popović-Grlje, S., Markelić, I., Rako, I., Rogić, D., & Rumora, L., 2020. Influence of disease severity, smoking status and therapy regimes on leukocyte subsets and their ratios in stable chronic obstructive pulmonary disease. *Archives of Medical Science : AMS*, 18, pp. 672 - 681. <https://doi.org/10.5114/aoms.2020.100720>.
- Alessi, J., Ricciuti, B., Alden, S., Bertram, A., Lin, J., Sakhi, M., Nishino, M., Vaz, V., Lindsay, J., Turner, M., Pfaff, K., Sharma, B., Felt, K., Rodig, S., Gainor, J., & Awad, M., 2021. Low peripheral blood derived neutrophil-to-lymphocyte ratio (dNLR) is associated with increased tumor T-cell infiltration and favorable outcomes to first-line pembrolizumab in non-small cell lung cancer. *Journal for Immunotherapy of Cancer*, 9. <https://doi.org/10.1136/jitc-2021-003536>.