

DAFTAR PUSTAKA

- Adiga, D. J., Veena, G. L., Thondaiman, V., & Babli, M. (2020). An overview of canopy management in cashew (*Anacardium occidentale* L.). *Journal of Horticultural Sciences*, 15(2), 127–135.
- Adu-Gyamfi, P. K. K., Barnor, M., Akpertey, A., Dadzie, A. M., Anyomi, E., Osei-Akoto, S., & Padi, F. (2021). Broadening the Gene Pool of Cashew (*Anacardium occidentale*) for Survival and Precocity. *Agricultural Research*, 10(4), 613–625. Scopus. <https://doi.org/10.1007/s40003-020-00521-z>
- Ahmad, R., Malik, W., & Anjum, M. A. (2019). Genetic Diversity and Selection of Suitable Molecular Markers for Characterization of *Indigenous Zizyphus* Germplasm. *Erwerbs-Obstbau*, 61(4), 345–353. <https://doi.org/10.1007/s10341-019-00438-0>
- Aidoo, R., Kwofie, E. M., & Ngadi, M. O. (2022). Circularity of Cashew Apples: Examining the Product-Process Pathways, Techno-Functional, Nutritional/Phytomolecular Qualities for Food Applications. *ACS Food Science and Technology*, 2(7), 1051–1066. Scopus. <https://doi.org/10.1021/acsfoodscitech.2c00093>
- Akyereko, Y. G., Yeboah, G. B., Wireko-Manu, F. D., Alemawor, F., Mills-Robertson, F. C., & Odoom, W. (2023). Nutritional value and health benefits of cashew apple. *JSFA Reports*, 3(3), 110–118. Scopus. <https://doi.org/10.1002/jsf2.107>
- Al-Gharaibeh, M. M., Hamasha, H. R., Rosche, C., Lachmuth, S., Wesche, K., & Hensen, I. (2017). Environmental gradients shape the genetic structure of two *Medicinal salvia* species in Jordan. *Plant Biology*, 19(2), 227–238. Scopus. <https://doi.org/10.1111/plb.12512>
- Aliyu, O. & Mashood. (2006). Application of tissue culture to cashew (*Anacardium occidentale* L.) breeding: An appraisal. *African Journal of Biotechnology*, 4, 1485–1489.
- Aliyu, O. M., & Awopetu, J. A. (2007). Multivariate analysis of cashew (*Anacardium occidentale* L.) germplasm in Nigeria. *Silvae Genetica*, 56(3–4), 170–179. <https://doi.org/10.1515/sg-2007-0026>
- Aliyu, O. M., & Awopetu, J. A. (2011). Variability study on nut size and number trade-off identify a threshold level for optimum yield in cashew (*anacardium occidentale* L.). *International Journal of Fruit Science*, 11(4), 342–363. Scopus. <https://doi.org/10.1080/15538362.2011.630297>
- Amirthavinayagam, N., & Meenakshi, K. (2025). Harnessing the Deep Learning techniques in Cashew plant disease management. *Int. Conf. Comput. Robot., Test. Eng. Eval., ICCRTEE*. International Conference on Computational Robotics, Testing and

- Archak, S., Gaikwad, A. B., Gautam, D., Rao, E. V. V. B., Swamy, K. R. M., & Karihaloo, J. L. (2003). Comparative assessment of DNA fingerprinting techniques (RAPD, ISSR and AFLP) for genetic analysis of cashew (*Anacardium occidentale* L.) accessions of India. *Genome*, 46(3), 362–369. <https://doi.org/10.1139/g03-016>
- Archak, S., Gaikwad, A. B., Gautam, D., Rao, E. V. V. B., Swamy, K. R. M., & Karihaloo, J. L. (2003). DNA fingerprinting of Indian cashew (*Anacardium occidentale* L.) varieties using RAPD and ISSR techniques. *Euphytica*, 230, 397–404.
- Archak, S., Gaikwad, A. B., Swamy, K. R. M., & Karihaloo, J. L. (2009). Genetic analysis and historical perspective of cashew (*Anacardium occidentale* L.) introduction into India. *Genome*, 52(3), 222–230. <https://doi.org/10.1139/G08-119>
- Asna, A. C., Menon, J. S., & Smitha, M. S. (2022). Phenotypic diversity and clustering of germplasm accessions of cashew for utilization and conservation. *Electronic Journal of Plant Breeding*, 12(4), 1218–1226. <https://doi.org/10.37992/2021.1204.167>
- Barros, L.M., Paiva, J.R., Crisóstomo, J.R. and Cavalcanti, J.J.V. (1999). Hybridization of the cashew tree. p.191-220. In: A. Borém (ed.) Artificial hybridization of plants. UFV, Viços
- Bartsch, D. (2010). Gene Flow in Sugar Beet. *Sugar Tech*, 12(3–4), 201–206. Scopus. <https://doi.org/10.1007/s12355-010-0053-1>
- Beguedou, E., Narra, S., Agboka, K., Armoo, E.A., Damgou, M.K. (2024). Cashew Nutshell and Raw Coal Co-grinding for Sustainability Goals in Clinkerization Pyro-Processes. In: Narra, MM., Narra, S. (eds) Innovations in Circular Economy and Renewable Energy in Africa. World Sustainability Series. Springer, Cham. https://doi.org/10.1007/978-3-031-68330-5_14
- Benjamin R. K, Alexander L.S., Umilaela A., Djoko T. I., Evy A., Christopher C. A., Jimmy A.M. (2020) "Crossing Lydekker's Line: Northern Water Dragons (*Tropicagama temporalis*) Colonized the Mollucan Islands of Indonesia from New Guinea. *Herpetologica*, 76(3), 344-350.
- Bocianowski, J., Niemann, J., Jagieniak, A., & Szwarc, J. (2024). Comparison of Six Measures of Genetic Similarity of Interspecific Brassicaceae Hybrids F2 Generation and Their Parental Forms Estimated on the Basis of ISSR Markers. *Genes*, 15(9). Scopus. <https://doi.org/10.3390/genes15091114>
- Boer, D., Ladati, N., Jumarno, Irwan, Arsyad, M. A., & Larekeng, S. H. (2021). Morphological diversity and relationship among cashew (*Anacardium occidentale* L.) individuals in three districts of Southeast Sulawesi, Indonesia. *IOP Conference Series: Earth and Environmental Science*, 886(1). <https://doi.org/10.1088/1755-1315/886/1/012056>

- Borges, A. N. C., Lopes, A. C. A., Britto, F. B., Vasconcelos, L. F. L., & Lima, P. S. C. (2018). Genetic diversity in a cajui (*Anacardium* spp.) germplasm bank as determined by *issr* markers. *Genetics and Molecular Research*, 17(4). <https://doi.org/10.4238/gmr18212>
- Bornet, B., & Branchard, M. (2001). Nonanchored Inter Simple Sequence Repeat (ISSR) Markers: Reproducible and Specific Tools for Genome Fingerprinting. *Plant Molecular Biology Reporter*, 19(3), 209–215. <https://doi.org/10.1007/BF02772892>
- Bradford, K. 2008. Comparing the ability of two PCR based techniques, RAPD and ISSR to detect low level of genetic diversity. In Chicago botanic Garden. <http://www.chicagobotanic.org/downloads/conservation>. Diakses 7 September 2022.
- Braasch, J. E., Di Santo, L. N., Tarble, Z. J., Prasifka, J. R., & Hamilton, J. A. (2021). Testing for evolutionary change in restoration: A genomic comparison between *ex situ*, native, and commercial seed sources of *Helianthus maximiliani*. *Evolutionary Applications*, 14(9), 2206–2220. <https://doi.org/10.1111/eva.13275>
- Buljević, N., Preiner, D., Šikuten, I., & Tomaz, I. (2025). Comparison of Different DNA Isolation Methods from Grapevine (*Vitis vinifera*) Leaves. *Separations*, 12(11). Scopus. <https://doi.org/10.3390/separations12110316>
- Campos, A. S., Jesus, E. D. C., Mendes Filho, P. F., Pereira, A. P. D. A., Castro, A. C. R. D., Coelho, A. C. M. A., Bordallo, P. D. N., Borges, W. L., & Carvalho, A. C. P. P. D. (2025). Soil quality changes in cashew orchard production in Brazilian semiarid areas: A comprehensive field survey across locations, seasons, tree types, and management practices. *Applied Soil Ecology*, 211. Scopus. <https://doi.org/10.1016/j.apsoil.2025.106102>
- Carneiro, L. A., Silva, L. S., Gomes, M. F. C., Santos, M. F., Valente, S. E. S., Gomes, R. L. F., & Costa, M. F. (2019). Morphological characterization and genetic divergence of a cashew population in Floriano, Piauí, Brazil. *Genetics and Molecular Research*, 18(3), 1–8. <https://doi.org/10.4238/gmr18348>
- Carvalho, R. S., Pinto, J. F. N., dos Reis, E. F., Santos, S. C., & Dias, L. A. S. (2012). Genetic variability of bushy cashew (*Anacardium humile* ST. HILL.) through RAPD markers. *Revista Brasileira de Fruticultura*, 34(1), 227–233. Scopus. <https://doi.org/10.1590/S0100-29452012000100030>
- Casazza, G., Macrì, C., Dagnino, D., Guerrina, M., Juin, M., Minuto, L., Thompson, J. D., Baumel, A., & Médail, F. (2021). When ecological marginality is not geographically peripheral: Exploring genetic predictions of the centre-periphery hypothesis in the endemic plant *Lilium pomponium*. *PeerJ*, 9. Scopus. <https://doi.org/10.7717/peerj.11039>
- Castro, A. C. R., Bordallo, P. N., Cavacanti, J. J. V., & Barros, L. M. (2011). Brazilian cashew germplasm bank. *Acta Horticulturae*, 918, 857–861. <https://doi.org/10.17660/ActaHortic.2011.918.113>
- Cavalcanti, J. J. V., & Wilkinson, M. J. (2007). The first genetic maps of cashew (*Anacardium occidentale* L.). *Euphytica*, 157(1–2), 131–143. <https://doi.org/10.1007/s10681-007-9403-9>

- Chipojola, F. M., Mwase, W. F., Kwapata, M. B., Bokosi, J. M., Njoloma, J. P., & Maliro, M. F. (2009). Morphological characterization of cashew (*Anacardium occidentale* L.) in four populations in Malawi. *African Journal of Biotechnology*, 8(20), 5173–5181.
- Czuppon B., Ifj. Papp L., Tóth Z., & Szépligeti M. (2019). A kereklevelű harmatfű (*Drosera rotundifolia*L.) szaporodásbiológiai vizsgálata, különös tekintettel az ex situ természetvédelmi módszerekre. *Botanikai Közlemények*, 106(1), 41–63. <https://doi.org/10.17716/BotKozlem.2019.106.1.41>
- Daras, U. (2007). Strategi dan Iovasi Teknologi Peingkatan Produktivitas Jambu Mete Di Nusa Tenggara. *Jurnal Litbang Pertanian*, 26(1), 25–34.
- Das, S., Ali, M. N., Bandyopadhyay, S., & Poduval, M. (2024). Multivariate analysis from maturase K (matK) gene-based markers and morphological indices in Indian cashew. *Plant Genetic Resources: Characterisation and Utilisation*, 22(1), 17–26. <https://doi.org/10.1017/S1479262123000874>
- Dasmohapatra, R., Rath, S., Pradhan, B., & Rout, G. R. (2014). Molecular and agromorphological assessment of cashew (*Anacardium occidentale* L.) genotypes of India. *Journal of Applied Horticulture*, 16(3), 215–221.
- de Brito, E. S., de Oliveira Silva, E., & Rodrigues, S. (2018). Caju—*Anacardium occidentale*. In *Exotic Fruits Reference Guide* (pp. 85–89). Elsevier; Scopus. <https://doi.org/10.1016/B978-0-12-803138-4.00012-5>
- Dhanaraj, A. L., Bhaskara Rao, E. V. V., Swamy, K. R. M., Bhat, M. G., Theertha Prasad, D., & Sondur, S. N. (2002). Using RAPDs to assess the diversity in Indian cashew (*Anacardium occidentale* L.) germplasm. *Journal of Horticultural Science and Biotechnology*, 77(1), 41–47. <https://doi.org/10.1080/14620316.2002.11511454>
- Dimoso, N., Aluko, A., Makule, E., & Kassim, N. (2021). Challenges and opportunities toward sustainable consumption and value addition of cashew apples in Tanzania. *Outlook on Agriculture*, 50(2), 169–177. Scopus. <https://doi.org/10.1177/0030727020941164>
- Ditjenbun. (2019). *Statistik Perkebunan Indonesia*. www.ditjenbun.pertanian.go.id. Diakses 7 September 2022.
- Dlugosch, K. M., & Parker, I. M. (2008). Invading populations of an ornamental shrub show rapid life history evolution despite genetic bottlenecks. *Ecology Letters*, 11(7), 701–709. Scopus. <https://doi.org/10.1111/j.1461-0248.2008.01181.x>
- dos Santos, J. O., Mayo, S. J., Bittencourt, C. B., & de Andrade, I. M. (2019). Genetic diversity in wild populations of the restinga ecotype of the cashew (*Anacardium occidentale*) in coastal Piauí, Brazil. *Plant Systematics and Evolution*, 305(10), 913–924. <https://doi.org/10.1007/s00606-019-01611-4>
- Douaihy B, Sobierajska K, Jasińska AK, Boratyńska K, Ok T, Romo A, Machon N, Didukh Y, Bou Dagher-Kharrat M, Boratyński A. (2012). Morphological versus molecular

markers to describe variability in *Juniperus excelsa* subsp. *excelsa* (Cupressaceae). *AoB Plants*. 2012:pls013. doi: 10.1093/aobpla/pls013.

- Ehoniyan, O. I., & Udo, E.-O. J. (2019). Variation studies of morphological characters of cashew trees (*Anacardium occidentale*) on Kogi State University Campus, Anyigba, Kogi State, Nigeria. *GSC Biological and Pharmaceutical Sciences*, 8(3), 017–022. <https://doi.org/10.30574/gscbps.2019.8.3.0161>
- Engler, A. (1879). *Monographiae: Phanerogamarum Burseraceae et Anacardiaceae* (Alphonso & D. C. Casimir, Eds.; Vol. 4). Sumptibus G. Masson.
- FAO. (2022). Cashew Production by Country 2022. <https://worldpopulationreview.com>. Diakses tanggal 27 November 2022.
- Fang, D.Q. & M.Z. Roose, 1997. Identification of closely related citrus cultivars with inter simple sequence repeat markers. *TheorAppl Genet* 95: 408–417
- Febriani, S., & Hartana, A. (2016). Klasifikasi Kultivar Jambu Mete (*Anacardium occidentale* L.). *Floribunda*, 5(4), 139–143.
- Ferry, Y. (2012) Pengembangan Industri Perbenihan Jambu Mete. *Perspektif*. 11 (1), 33–44
- Frercks, B., Stanys, V., Siksniene, J. B., Stepulaitiene, I., Gelvonauskiene, D., Staniene, G., Rugienius, R., & Siksniene, T. (2014). Efficiency of AFLP marker attributes in the genetic analysis of sweet cherry cultivars. *Journal of Food, Agriculture and Environment*, 12(1), 122–127.
- Gamermann, D., Montagud, A., Alberto Conejero, J., de Córdoba, P. F., & Urchueguía, J. F. (2019). Large scale evaluation of differences between network-based and pairwise sequence-alignment-based methods of dendrogram reconstruction. *PLoS ONE*, 14(9). Scopus. <https://doi.org/10.1371/journal.pone.0221631>
- Garcia Gonçalves, A. P., Bassi Scarpim, L., Alves Garcia, C., Candido da Silva, T., Goloni, C., de Souza Ávida Castro, T., Carciofi, A. C. (2025). Cashew (*Anacardium occidentale*) nut co-product in extruded foods for dogs. *Archives of Animal Nutrition*, 79(2), 95–110. <https://doi.org/10.1080/1745039X.2025.2503172>
- Gateva, S. P., Jovtchev, G., Angelova, T. V., Nonova, T. P., Tyutyundzhiev, N., Geleva, E. G., Katrandzhiev, K., Nikolova, N. A., Dimitrov, D., & Angelov, C. V. (2022). Effect of uv radiation and other abiotic stress factors on dna of different wild plant species grown in three successive seasons in alpine and subalpine regions. *Phyton-International Journal of Experimental Botany*, 91(2), 293–313. Scopus. <https://doi.org/10.32604/phyton.2022.016397>
- Germano, T. A., de Oliveira, M. F. R., Aziz, S., Oliveira, A. E. R., da Cruz Saraiva, K. D., dos Santos, C. P., Moura, C. F. H., & Costa, J. H. (2022). Transcriptome profiling of cashew apples (*Anacardium occidentale*) genotypes reveals specific genes linked to firmness and color during pseudofruit development. *Plant Molecular Biology*, 109(1–2), 83–100. <https://doi.org/10.1007/s11103-022-01257-w>

Gomes, M. F. C., Borges, A. N. C., Batista, G., Luz, G., Oliveira, M., Lopes, Â., de ARAÚJO, A., Gomes, R. L. F., Britto, F., Lima, P., & Valente, S. E. S. (2021). Genetic Diversity and Structure In Natural Populations OF Cajui From Brazillian Cerrado. *Bioscience Journal*, 37. <https://doi.org/10.14393/BJ-V37N0A2021-53974>

Guterres, L., Barnabé, J., Barros, A., Charrua, A. B., Duarte, M. C., Romeiras, M. M., & Monteiro, F. (2023). East Timor as an important source of cashew (*Anacardium occidentale* L.) genetic diversity. *PeerJ*, 11. <https://doi.org/10.7717/peerj.14894>

Gupta, M., Y.S. Chyi, J. Romero-Severson & J.L. Owen, 1994. Amplification of DNA markers from evolutionarily diverse gen-omes using single primers of simple-sequence repeats. *TheorAppl Genet* 89: 998–1006

Hadad, E. A., Syakir, M., & Rosman, R. (2006). *Karakterisasi Plasma Nutfah Jambu Mete*. Bogor: Balai Penelitian Tanaman Rempah dan Obat (Balittro).

Hamrick, J. L., & Godt, M. J. W. (1996). Effects of life history traits on genetic diversity in plant species. *Philosophical Transactions of the Royal Society of London. Series B: Biological Sciences*, 351(1345), 1291–1298. <https://doi.org/10.1098/rstb.1996.0112>

Hansen, M., Halldén, C., and Säll, T. 1998. Error rates and polymorphism frequencies for three RAPD protocols. *Plant Mol Biol Rep* 16: 139-146.

Haryudin, W., & Rostiana, O. (2017). Keragaman 16 Aksesi Jambu Mete Hasil Grafting Berdasarkan Karakter Morfologis. *Buletin Penelitian Tanaman Rempah Dan Obat*, 27(2), 93. <https://doi.org/10.21082/bullittro.v27n2.2016.93-104>

Hatai, L. D. (2018). Economic feasibility of on-farm processing and value chain of cashewnut in west garo hills of meghalaya. *Journal of Plantation Crops*, 46(3), 196–203. Scopus. <https://doi.org/10.25081/jpc.2018.v46.i3.2078>

Hawerroth, M. C., do Nascimento Bordallo, P., Oliveira, L. C. P., Vale, E. H., das Chagas Vidal Neto, F., & Melo, D. S. (2019). Genetic variability among cashew hybrids and prediction of superior combinations based on agronomic performance. *Pesquisa Agropecuaria Brasileira*, 54. Scopus. <https://doi.org/10.1590/S1678-3921.PAB2019.V54.00725>

IBPGR. (1986). *International Board for Plant Genetic Resources: Cashew Descriptors* (pp. 1–36). International Board for Plant Genetic Resources (IPBGR).

Jena, R. C., Samal, K. C., Pal, A., Das, B. K., & Chand, P. K. (2016). Genetic Diversity Among Some Promising Indian Local Selections and Hybrids of Cashew Nut Based on Morphometric and Molecular Markers. *International Journal of Fruit Science*, 16(1), 69–93. <https://doi.org/10.1080/15538362.2015.1046321>

Jeyavishnu, K., Thulasidharan, D., Shereen, M. F., & Arumugam, A. (2021). Increased Revenue with High Value-Added Products from Cashew Apple (*Anacardium occidentale* L.) Addressing Global Challenges. *Food and Bioprocess Technology*, 14(6), 985–1012. <https://doi.org/10.1007/s11947-021-02623-0>

Karmawati, E., Penelitian, P., & Perkebunan, P. (2008). *Perkembangan Jambu Mete dan Strategi Pengendalian Hama Utamanya*. 7(2), 102–111.

Karp, A. . (1997). *Molecular tools in plant genetic resources conservation: A guide to the technologies*. IPGRI.

Kaunang, I. R. B., Haliadi., & Rabani, O. 2016. *Jaringan Maritim Indonesia : Sejarah Toponim Kota Pantai Di Sulawesi* (J. Bangun, Ed.). Direktorat Sejarah Direktorat Jendral Kebudayaan Kementrian Pendidikan dan Kebudayaan.

Kebede, M., & Bekele, E. (2018). Genetic diversity and population structure of Ethiopian sesame (*Sesamum indicum* L.) germplasm as revealed by ISSR markers. *African Journal of Agricultural Research*, 13(45), 2575–2586. <https://doi.org/10.5897/AJAR2018.13490>

Khatun, M., Sarker, A., Akhter, S., & Chowdhury, M. A. Z. (2022). Morphological and molecular characterization of Lablab (*Lablab purpureus* L.) germplasm. *Asian Journal of Agricultural and Biological Research*, 8(1), 25–37.

Konan K, C., Adopo, da, Jean-Baptiste Djaha, A., Pokou, D. N., Annicet, H. N., Arsène Zoro Bi, I., Kévin Koffi, K., Djidji, H., Yves Minhibo, M., Dosso, M., & Évariste, A. N. (2020). Genetic characterization of promising high-yielding cashew (*Anacardium occidentale* L.) cultivars from Côte d'Ivoire. *Biotechnol. Agron. Soc. Environ.* 2020, 24(1), 46–58.

Konate, M., Tarpaga, W. V., Bourgou, L., & Wonni, I. (2022). Genetic diversity assessment among 18 elite cashew tree genotypes (*Anacardium occidentale* L) selected in Western Burkina Faso. *Journal of Plant Breeding and Crop Science*, 14(1), 1–11. <https://doi.org/10.5897/jpbcs2021.0986>

Kouakou, C. K., Konan, J. A., Konan, D., & Kouadio, A. B. R. (2021). Genetic Diversity of Cashew. In *Cash Crops: Genetic Diversity, Erosion, Conservation and Utilization* (pp. 523–555). Springer International Publishing; Scopus. https://doi.org/10.1007/978-3-030-74926-2_14

Kumar, A., Sharma, N., Panwar, S., & Kumar, M. (2014). Polymorphic Information Content (PIC) and marker utility of molecular markers. *Indian Journal of Genetics and Plant Breeding*, 74(1), 95–99.

Kumar, S.(2014). *The Role of Bioinformatics in Agriculture* (1st ed.). Apple Academic Press. <https://doi.org/10.1201/b16568>

Lohman, D. J., De Bruyn, M., Page, T., Von Rintelen, K., Hall, R., Ng, P. K. L., Shih, H. T., Carvalho, G. C., & Von Rintelen, T. (2011). Beyond Wallaces line: Genes and biology inform historical biogeographical insights in the Indo-Australian archipelago. *Annual Review of Ecology, Evolution, and Systematics*, 42. <https://doi.org/10.1146/annurev-eolsys-102710-145001>

Lovatel, Q. C., Rosa, G. T. D., Sá, A. C. S., Lopes, B. C., Tonett, E. L., Dias, R. A. R., Pereira, M. D. O., & Navroski, M. C. (2021). Impacts of genetic selection on Sequoia

- Lucena-Aguilar, G., Sánchez-López, A. M., Barberán-Aceituno, C., Carrillo-Ávila, J. A., López-Guerrero, J. A., & Aguilar-Quesada, R. (2016). DNA Source Selection for Downstream Applications Based on DNA Quality Indicators Analysis. *Biopreservation and Biobanking*, 14(4), 264–270. <https://doi.org/10.1089/bio.2015.0064>
- Malik, J.A., Bhadauria, M. (2020). Cashew Nut (*Anacardium occidentale*). In: Nayik, G.A., Gull, A. (eds) *Antioxidants in Vegetables and Nuts - Properties and Health Benefits*. Springer, Singapore. https://doi.org/10.1007/978-981-15-7470-2_28
- Manjusha, A. V. M., Nair, A. S., & Rethesh, P. K. (2015). Cashew based cropping systems: An effective approach for integrated land use planning in hard lateritic areas of northern kerala. In *Integrated Land Use Planning for Sustainable Agriculture and Rural Development* (pp. 237–244). Apple Academic Press; Scopus. <https://doi.org/10.1201/b18816>
- Mathiang, J. M., Nyirenda, M., & Kamwene, C. (2023). Morphological characterization and genetic diversity of maize (*Zea mays* L.) landraces from Zambia. *Journal of Agricultural Science*, 15(3), 112-128.
- Matos-Filho, C. H. A., De Carvalho-Junior, J., Costa, G., Costa, M., Nunes, J., Lopes, Â., & Ferreira-Gomes, R. (2023a). Estimation of the extent of phenotypic diversity among cashew (*Anacardium occidentale* L.) genotypes based on agro-morphological and physicochemical traits. *Plant Genetic Resources: Characterisation and Utilisation*, 175. <https://doi.org/10.1017/S1479262123000102>
- Mint Abdelaziz, S., Medraoui, L., Alami, M., Pakhrou, O., Makkaoui, M., Ould Mohamed Salem Boukhary, A., & Filali-Maltouf, A. (2020). Inter simple sequence repeat markers to assess genetic diversity of the desert date (*Balanites aegyptiaca* Del.) for Sahelian ecosystem restoration. *Scientific Reports*, 10(1). <https://doi.org/10.1038/s41598-020-71835-9>
- Mog, B., Janani, P., Nayak, M. G., Adiga, J. D., & Meena, R. (2019). Manipulation of vegetative growth and improvement of yield potential of cashew (*Anacardium occidentale* L.) by Paclobutrazol. *Scientia Horticulturae*, 257. Scopus. <https://doi.org/10.1016/j.scienta.2019.108748>
- Mog, B., Adiga, J. D., Preethi, P., & Nayak, M. G. (2023). Changes in Growth and Developmental Stages of Nuts of Selected Cashew (*Anacardium occidentale*) Varieties in West Coast Region of Karnataka. *Agricultural Research*, 12(1), 62–71. Scopus. <https://doi.org/10.1007/s40003-022-00620-z>
- Mohammadi, S. A., & Prasanna, B. M. (2003). Analysis of genetic diversity in crop plants—salient statistical tools and considerations. *Crop Science*, 43(4), 1235–1248. <https://doi.org/10.2135/cropsci2003.1235>
- Mohod, A., Jain, S., Powar, A. G. (2011). Cashew nut processing: sources of environmental pollution and standards. *BioInfo Env. Pollution*, 1 (1): 5-11

- Mondini, L., Noorani, A., & Pagnotta, M. A. (2009). Assessing plant genetic diversity by molecular tools. *Diversity*, 1(1), 19–35. <https://doi.org/10.3390/d1010019>
- Morton, J. (1987). *Fruits of Warm Climates*. Julia F Morton.
- Mukhebi, D. W., Gachanja, P. W., Karan, D. J., Kamau, B. M., King'ori, P. W., Juma, B. S., & Mbinda, W. M. (2025). DArTseq-based silicoDArT and SNP markers reveal the genetic diversity and population structure of Kenyan cashew (*Anacardium occidentale* L.) landraces. *PLoS ONE*, 20(1 January). Scopus. <https://doi.org/10.1371/journal.pone.0313850>
- Mukhlif, F. H., Ramadan, A. S. A., Hammody, D. T., Mousa, M. O., & Shahatha, S. S. (2023). Molecular assessment of genetic divergence among maize genotypes. *Sabrao Journal of Breeding and Genetics*, 55(3), 739–748. Scopus. <https://doi.org/10.54910/sabrao2023.55.3.12>
- Muniz, C. R., Freire, F. C. O., Soares, A. A., Cooke, P. H., & Guedes, M. I. F. (2013). The ultrastructure of shelled and unshelled cashew nuts. *Micron*, 54–55, 52–56. Scopus. <https://doi.org/10.1016/j.micron.2013.08.006>
- Nakasone, H.K. and Paull, R.E. 1998. Tropical Fruits. CAB International, Wallingford, 132–148
- Nalini, E., Jawali, N., & Bhagwat, S. G. (2003). *A Simple Method For Isolation Of DNA From Plants Suitable For Long Term Storage And DNA Marker Analysis* (pp. 208–214). Molecular Biology and Agriculture Division Bhabha Atomic Research Centre.
- Nanami, S. (2023). Multiplexed ISSR Genotyping by Sequencing (MIG-Seq). In: Shavrukov, Y. (eds) Plant Genotyping. Methods in Molecular Biology, vol 2638. Humana, New York, NY. https://doi.org/10.1007/978-1-0716-3024-2_29
- Nayak, M. G., Mohana, G. S., Bhat, P. S., Saroj, P. L., Swamy, K. R. M., & Bhat, M. G. (2014). *Catalogue-V Minimum Descriptors of Cashew Germplasm Accessions*. ICAR - Directorate of Cashew Research.
- N'Djolossè, K., Adoukonou-Sagbadja, H., Gbèmavo, C. D. S. J., Kodjo, S., Badou, A., Maliki, R., & Ahoyo Adjovi, R. N. (2019). Agro-morphological characterization of preselected cashew (*Anacardium occidentale* L.) mother trees in Benin farmer's plantations. *Journal of Agriculture and Environment for International Development*, 113(1), 17–34. <https://doi.org/10.12895/jaeid.20191.841>
- Nuri, R, N, I., & Lak, L,H,N, E. (2016). Strategi peningkatan perekonomian masyarakat melalui diversifikasi produk makanan olahan buah jambu mete sebagai basis produk unggulan kabupaten wonogiri. Prosiding seminar nasional: *Pemberdayaan Sektor Riil Melalui Pengembangan Keuangan*.2016: 159-177
- Orwa, C., & Simons, A. J. (2009). *ORWA et al. Pdf maxlike View project COST Action CA15226-Climate-Smart Forestry in Mountain Regions (CLIMO) View project*. <http://www.worldagroforestry.org/output/agroforestree->

- Qian, W., S. Ge & D.Y. Hong, 2001. Genetic variation within and among populations of a wild rice *Oryza granulata* from China detected by RAPD and ISSR markers. *Theor Appl Genet* 102:440–449.
- Pereira, L.D., Costa, M.L., Pinto, J.F.N., Assunção, H.F. 2016. Diversidade genética do florescimento de *Anacardium humile* A. St. Hill no Sudoeste Goiano. *Rev. Bras. Agropecu. Sustent.* 6(4): 19-25. <http://dx.doi.org/10.21206/rbas.v6i4.373>
- Pereira, A. L. F., & Rodrigues, S. (2012). Fermentation and cashew apple juice. In *Handb. Of Plant-Based Fermented Food and Beverage Technology, Second Edition* (pp. 237–248). CRC Press; Scopus. <https://doi.org/10.1201/b12055>
- Pessoni, L. A. (2007). Genetic diversity in cashew (*Anacardium* spp.): A review. *Genetics and Molecular Research*, 6(4), 1035-1048.
- Pissard, A., Arbizu, C., Ghislain, M., Faux, A.-M., Paulet, S., & Bertin, P. (2007). Congruence between morphological and molecular markers inferred from the analysis of the intra-morphotype genetic diversity and the spatial structure of *Oxalis tuberosa* Mol. *Genetica*, 132(1), 71–85. <https://doi.org/10.1007/s10709-007-9150-9>
- Pitono, J., Syakir, M., Nurhayati, H., Sugiartini, E., & Suhirman, S. (2024). Hydraulic Redistribution: Co-Benefit Potential on Cashew (*Anacardium occidentale* L.) Cultivation—A Review. In H. Kurniawan, B. A. Atmoko, W. Harsonowati, W. null, S. Widodo, M. F. Hudaya, R. D. Purba, & E. M. Putri (Eds.), *AIP Conference Proceedings* (Vol. 2957, Issue 1). American Institute of Physics Inc. <https://doi.org/10.1063/5.0184333>
- Prevost, A. & M.J. Wilkinson, 1999. A new system of comparing PCR primers applied to ISSR fingerprinting of potato cultivars. *Theor Appl Genet* 98: 107–112.
- Puntieri, J., Torres, C., & Ghirardi, S. (2013). Growth and branching in *Nothofagus alpina* and *Nothofagus obliqua* (Nothofagaceae) under different light conditions. *Boletín de la Sociedad Argentina de Botánica*, 48(2), 277–293. Scopus. <https://doi.org/10.31055/1851.2372.v48.n2.6262>
- Ramteke, V., Nirala, Y. S., Nayak, M. G., & Mohana, G. S. (2020). Evaluation of apple and nut characters of cashew germplasm from Bastar region, Chhattisgarh, India. *Journal of Plantation Crops*, 48(2), 142–145. <https://doi.org/10.25081/jpc.2020.v48.i2.6374>
- Randriani, E., Listyati, D., & Syafaruddin. (2011). Kekerabatan Plasma Nutfah Jambu Mete (*Anacardium occidentale* L.) Berdasarkan Marka Random Amplified Polymorphic Dna (RAPD). *Buletin RISTRI*, 2(2), 143–150.
- Randriani, E., Tresniawati, & Syafaruddin. (2012). Pemanfaatan Teknik Random Amplified Polymorphic DNA (RAPD) Untuk Pengelompokan Secara Genetik Plasma Nutfah Jambu Mete. *Buletin RISTRI*, 3(1), 1–6.
- Rocha, S. S., Oliveira, A. C., & Santos, R. P. (2020). Congruence between morphological and molecular characterization in plants. *Genetica*, 148(2), 89-98.

- Rosman, R. (2018). *Increasing National Cashew Production Through Improved Ecology-Based Cultivation Technology*. 17(2), 166–174. <https://doi.org/10.21082/psp.v17n2.2018>
- Rostiana, O., Haryudin, W., & Darajat, J. (2017). The Spreading of Cashew Superior Variety-seeds in the Eastern and Western Regions of Indonesia. *Buletin Littro*, 28(1), 1–14.
- Rucińska, A., Olszak, M., Świerszcz, S., Nobis, M., Zubek, S., Kusza, G., Boczkowska, M., & Nowak, A. (2021). Looking for hidden enemies of metabarcoding: Species composition, habitat and management can strongly influence dna extraction while examining grassland communities. *Biomolecules*, 11(2), 1–19. Scopus. <https://doi.org/10.3390/biom11020318>
- S R, V. K., Vermani, A., Rangarajan, V., & Manjare, S. D. (2025). Production of Feni Beverage from Cashew Apple and Assessment of Bioethanol Production Potential from the Residual Pulp. In *Ref. Ser. Phytochem.: Vol. Part F3607* (pp. 1101–1128). Springer Science and Business Media B.V.; Scopus. https://doi.org/10.1007/978-3-031-38663-3_185
- Sabran, M., Lestari, P., & Mastur, M. (2019). Conservation and Sustainable Use of PGRFA in Indonesia. *Buletin Plasma Nutfah*, 25(2), 67. <https://doi.org/10.21082/blpn.v25n2.2019.p67-78>
- Salehi, B., Gültekin-Özgüven, M., Kırkın, C., Özçelik, B., Morais-Braga, M. F. B., Carneiro, J. N. P., Bezerra, C. F., Silva, T. G. d., Coutinho, H. D. M., Amina, B., Armstrong, L., Selamoglu, Z., Sevindik, M., Yousaf, Z., Sharifi-Rad, J., Muddathir, A. M., Devkota, H. P., Martorell, M., Jugran, A. K., Cho, W. C. (2019). *Anacardium* Plants: Chemical, Nutritional Composition and Biotechnological Applications. *Biomolecules*, 9(9), 465. <https://doi.org/10.3390/biom9090465>.
- Samal, S., Lenka, P. C., Nanda, R. M., Nayak, S., Rout, G. R., & Das, P. (2004). Genetic relatedness in cashew (*Anacardium occidentale* L.) germplasm collections as determined by randomly amplified polymorphic DNA. *Genetic Resources and Crop Evolution*, 51(2), 161–166. <https://doi.org/10.1023/B:GRES.0000020856.77201.7d>
- Samal, S., Rout, G. R., & Lenka, P. C. (2003). Analysis of genetic relationships between populations of cashew (*Anacardium occidentale* L.) by using morphological characterisation and RAPD markers. *Plant Soil Environ*, 49(4), 176–182.
- Sankhare, M., Diallo, A. M., Ba Halimatou Sadyaneand Diatta, S., Samb, C. O., & Toure Mamoudou Abdouland Badiane, S. (2025). Phenotypic diversity of growth, leaf and yield-related traits in cashew (*Anacardium occidentale* L.): Implications for the development of a cashew breeding program in Senegal. *GENETIC RESOURCES AND CROP EVOLUTION*. <https://doi.org/10.1007/s10722-025-02367-2>
- Santana, I. B., Paiva, J. R., Barros, L. M., Rosado, A. S., & Peixoto, A. R. (2011). Genetic diversity of cashew (*Anacardium occidentale* L.) germplasm using ISSR markers. *Genetics and Molecular Research*, 10(2), 817–828. <https://doi.org/10.4238/vol10-2gmr1117>

- Savadi, S., Manoj, K., Ashwitha, K., Muralidhara, B. M., Manjunatha, K., Eradasappa, E., Bhagya, H. P., Mohana, G. S., & Adiga, J. D. (2024). Morpho-biochemical and molecular characterization of accessions with unique traits as potential genetic stocks for cashew improvement. *Genetic Resources and Crop Evolution*. <https://doi.org/10.1007/s10722-024-02036-w>
- Schut, J.W., Qi, X., and Stam, P. 1997. Association between relationship measures based on AFLP markers, pedigree data and morphological traits in barley. *Theoretical and Applied Genetics*, 95, 1161- 1168.
- Sethi, K., Dash, M., Panda, P. K., G. S. , M., & Adiga, J. D. (2024). Assessment of adaptability of cashew varieties to changing environmental conditions of Odisha. *Scientia Horticulturae*, 324, 1–9. <https://doi.org/10.1016/j.scienta.2023.112577>
- Shadenova, E. A., Mamirova, A. A., Sembekov, M. T., & Imanbayeva, A. A. (2019). DNA extraction from leaves of woody plants without liquid nitrogen. *Annals of Agri Bio Research*, 24(2), 153–159. Scopus.
- Shahrajabian, M. H., & Sun, W. (2023). The Important Nutritional and Wonderful Health Benefits of Cashew (*Anacardium occidentale* L.). *Natural Products Journal*, 13(4), 2–10. Scopus. <https://doi.org/10.2174/2210315512666220427113702>
- Sheeba, A., & Mohan, S. (2025). Genetic Diversity Studies in Drought Tolerant Rice (*Oryza sativa* L.) Genotypes. *Agricultural Science Digest*, 45(1), 86–90. <https://doi.org/10.18805/ag.D-5434>
- Sika, K., Adoukonou-Sagbadja, H., Ahoton, L., Saidou, A., Ahanchede, A., Kefela, T., Gachomo, E., Baba, M., & Kotchoni, S. O. 2015. Genetic Characterization of Cashew (*Anacardium occidentale* L.) Cultivars from Benin. *Journal of Horticulture*, 02(04), 1–7. <https://doi.org/10.4172/2376-0354.1000153>
- Singh, G. (2010). *Plant systematics: An integrated approach* (3rd ed). Science Publishers.
- Singh, P., Singh, M. P., Sharma, R., & Kumar, A. (2016). Morphological variability and principal component analysis in *Cajanus* species accessions. *International Journal of Plant Research*, 6(2), 45-58.
- Sivaprakasam, S., Chinnaiyan, U., Varatharajan, S., Uthayasurian, E., Chandrasekaran, M., Paramasivam, S., & Ramasamy, P. (2025). Unveiling the Physicochemical Attributes and Antioxidant Potential of Cashew Apple Juice Varieties. *Applied Fruit Science*, 67(2). <https://doi.org/10.1007/s10341-025-01303-z>
- Soelkhan, M., Odit Kudratun, M., Novia Miranda, S., Evanzi, W., Astuti, I., Ahmad, J. A., Fauzan, M., Abdillah, R. D., Ihwanul, A., Rakasiwi, R. G., & Hidayat, A. R. (2022). Pemanfaatan Daging Jambu Mete Sebagai Bahan Baku Pembuatan Abon Di Desa Batu Rakit Kecamatan Bayan Kabupaten Lombok Utara. *Jurnal Sinergi*, 5(1), 27–30.
- Sophian, A., & Syukur, A. (2021). Analysis of Purity and Concentration of Isolated DNA in Making Raw DNA of Rat Species: Analysis of Purity and Concentration of Isolated

DNA in Making Raw DNA of Rat Species. *Eruditio : Indonesia Journal of Food and Drug Safety*, 1(2), 1–5. <https://doi.org/10.54384/eruditio.v1i2.75>

Sudaryanto, A., Hidayat, D. D., Sagita, D., Darmajana, D. A., Indriati, A., & Kurniawan, Y. R. (2024). Engineering Properties of Cashew Nut in Context to Design of Post-Harvest Handling and Processing Machinery. In Setiawan W., Anggono A.D., Hidayati N., & Kusban M. (Eds.), *AIP Conf. Proc.* (Vol. 2838, Issue 1). American Institute of Physics; Scopus. <https://doi.org/10.1063/5.0199495>

Swamy, K. R. M. (2021). Technological Innovations in Cashewnut Production. In *The Basics of Hum. Civilization: Food, Agriculture and Humanity Volume-I : Present Scenario* (pp. 383–417). Taylor and Francis; Scopus. <https://doi.org/10.1201/9781003246237-30>

Syed, N. H., Sureshsundar, S., Wilkinson, M. J., Bhau, B. S., Cavalcanti, J. J. V., & Flavell, A. J. (2005). Ty1-copia retrotransposon-based SSAP marker development in cashew (*Anacardium occidentale* L.). *Theoretical and Applied Genetics*, 110(7), 1195–1202. <https://doi.org/10.1007/s00122-005-1948-1>

Tamburaka, I. P. (2016). Strategi Pengembangan Jambu Mete Untuk Mendukung Perekonomian Petani Di Kabupaten Muna. *Jurnal Mega Aktiva*, 5(1).

Tautz, D. 1989. Hypervariability of simple sequences as a general source for polymorphic DNA markers. *Nucleic Acids Res* 17(16): 6463-6471.

Thakkar, J. R., Mandavia, M. K., Golakiya, B. A., & Sabara, P. H. (2014). Analysis of genetic diversity among brinjal (*Solanum melongena* L.) genotypes through dominant markers. *Indian Journal of Agricultural Biochemistry*, 27(2), 165–169.

Thimmappaiah, Santhosh, W. G., Shobha, D., & Melwyn, G. S. (2009). Assessment of genetic diversity in cashew germplasm using RAPD and ISSR markers. *Scientia Horticulturae*, 120(3), 411–417. <https://doi.org/10.1016/j.scienta.2008.11.022>

Turrión, D., & Bautista, S. (2025). Tree-architectural response to restoration thinning pattern in post-fire Aleppo pine forests – A modular view of tree plasticity. *Forest Ecology and Management*, 595. Scopus. <https://doi.org/10.1016/j.foreco.2025.123025>

Uller, T., & Leimu, R. (2011). Founder events predict changes in genetic diversity during human-mediated range expansions. *Global Change Biology*, 17(11), 3478–3485. Scopus. <https://doi.org/10.1111/j.1365-2486.2011.02509.x>

Van Beuningen, L.T. and Busch, R.H. 1997. Genetic diversity among North American spring wheat cultivars. In: Analysis of the coefficient of parentage matrix. *Crop Science*, 37, 564-573.

Veena, G. L., Adiga, J. D., Mohana, G., Eradasappa, E., Savadi, S., Vanitha, K., Thondaiman, V., & Shamsudheen, M. (2023). Purple Cashew (*Anacardium occidentale* L.): A Unique Cashew Type. *National Academy Science Letters*, 46(6), 559–562. Scopus. <https://doi.org/10.1007/s40009-023-01307-x>

- Virk, P.S., Zhu, J., Newbury, H.J., Bryan, G.J., Jackson, M.T., & Ford-Lloyd, B.V. 2000 Effectiveness of different classes of molecular markers for classifying and revealing variations in rice (*Oryza sativa*) germplasm. *Euphytica* 112: 275-284.
- Vos P, Hogers R, Bleeker M, Reijans M, Van der Lee T, Hornes M, Frijters A, Pot J, Peleman J, Kuiper M and Zabeau M (1995) AFLP: a new technique for DNA fingerprinting. *Nucleic Acids Res* 23: 4407-4414.
- Wahyuni, S., D.H. Xu, N. Bermawie, H. Tsunematsu & T. Ban. 2004. Skrining ISSR primer studi pendahuluan Kekerabatan antar jahe merah, jahe emprit dan jahe besar. *Buletin TRO*, 15: 33-40.
- Wahyuni, S. (2006). Kekerabatan Plasma Nutfah Jambu Mete Berdasar Sifat Morfologis. *Jurnal Littri*, 12(2), 58–66.
- Wan, Y., Mu, H., Luo, N., Yang, J., Tian, Y., Hong, N., & Dong, H. (2023). Rapid accuracy determining DNA purity and concentration in heavy oils by spectrophotometry methods. *Petroleum Science*, 20(6), 3394–3399. Scopus. <https://doi.org/10.1016/j.petsci.2023.07.019>
- Wang, A., Zhang, P., Liu, X., Liang, J., & Li, W. (2016). Genetic structure and diversity of *Glehnia littoralis*, an endangered medicinal plant in China. *Biochemical Systematics and Ecology*, 66, 265–271. <https://doi.org/10.1016/j.bse.2016.04.019>
- Williams, J.G.K., Kubelik, A.R., Livak, K.J., Rafalski, J.A., & Tingey, S.V. 1990. DNAPolymorphisms amplified by arbitrary primers are useful as genetic markers. *Nucleic Acids Res* 18: 6231-6235.
- Xie, X., Qiu, Y., Ke, L., Zheng, X., Wu, G., Chen, J., Qi, X., & Ahn, S. (2011). Microsatellite primers in red bayberry, *Myrica rubra* (myricaceae). *American Journal of Botany*, 98(4), e93–e95. Scopus. <https://doi.org/10.3732/ajb.1000271>
- Xu, Z., Xiao, D., Shen, B., Xiao, J., Chen, X., & Zhou, Z. (2021). Variation and correlation analyses on leaf color parameters and leaf shape traits of *Cornus hongkongensis* subsp. *Elegans*. *Journal of Plant Resources and Environment*, 30(1), 61–68. Scopus. <https://doi.org/10.3969/j.issn.1674-7895.2021.01.08>
- Zehdi, S., Sakka, H., Rhouma, A., Ould Mohamed Salem, A., Marrakchi, M., & Trifi, M. (2004). Analysis of Tunisian date palm germplasm using simple sequence repeat primers. *African Journal of Biotechnology*, 3(4), 215–219. Scopus. <https://doi.org/10.5897/AJB2004.000-2040>
- Zie, M., Karoui, R., Alabi, T., Jacquet, N., Richel, A., & Blecker, C. (2025). Ultrasound-enhanced extraction and structural dynamics of protein-polysaccharide complexes from cashew apple bagasse: Structure–function relationship. *Food Hydrocolloids*, 171. Scopus. <https://doi.org/10.1016/j.foodhyd.2025.111843>
- Zietkiewicz, E., Rafalski, A. & Labuda, D. 1994. Genome finger-printing by simple sequence repeat (SSR)-anchored polymerase chain reaction amplification. *Genomics* 20: 176–183