

DAFTAR PUSTAKA

- Dušejovská M, Vecka M, Rychlík I, Žák A. Dyslipidemia in patients with chronic kidney disease: Etiology and management. *Vnitr Lek.* 2020;66(5):275–81.
- Acosta-Ochoa I, Bustamante-Munguira J, Mendiluce-Herrero A, Bustamante-Bustamante J, Coca-Rojo A. Impact on outcomes across KDIGO-2012 AKI criteria according to baseline renal function. *J Clin Med.* 2019;8(9):19–23.
- Bi JB, Zhang J, Ren YF, Du ZQ, Wu Z, Lv Y, et al. Neutrophil-to-lymphocyte ratio predicts acute kidney injury occurrence after gastrointestinal and hepatobiliary surgery. *World J Gastrointest Surg.* 2020;12(7):326–35.
- Makris K, Spanou L. Acute Kidney Injury : Definition , Pathophysiology and Clinical Phenotypes. 2016;37(2):85–98.
- MacEdo E, Malhotra R, Bouchard J, Wynn SK, Mehta RL. Oliguria is an early predictor of higher mortality in critically ill patients. *Kidney Int.* 2011;80(7):760–7.
- Forget P, Khalifa C, Defour JP, Latinne D, Van Pel MC, De Kock M. What is the normal value of the neutrophil-to-lymphocyte ratio? *BMC Res Notes.* 2017;10(1):1–4.
- Abecassis M, Bartlett ST, Collins AJ, Davis CL, Delmonico FL, Friedewald JJ, et al. Kidney transplantation as primary therapy for end-stage renal disease: A National Kidney Foundation/Kidney Disease Outcomes Quality Initiative (NKF/KDOQI™) conference. *Clinical Journal of the American Society of Nephrology.* 2008;3(2):471–80.
- Kemenkes RI. Hasil Riset Kesehatan Dasar Tahun 2018. Kementrian Kesehatan RI. 2018;53(9):1689–99.
- Boveland E, Weltings S, Rad M, Van Kampen P, Pelger RCM, Roshani H. The Influence of Pain on the Outcome of Extracorporeal Shockwave Lithotripsy. *Curr Urol.* 2019;12(2):81–7.

- Inker LA, Astor BC, Fox CH, Isakova T, Lash JP, Peralta CA, et al. KDOQI US commentary on the 2012 KDIGO clinical practice guideline for the evaluation and management of CKD. *American Journal of Kidney Diseases*. 2014;63(5):713–35.
- Son HE, Ryu JY, Go S, Yi Y, Kim K, Oh YK, et al. Association of ambulatory blood pressure monitoring with renal outcome in patients with chronic kidney disease. *Kidney Res Clin Pract*. 2020;39(1):70–80.
- Gilbert J, Lovibond K, Mooney A, Dudley J. Renal replacement therapy: Summary of NICE guidance. *BMJ (Online)*. 2018;363(October):1–4.
- Hall KS, Kusunoki Y, Gatny H, Barber J. Stress symptoms and frequency of sexual intercourse among young women. *Journal of Sexual Medicine*. 2014;11(8):1982–90.
- Ishisawa N. Complications of renal transplantation. *The Japanese Journal of Nephrology*. 1971;13(1):93.
- Er RE, Okyay GU. KIDNEY DISEASES Comparison Between RIFLE , AKIN , and KDIGO : Acute Kidney Injury Definition Criteria for Prediction of In-hospital Mortality in Critically Ill Patients.
- Ostermann M, Joannidis M. Acute kidney injury 2016: Diagnosis and diagnostic workup. *Crit Care*. 2016;20(1):1–13.
- Grynberg K, Polkinghorne KR, Ford S, Stenning F, Lew TE, Barrett JA, et al. Early serum creatinine accurately predicts acute kidney injury post cardiac surgery. *BMC Nephrol*. 2017;18(1):1–7.
- Qureshi AI, Huang W, Lobanova I, Hanley DF, Hsu CY, Malhotra K, et al. Systolic Blood Pressure Reduction and Acute Kidney Injury in Intracerebral Hemorrhage. *Stroke*. 2020;(October):3030–8.
- Palmisano A, Gandolfini I, Delsante M, Cantarelli C, Fiaccadori E, Cravedi P, et al. Acute kidney injury (Aki) before and after kidney transplantation: Causes, medical approach, and implications for the long-term outcomes. *J Clin Med*. 2021;10(7).

Washburn KK, Zappitelli M, Arikan AA, Loftis L, Yalavarthy R, Parikh CR, et al. Urinary interleukin-18 is an acute kidney injury biomarker in critically ill children. *Nephrology Dialysis Transplantation*. 2008;23(2):566–72.

Kaddourah A, Basu RK, Goldstein SL, Sutherland SM. Oliguria and acute kidney injury in critically ill children: Implications for diagnosis and outcomes. *Pediatric Critical Care Medicine*. 2019;20(4):332–9.

Cooper JE, Wiseman AC. Acute kidney injury in kidney transplantation. 2013;

Jawdeh BGA, Govil A. Acute Kidney Injury in Transplant Setting : Differential Diagnosis and Impact on Health and Health Care. *Adv Chronic Kidney Dis*. 2017;24(4):228–32.

Schiffl H, Lang SM. Update on Biomarkers of Acute Kidney Injury. *Mol Diagn Ther*. 2012;16(4):199–207.

Liu J xing, Yang C, Zhang W huang, Su H yong, Liu Z jian, Pan Q, et al. Disturbance of mitochondrial dynamics and mitophagy in sepsis-induced acute kidney injury. *Life Sci*. 2019;235(July):116828.

Gaytán-Campos IM, Chávez-Meza A, Juárez-Sánchez JO, et al. Elevated neutrophil-lymphocyte ratio and delayed graft function after kidney transplantation. *Zenodo*. 2025 Mar 10;v1.

Kania E, Wiśniewska N, Kania A, Głowiński J. The impact of hematological indices on the occurrence of delayed graft function after kidney transplantation. *J Clin Med*. 2023;12(24):7514.

Zhang Y, Liu R, Zhao X, Ou Z, Wang S, Huang K, et al. Dynamic changes of neutrophil-to-lymphocyte ratio in brain-dead donors and delayed graft function in kidney transplant recipients. *Ren Fail*. 2022;44(1):1907–1913.

van den Berg E, Geleijnse JM, Brink EJ, et al. Chronic low-grade inflammation in renal transplant recipients is associated with reduced renal function. *Am J Clin Nutr*. 2011;93(3):641–647.

Kwon YJ, Lee YJ, Park BJ, et al. Association of preoperative prognostic nutritional index with early graft function after kidney transplantation. *Transplant Proc.* 2020;52(6):1751–1756.

Işıktas Sayilar E, Yilmaz VT, Yilmaz S, et al. The post-transplant neutrophil-to-lymphocyte ratio and platelet-to-lymphocyte ratio in relation to graft function among renal transplant recipients. *Ann Transplant.* 2020;25:e919216.

Kolonko A, Słabiak-Błaż N, Karkoszka H, et al. Neutrophil-to-lymphocyte ratios in differential diagnosis of kidney transplant acute rejection. *Ann Transplant.* 2022;27:e937239.

Bu X, Zhang L, Chen P, Wu X. Relation of neutrophil-to-lymphocyte ratio to acute kidney injury in patients with sepsis and septic shock: A retrospective study. *Int Immunopharmacol.* 2019 May 1;70:372–7.

Yilmaz H, Cakmak M, Inan O, Darcin T, Akcay A. Can neutrophil-lymphocyte ratio be independent risk factor for predicting acute kidney injury in patients with severe sepsis? *Ren Fail.* 2015 Mar 1;37(2):225–9.

Tang Y, Chen L, Li B, Yang L, Ouyang W, Li D. Preoperative Neutrophil-Lymphocyte Ratio for predicting surgery-related acute kidney injury in non-cardiac surgery patients under general anaesthesia: A retrospective cohort study. *PLoS One.* 2022 Jul 1;17(7 July).

Andrade-Oliveira V, Foresto-Neto O, Watanabe IKM, Zatz R, Câmara NOS. Inflammation in renal diseases: New and old players. *Front Pharmacol.* 2019;10.

Zhang WR, Garg AX, Coca SG, Devereaux PJ, Eikelboom J, Kavsak P, et al. Plasma IL-6 and IL-10 concentrations predict AKI and long-term mortality in adults after cardiac surgery. *Journal of the American Society of Nephrology.* 2015 Dec 1;26(12):3123–32.

Valga F, Monzón T, Henriquez F, Antón-Pérez G. Índices neutrófilo-linfocito y plaqueta-linfocito como marcadores biológicos de interés en la enfermedad renal. *Nefrología*. 2019 May;39(3):243–9.

Kim WH, Park JY, Ok SH, Shin IW, Sohn JT. Association between the neutrophil/lymphocyte ratio and acute kidney injury after cardiovascular surgery:a retrospective observational study. *Medicine (United States)*. 2015;94(43).

Palmisano A, Gandolfini I, Delsante M, Cantarelli C, Fiaccadori E, Cravedi P, et al. Acute kidney injury (Aki) before and after kidney transplantation: Causes, medical approach, and implications for the long-term outcomes. Vol. 10, *Journal of Clinical Medicine*. MDPI; 2021.