

DAFTAR PUSTAKA

- Aldrian, E., & Susanto, R. D. (2003). Identification of Three Dominant Rainfall Regions Within Indonesia and Their Relationship to Sea Surface Temperature. *International Journal of Climatology*, 23(12), 1435–1452. <https://doi.org/10.1002/joc.950>
- Amone-Mabuto, M., Bandeira, S., Hollander, J., Hume, D., Campira, J., & Adams, J. B. (2024). Monitoring seagrass meadows in Maputo Bay using integrated remote sensing techniques and machine learning. *Regional Studies in Marine Science*, 79, 1–11. <https://doi.org/10.1016/j.rsma.2024.103816>
- Anggoro, A., Siregar, V. P., Bahri Agus, S., & Agus, S. B. (2016). The Effect of Sunlight on Benthic Habitats Mapping in Pari Island Using Worldview-2 Imagery. *Procedia Environmental Sciences*, 33(April), 487–495. <https://doi.org/10.1016/j.proenv.2016.03.101>
- Badan Meteorologi, Klimatologi, dan Geofisika. (2022). *Pemanfaatan Data Iklim di Stasiun Klimatologi Kairatu, Ambon*. <https://content.bmkg.go.id/wp-content/uploads/PEMANFAATAN-DATA-IKLIM-DI-STASIUN-KLIMATOLOGI-KAIRATU-AMBON.pdf>
- Badan Pusat Statistik Provinsi Maluku. (2025). *Jumlah Penduduk Menurut Kabupaten/Kota di Maluku (Jiwa), 2025*. <https://maluku.bps.go.id/id>
- Barazzetti, L. (2024). Spatio-temporal analysis of georeferenced time-series applied to structural monitoring. *Journal of Civil Structural Health Monitoring*, 14(1), 163–188. <https://doi.org/10.1007/s13349-023-00743-z>
- Belgiu, M., & Drăguț, L. (2016). Random forest in remote sensing: A review of applications and future directions. *ISPRS Journal of Photogrammetry and Remote Sensing*, 114, 24–31. <https://doi.org/10.1016/j.isprsjprs.2016.01.011>
- Benesty, J., Chen, J., Huang, Y., & Cohen, I. (2009). Pearson correlation coefficient. In *Noise Reduction in Speech Processing* (pp. 1–4). Springer.

https://doi.org/10.1007/978-3-642-00296-0_5

Br Ginting, D. N., Wicaksono, P., & Farda, N. M. (2024). MAPPING SEAGRASS PERCENT COVER AND BIOMASS IN NUSA LEMBONGAN, BALI, INDONESIA. *Geography, Environment, Sustainability*, 17(1), 16–27.
<https://doi.org/10.24057/2071-9388-2023-2886>

Bray, J. R., & Curtis, J. T. (1957). An ordination of the upland forest communities of southern Wisconsin. *Ecological Monographs*, 27(4), 325–349.
<https://doi.org/10.2307/1942268>

Breiman, L. (2001). Random Forests. *Machine Learning*, 45(1), 5–32.
<https://doi.org/10.1023/A:1010933404324>

Campbell, J. B., & Wynne, R. H. (2011). *Introduction to Remote Sensing*. Guilford Publications. <https://books.google.co.id/books?id=zgQDZEya6foC>

Chai, T., & Draxler, R. R. (2014). Root mean square error (RMSE) or mean absolute error (MAE)? *Geoscientific Model Development*, 7(3), 1247–1250.
<https://doi.org/10.5194/gmd-7-1247-2014>

Choto, M., Higa, H., Salem, S., Siswanto, E., Suzuki, T., & Mäll, M. (2025). Performance Evaluation of Inherent Optical Property Algorithms and Identification of Potential Water Quality Indicators Using GCOM-C Data in Eutrophic Lake Kasumigaura, Japan. *Remote Sensing*, 17(9), 1621.
<https://doi.org/10.3390/rs17091621>

Cleveland, R. B. (1990). STL : A Seasonal-Trend Decomposition Procedure Based on Loess. *Journal of Official Statistics*, 6(1), 3-73.

Cullen-Unsworth, L., & Unsworth, R. (2013). Seagrass meadows, ecosystem services, and sustainability. *Environment*, 55(3), 14–28.
<https://doi.org/10.1080/00139157.2013.785864>

Danoedoro, P. (2009). *LAND-USE INFORMATION FROM THE SATELLITE*

IMAGERY: Versatility and Contents for Local Physical Planning. LAP
LAMBERT Academic Publishing.

Danoedoro, P. (2012). *Pengantar Penginderaan Jauh Digital* (1st ed.). Andi.

Davies, B. F. R., Oiry, S., Rosa, P., Zoffoli, M. L., Sousa, A. I., Thomas, O. R., Smale, D. A., Austen, M. C., Biermann, L., Attrill, M. J., Roman, A., Navarro, G., Barillé, A. L., Harin, N., Clewley, D., Martinez-Vicente, V., Gernez, P., & Barillé, L. (2024). A sentinel watching over inter-tidal seagrass phenology across Western Europe and North Africa. *Communications Earth and Environment*, 5(1), 1–13. <https://doi.org/10.1038/s43247-024-01543-z>

Davies, B., Oiry, S., Rosa, P., Zoffoli, M., Sousa, A., Thomas, O., Smale, D., Austen, M., Biermann, L., Attrill, M., Román, A., Navarro, G., Barillé, A.-L., Harin, N., Clewley, D., Martinez-Vicente, V., Gernez, P., & Barillé, L. (2024). Intertidal seagrass extent from Sentinel-2 time-series show distinct trajectories in Western Europe. *Remote Sensing of Environment*. <https://doi.org/10.1016/j.rse.2024.114340>

Dimitriadis, S. I., & Liparas, D. (2018). How random is the random forest? Random forest algorithm on the service of structural imaging biomarkers for Alzheimer's disease: From Alzheimer's disease neuroimaging initiative (ADNI) database. In *Neural Regeneration Research* (Vol. 13, Issue 6, pp. 962–970). Wolters Kluwer Medknow Publications. <https://doi.org/10.4103/1673-5374.233433>

Drusch, M., Del Bello, U., Carlier, S., Colin, O., Fernandez, V., Gascon, F., Hoersch, B., Isola, C., Laberinti, P., Martimort, P., Meygret, A., Spoto, F., Sy, O., Marchese, F., & Bargellini, P. (2012). Sentinel-2: ESA's Optical High-Resolution Mission for GMES Operational Services. *Remote Sensing of Environment*, 120, 25–36. <https://doi.org/10.1016/j.rse.2011.11.026>

Duarte, C. M., Dennison, W. C., Orth, R. J. W., & Carruthers, T. J. B. (2008). The charisma of coastal ecosystems: Addressing the imbalance. In *Estuaries and Coasts* (Vol. 31, Issue 2, pp. 233–238). Springer New York.

<https://doi.org/10.1007/s12237-008-9038-7>

Duarte, C. M., & Kirkman, H. (2001). *Chapter 7 - Methods for the measurement of seagrass abundance and depth distribution* (F. T. Short & R. G. B. T.-G. S. R. M. Coles (eds.); pp. 141–153). Elsevier Science.
<https://doi.org/https://doi.org/10.1016/B978-044450891-1/50008-6>

Duarte, C. M., Marba, N., Agawin, N., Cebrian, J., Enriquez, S., Fortes, M. D., Gallegos, M. E., Merino, M., Olesen, B., & Sandjensen, K. (1994). Reconstruction of seagrass dynamics - Age determinations and associated tools for the seagrass ecologist. *Marine Ecology Progress Series*, 107(1–2), 195.
<https://doi.org/10.3354/meps107195>

European Space Agency. (2024). *Sentinel-2 Mission*.

Fauzan, M. A., Wicaksono, P., & Hartono. (2021). Characterizing Derawan seagrass cover change with time-series Sentinel-2 images. *Regional Studies in Marine Science*, 48. <https://doi.org/10.1016/j.rsma.2021.102048>

Foody, G. M. (2002). Status of land cover classification accuracy assessment. *Remote Sensing of Environment*, 80(1), 185–201. [https://doi.org/10.1016/S0034-4257\(01\)00295-4](https://doi.org/10.1016/S0034-4257(01)00295-4)

Friedl, M. A., Henebry, G. M., Reed, B. C., Huete, A., White, M. A., Morisette, J. T., Nemani, R. R., Zhang, X., & Myneni, R. B. (2006). *Land Surface Phenology*.

Gao, B.-C., & Li, R.-R. (2021). Correction of Sunlint Effects in High Spatial Resolution Hyperspectral Imagery Using SWIR or NIR Bands and Taking Account of Spectral Variation of Refractive Index of Water. *Advances in Environmental and Engineering Research*, 02(03), 17.
<https://doi.org/10.21926/aeer.2103017>

Green, E. P., Mumby, P., & Edwards, Alasdair James Clark, C. D. (2000). *Remote Sensing Handbook for Tropical Coastal Management*.
<https://www.researchgate.net/publication/257395007>

- Grolemund, G., & Wickham, H. (2011). Dates and times made easy with lubridate. *Journal of Statistical Software*, 40(3), 1–25. <https://doi.org/10.18637/jss.v040.i03>
- Ha, N. T., Manley-Harris, M., Pham, T. D., & Hawes, I. (2020). A Comparative Assessment of Ensemble-Based Machine Learning and Maximum Likelihood Methods for Mapping Seagrass Using Sentinel-2 Imagery in Tauranga Harbor, New Zealand. *Remote Sensing*, 12(3), 1–16. <https://doi.org/10.3390/rs12030355>
- Hanes, J., Li, L., & Morisette, J. (2013). Land surface phenology. In *Biophysical Applications of Satellite Remote Sensing* (pp. 99–125). Springer-Verlag Berlin Heidelberg. https://doi.org/10.1007/978-3-642-25047-7_4
- Harahap, S. D., & Wicaksono, P. (2024). Relative Water Column Correction Methods for Benthic Habitat Mapping in Optically Shallow Coastal Water. *Advances in Science, Technology and Innovation*, 181–183. https://doi.org/10.1007/978-3-031-43218-7_42
- Harjuna, R. A., Riniatsih, I., & Suryono, C. A. (2020). Kondisi Padang Lamun di Pulau Panjang dan Pulau Lima, Banten. *Journal of Tropical Marine Science*, 3(2), 89–93. <https://doi.org/10.33019/jour.trop.mar.sci.v3i2.1928>
- Hartoni, Siregar, V. P., Wouthuyzen, S., & Agus, S. B. (2022). Object based classification of benthic habitat using Sentinel 2 imagery by applying with support vector machine and random forest algorithms in shallow waters of Kepulauan Seribu, Indonesia. *Biodiversitas*, 23(1), 514–520. <https://doi.org/10.13057/biodiv/d230155>
- Hastie, T., Tibshirani, R., & Friedman, J. (2009). *The Elements of Statistical Learning: Data Mining, Inference, and Prediction* (2nd ed.). Springer. <https://doi.org/10.1007/978-0-387-84858-7>
- Hedley, J. D., Harborne, A. R., & Mumby, P. J. (2005). Simple and robust removal of sun glint for mapping shallow-water benthos. *International Journal of Remote Sensing*, 26(10), 2107–2112. <https://doi.org/10.1080/01431160500034086>

- Helmi, M., Hartoko, A., S., H., Munasik, & Wouthuyzen, S. (2011). Analisis Respon Spektral dan Ekstraksi Nilai Spektral Terumbu Karang Pada Citra Digital Multispektral Satelit ALOS-AVNIR di Perairan Gugus Pulau Pari, Kepulauan Seribu, Jakarta. *Buletin Oseanografi Marina*, 1, 10–12.
- Hijmans, R. J. (2024). *terra: Spatial Data Analysis*. <https://cran.r-project.org/package=terra>
- Huang, Z., Brooke, B., & Harris, P. (2011). A new approach to mapping marine benthic habitats using physical environmental data. *Continental Shelf Research*, 31. <https://doi.org/10.1016/J.CSR.2010.03.012>
- Jayanti, A. R. (2020). MANFAAT PADANG LAMUN SEBAGAI PENYEIMBANG EKOSISTEM LAUT DI PULAU PRAMUKA, KEPULAUAN SERIBU. *JURNAL GEOGRAFI Geografi Dan Pengajarannya*. <https://api.semanticscholar.org/CorpusID:225849359>
- Jönsson, P., & Eklundh, L. (2004). TIMESAT—A program for analyzing time-series of satellite sensor data. *Computers & Geosciences*, 30(8), 833–845. <https://doi.org/10.1016/j.cageo.2004.05.006>
- Kirk, J. T. O. (2011). *Light and Photosynthesis in Aquatic Ecosystems* (3rd ed.). Cambridge University Press. <https://doi.org/10.1017/CBO9781139168212>
- Kohler, K. E., & Gill, S. M. (2006). Coral Point Count with Excel extensions (CPCe): A Visual Basic program for the determination of coral and substrate coverage using random point count methodology. *Computers and Geosciences*, 32(9), 1259–1269. <https://doi.org/10.1016/j.cageo.2005.11.009>
- Kritzer, J., Delucia, M., Greene, E., Shumway, C., Topolski, M., Thomas-Blate, J., Chiarella, L., Davy, K., & Smith, K. (2016). The Importance of Benthic Habitats for Coastal Fisheries. *BioScience*, 66, 274–284. <https://doi.org/10.1093/BIOSCI/BIW014>
- Kuhn, M. (2008). Building predictive models in R using the caret package. *Journal of*

Statistical Software, 28(5), 1–26. <https://doi.org/10.18637/jss.v028.i05>

Kuhn, M., & Johnson, K. (2013). *Applied Predictive Modeling*. Springer.
<https://doi.org/10.1007/978-1-4614-6849-3>

Kurniawan, A., Khakhim, N., & Wicaksono, P. (2025). Mapping of Nearshore Bathymetry Based on Random Forest Machine Learning in Kemujan Island Waters. *International Journal of Geoinformatics*, 21(2), 25–41.
<https://doi.org/10.52939/ijg.v21i2.3933>

Latuconsina, H., & Dawar, L. (2012). Telaah ekologi komunitas lamun (seagrass) perairan Pulau Osi Teluk Kotania Kabupaten Seram Bagian Barat. *Agrikan: Jurnal Agribisnis Perikanan*, 5(2), 12–19.
<https://doi.org/10.29239/j.agrikan.5.2.12-19>

Legendre, P., & Legendre, L. (2012). *Numerical Ecology* (3rd ed.). Elsevier.

Liaw, A., & Wiener, M. (2002). Classification and regression by randomForest. *R News*, 2(3), 18–22.

Lieth, H. (1973). Phenology in Productivity Studies. In D. E. Reichle (Ed.), *Analysis of Temperate Forest Ecosystems* (pp. 29–46). Springer Berlin Heidelberg.
https://doi.org/10.1007/978-3-642-85587-0_4

Lieth, H. (1974). Purposes of a Phenology Book. In H. Lieth (Ed.), *Phenology and Seasonality Modeling* (pp. 3–19). Springer Berlin Heidelberg.
https://doi.org/10.1007/978-3-642-51863-8_1

Liu, Y., Liu, W., Ye, T., Chen, S., Chen, X., Li, Z., Zhan, N., & Sun, R. (2025). Incorporating climate data with machine learning can improve rice phenology estimation. *Environmental Research Letters*, 20(2), 24020.
<https://doi.org/10.1088/1748-9326/ada56e>

Lyzenga, D. R. (1978). Passive remote sensing techniques for mapping water depth and bottom features. *Applied Optics*, 17(3), 379.

<https://doi.org/10.1364/ao.17.000379>

Lyzenga, D. R. (1981). Remote sensing of bottom reflectance and water attenuation parameters in shallow water using aircraft and landsat data. *International Journal of Remote Sensing*, 2(1), 71–82. <https://doi.org/10.1080/01431168108948342>

Maechler, M., Rousseeuw, P., Struyf, A., Hubert, M., & Hornik, K. (2023). *cluster: Cluster Analysis Basics and Extensions*. <https://cran.r-project.org/package=cluster>

Main-Knorn, M., Pflug, B., Louis, J., Debaecker, V., Müller-Wilm, U., & Gascon, F. (2017). Sen2Cor for Sentinel-2. *Proceedings of SPIE*, 10427, 1042704. <https://doi.org/10.1117/12.2278218>

McKenzie, L. J., Finkbeiner, M. A., & Kirkman, H. (2001). Chapter 5 - Methods for mapping seagrass distribution. In F. T. Short & R. G. B. T.-G. S. R. M. Coles (Eds.), *Global Seagrass Research Methods* (pp. 101–121). Elsevier Science. <https://doi.org/https://doi.org/10.1016/B978-044450891-1/50006-2>

McKenzie, L. J., Yaakub, S. M., Tan, R., Seymour, J., & Yoshida, R. L. (2016). Seagrass habitats of Singapore: Environmental drivers and key processes. *Raffles Bulletin of Zoology*, 2016(Part I), 60–77.

Murtagh, F., & Legendre, P. (2014). Ward's hierarchical agglomerative clustering method: Which algorithms implement Ward's criterion? *Journal of Classification*, 31(3), 274–295. <https://doi.org/10.1007/s00355-014-0381-z>

Nakagawa, S., & Schielzeth, H. (2013). A general and simple method for obtaining R^2 from generalized linear mixed-effects models. *Methods in Ecology and Evolution*, 4(2), 133–142.

Nalepa, J. (2021). Recent Advances in Multi- and Hyperspectral Image Analysis. *Sensors (Basel, Switzerland)*, 21. <https://doi.org/10.3390/s21186002>

Oksanen, J., Simpson, G. L., Blanchet, F. G., Kindt, R., Legendre, P., & others. (2022).

vegan: Community Ecology Package. <https://cran.r-project.org/package=vegan>

Orth, R. J., Carruthers, T. J. B. B., Dennison, W. C., Duarte, C. M., Fourqurean, J. W., Heck, K. L., Hughes, A. R., Kendrick, G. A., Kenworthy, W. J., Olyarnik, S., Short, F. T., Waycott, M., Williams, S. L., Heck Jr., K. L., Hughes, A. R., Kendrick, G. A., Kenworthy, W. J., Olyarnik, S., Short, F. T., ... Williams, S. L. (2006). A global crisis for seagrass ecosystems. *BioScience*, 56(12), 987 – 996. [https://doi.org/10.1641/0006-3568\(2006\)56\[987:AGCFSE\]2.0.CO;2](https://doi.org/10.1641/0006-3568(2006)56[987:AGCFSE]2.0.CO;2)

Pebesma, E. (2018). Simple features for R: Standardized support for spatial vector data. *The R Journal*, 10(1), 439–446. <https://doi.org/10.32614/RJ-2018-009>

Picard, R. R., & Cook, R. D. (1984). Cross-validation of regression models. *Journal of the American Statistical Association*, 79(387), 575–583. <https://doi.org/10.1080/01621459.1984.10478083>

Probst, P., Wright, M. N., & Boulesteix, A.-L. (2019). Hyperparameters and tuning strategies for random forest. *WIREs Data Mining and Knowledge Discovery*, 9(3), e1301. <https://doi.org/10.1002/widm.1301>

Rahman, I., & Astriana, B. H. (2019). Penyuluhan Mengenai Ekosistem Lamun Sebagai Upaya Pelestarian Ekosistem Di Perairan Pantai Sire, Kabupaten Lombok Utara. *Abdi Insani*, 6(2), 245. <https://doi.org/10.29303/abdiinsani.v6i2.245>

Ririhena, R. E., Wattimena, A. Y., & Tuanaya, A. F. (2023). The Impact of Climate Change on the Productivity of Nutmeg (*Myristica fragrans* Houtt.) in Kamal Village, West Kairatu District. *JURNAL AGRIKAN (Agribisnis Perikanan)*, 16(1), 43–49.

Roelfsema, C. M., Lyons, M., Kovacs, E. M., Maxwell, P., Saunders, M. I., Samper-Villarreal, J., & Phinn, S. R. (2014). Multi-temporal mapping of seagrass cover, species and biomass: a semi-automated object based image analysis approach. *Remote Sensing of Environment*, 150, 172–187. <https://doi.org/10.1016/j.rse.2014.05.001>

- Roelfsema, C., & Phinn, S. (2008). Evaluating eight field and remote sensing approaches for mapping the benthos of three different coral reef environments in Fiji. *Remote Sensing of Inland, Coastal, and Oceanic Waters*, 7150, 71500F. <https://doi.org/10.1117/12.804806>
- Roelfsema, C., & Phinn, S. (2009). *A Manual for Conducting Georeferenced Photo Transects Surveys to Assess the Benthos of Coral Reef and Seagrass Habitats*.
- Rousseeuw, P. J. (1987). Silhouettes: A graphical aid to the interpretation and validation of cluster analysis. *Journal of Computational and Applied Mathematics*, 20, 53–65. [https://doi.org/10.1016/0377-0427\(87\)90125-7](https://doi.org/10.1016/0377-0427(87)90125-7)
- Samper-Villarreal, J., Lovelock, C. E., Saunders, M. I., Roelfsema, C., & Mumby, P. J. (2016). Organic carbon in seagrass sediments is influenced by seagrass canopy complexity, turbidity, wave height, and water depth. *Limnology and Oceanography*, 61(3), 938–952. <https://doi.org/10.1002/lno.10262>
- Setyawan, I., Siregar, V. P., Pramono, G. H., & Yuwono, D. M. (2014). PEMETAAN PROFIL HABITAT DASAR PERAIRAN DANGKAL BERDASARKAN BENTUK TOPOGRAFI: Studi Kasus Pulau Panggang, Kepulauan Seribu Jakarta. *Majalah Ilmiah Globë*, 16(2), 125–132. <https://api.semanticscholar.org/CorpusID:132008916>
- Short, F. T., & Duarte, C. M. (2001). *Chapter 8 - Methods for the measurement of seagrass growth and production* (F. T. Short & R. G. B. T.-G. S. R. M. Coles (eds.); pp. 155–182). Elsevier Science. <https://doi.org/https://doi.org/10.1016/B978-044450891-1/50009-8>
- Simanjuntak, T., & Soselisa, H. L. (2021). *Menganyam pengetahuan dari tanah Seram, Maluku, Indonesia* (L. Wattimena (ed.)). PT Pustaka Obor Indonesia. <https://books.google.co.id/books?id=rfCW0QEACAAJ>
- Wang, Y., He, X., Bai, Y., Wang, D., Zhu, Q., Gong, F., Yang, D., & Li, T. (2022). Satellite retrieval of benthic reflectance by combining lidar and passive high-

resolution imagery: Case-I water. *Remote Sensing of Environment*, 272, 112955.
<https://doi.org/10.1016/j.rse.2022.112955>

Ward, J. H. (1963). Hierarchical grouping to optimize an objective function. *Journal of the American Statistical Association*, 58(301), 236–244.
<https://doi.org/10.1080/01621459.1963.10500845>

Wicaksono, P. (2014). THE USE OF IMAGE ROTATIONS ON MULTISPECTRAL-BASED BENTHIC HABITATS MAPPING. *12th Biennial Conference of Pan Ocean Remote Sensing Conference (PORSEC 2014)*, 10.
<https://doi.org/10.13140/2.1.3877.6006>

Wicaksono, P., Hafizt, M., Harahap, S. D., & Nandika, M. R. (2024a). Integrating Sentinel-2 and PlanetScope Image with Drone-based Seagrass Data for Seagrass Percent Cover Mapping. *IOP Conference Series: Earth and Environmental Science*, 1291(1), 12012. <https://doi.org/10.1088/1755-1315/1291/1/012012>

Wicaksono, P., Hafizt, M., Harahap, S. D., & Nandika, M. R. (2024b). Integrating Sentinel-2 and PlanetScope Image with Drone-based Seagrass Data for Seagrass Percent Cover Mapping. *IOP Conference Series: Earth and Environmental Science*, 1291(1). <https://doi.org/10.1088/1755-1315/1291/1/012012>

Wicaksono, P., Harahap, S. D., Hafizt, M., Maishella, A., & Yuwono, D. M. (2023). Seagrass ecosystem biodiversity mapping in part of Rote Island using multi-generation PlanetScope imagery. *Carbon Footprints*, 2(4).
<https://doi.org/10.20517/cf.2023.9>

Wicaksono, P., Lazuardi, W., & Munir, M. (2019). INTEGRATING IMAGE at DIFFERENT SPATIAL RESOLUTIONS and FIELD DATA for SEAGRASS PERCENT COVER MAPPING. *International Archives of the Photogrammetry, Remote Sensing and Spatial Information Sciences - ISPRS Archives*, 42(4/W19), 487–492. <https://doi.org/10.5194/isprs-archives-XLII-4-W19-487-2019>

Wicaksono, P., Maishella, A., Arjasakusuma, S., Lazuardi, W., & Harahap, S. D.

- (2022). Assessment of WorldView-2 images for aboveground seagrass carbon stock mapping in patchy and continuous seagrass meadows. *International Journal of Remote Sensing*, 43(8), 2915–2941. <https://doi.org/10.1080/01431161.2022.2074809>
- Wicaksono, P., Maishella, A., Lazuardi, W., & Muhammad, F. H. (2022). Consistency assessment of multi-date PlanetScope imagery for seagrass percent cover mapping in different seagrass meadows. *Geocarto International*, 37(27), 15161–15186. <https://doi.org/10.1080/10106049.2022.2096122>
- Wicaksono, P., Maishella, A., & Ramadhan. (2025). Capturing the dynamics of aboveground carbon stock in intertidal seagrass meadows using Sentinel-2 time-series imagery. *Remote Sensing Applications: Society and Environment*, 38. <https://doi.org/10.1016/j.rsase.2025.101552>
- Wicaksono, P., Ramadhan, Harahap, S. D., Kartika, C. S. D., & Kamal, M. (2025). Mapping the dynamics of seagrass aboveground carbon stock in a developing coastal area: case study of Kuta Mandalika, Lombok Island, Indonesia. *Environmental Monitoring and Assessment*, 197(10), 1–24. <https://doi.org/10.1007/s10661-025-14534-0>
- Wicaksono, P., Wulandari, S. A., Lazuardi, W., & Munir, M. (2021). Sentinel-2 images deliver possibilities for accurate and consistent multi-temporal benthic habitat maps in optically shallow water. *Remote Sensing Applications: Society and Environment*, 23. <https://doi.org/10.1016/j.rsase.2021.100572>
- Wickham, H. (2016). *ggplot2: Elegant Graphics for Data Analysis*. Springer-Verlag New York. <https://ggplot2.tidyverse.org>
- Wijaya, J., & Wicaksono, P. (2024). *Lifeform-based seagrass species composition mapping of Pari Island using WorldView-2 multispectral imagery*. 78. <https://doi.org/10.1117/12.3009770>
- Zeileis, A., & Grothendieck, G. (2005). zoo: S3 infrastructure for regular and irregular

time series. *Journal of Statistical Software*, 14(6), 1–27.
<https://doi.org/10.18637/jss.v014.i06>

Zoffoli, M. L., Frouin, R., & Kampel, M. (2014). Water column correction for coral reef studies by remote sensing. In *Sensors (Switzerland)* (Vol. 14, Issue 9).
<https://doi.org/10.3390/s140916881>

Zoffoli, M. L., Gernez, P., Rosa, P., Le Bris, A., Brando, V. E., Barillé, A.-L., Harin, N., Peters, S., Poser, K., Spaias, L., Peralta, G., & Barillé, L. (2020). Sentinel-2 remote sensing of *Zostera noltei*-dominated intertidal seagrass meadows. *Remote Sensing of Environment*, 251, 112020.
<https://doi.org/https://doi.org/10.1016/j.rse.2020.112020>