

DAFTAR PUSTAKA

- Aapkes, S.E., Barten, T.R.M., Coudyzer, W., Drenth, J.P.H., Geijselaers, I.M.A., ter Grote, S.A.M., *et al.* (2023). Validation of a semi-automatic method to measure total liver volumes in polycystic liver disease on computed tomography — high speed and accuracy. *Eur. Radiol.* 33 : 3222–3231.
- Agarwal, P.D., Phillips, P., Hillman, L., Lucey, M.R., Lee, F., Mezrich, J.D., *et al.* (2017). Multidisciplinary Management of Hepatocellular Carcinoma Improves Access to Therapy and Patient Survival. *J. Clin. Gastroenterol.* 51 : 845–849.
- Ahmad, J., & Lau, T. (2013). Clinical applications of the Model for End-Stage Liver Disease (MELD) in hepatic medicine. *Clin. Liver Dis.* 2(1) : 1–10.
- Ahmad, M.S., Alkhatib, S.G., & Hjouj, M. (2023). A New Algorithm for Assessing Hepatomegaly Through CT Scan of the Abdomen. *ACM Int. Conf. Proceeding Ser.* 82–87.
- American College of Radiology (2018). CT/MRI LI-RADS v2018. American Collage of Radiology. Reston, VA : 5-18
- Attwa, M.H., & El-Etreby, S.A. (2015). Guide for diagnosis and treatment of hepatocellular carcinoma. *World J. Hepatol.* 7 : 1632–1651.
- Baffy, G. (2015). Decoding multifocal hepatocellular carcinoma: an opportune pursuit. *Hepatobiliary Surg. Nutr.* 4 : 206–20610.
- Bartlett, S., Hong, T., & Valaydon, Z. (2025). Nutritional assessment of patients with liver cirrhosis in the outpatient setting : A narrative review. *Nut*

Article MELD score is the better predictor for 30-day mortality in patients with ruptured hepatocellular carcinoma treated by trans-arterial embolization 11 : 3726–3734.

- Choi, J.Y., Lee, J.M., & Sirlin, C.B. (2014). CT and MR imaging diagnosis and staging of hepatocellular carcinoma: Part I. Development, growth, and spread: Key pathologic and imaging aspects. *Radiology* 272 : 635–654.
- Delis, S.G., Bakoyiannis, A., Biliatis, I., Athanassiou, K., Tassopoulos, N., & Dervenis, C. (2009). Model for end-stage liver disease (MELD) score , as a prognostic factor for post-operative morbidity and mortality in cirrhotic patients , undergoing hepatectomy for hepatocellular carcinoma 351–357.
- Dumont, C., Wuestenberghs, F., Lanthier, N., Piessevaux, H., & Dahlqvist, G. (2022). Malnutrition is highly prevalent in hospitalized cirrhotic patients and associates with a poor outcome 85 : 311–319.
- Eisenhauer, E.A., Therasse, P., Bogaerts, J., Schwartz, L.H., Sargent, D., Ford, R., *et al.* (2009). New response evaluation criteria in solid tumours: Revised RECIST guideline (version 1.1). *Eur. J. Cancer* 45 : 228–247.
- El Serag, Hashem, B. (2013). Epidemiology of Viral Hepatitis and Hepatocellular Carcinoma 142 : 1264–1273.
- Elsawy, A.A., Dawoud, M.M., Elarabawy, R.A., Mohamed, W.S., & Dawoud, R.M. (2020). Role of residual liver volumetry and function in prediction of liver tolerability after transarterial chemoembolization for hepatocellular carcinoma in cirrhotic patients: deriving a clinical decision support score. *Egypt. J. Radiol. Nucl. Med.* 51.
- Favaloro, E.J., Bahraini, M., & Rad, F. (2021). Standardization of Prothrombin Time / International Normalized Ratio (PT / INR) 21–2

- Gomaa, A.I., Khan, S.A., Leen, E.L.S., Waked, I., & Taylor-Robinson, S.D. (2009). Diagnosis of Hepatocellular carcinoma. *World J. Gastroenterol.* 15 : 1301–1314.
- Gotra, A., Sivakumaran, L., Chartrand, G., Vu, K.N., Vandenbroucke-Menu, F., Kauffmann, C., *et al.* (2017). Liver segmentation: indications, techniques and future directions. *Insights Imaging* 8 : 377–392.
- Hasan, I., Loho, I.M., Setiawan, P.B., Djumhana, A., Punmomo, H.D., Siregar, L., *et al.* (2021). Konsensus Nasional Terapi Sistemik Karsinoma Sel Hati di Indonesia, 1st ed. Jakarta : PERHIMPUNAN PENELITIAN HATI INDONESIA.
- Hosmer, D.W., & Lemeshow, S. (2020). Applied Logistic Regression, 2nd ed. New York : John Wiley & Sons.
- Kashkoush, S., El Moghazy, W., Kawahara, T., Gala-Lopez, B., Toso, C., & Kneteman, N.M. (2014). Three-dimensional tumor volume and serum alpha-fetoprotein are predictors of hepatocellular carcinoma recurrence after liver transplantation: Refined selection criteria. *Clin. Transplant.* 28 : 728–736.
- Kelley, R.K., Villanueva, A., Rossi, J.Z., Llovet, J.M., Singal, A.G., Pikarsky, E., *et al.* (2021). Hepatocellular carcinoma 0123456789.
- Kementerian Kesehatan Republik Indonesia (2021). Keputusan Menteri Kesehatan Republik Indonesia Nomor Hk.01.07/Menkes/4722/2021 Tentang Pedoman Nasional Pelayanan Kedokteran Tata Laksana Karsinoma Sel Hati. *Jdih.Kemkes.Go.Id* 1–55.
- Khalidy, N. Al, & Abeerhm Hasan, A.A Jazim, Z.K. (2022). Assessment of malnutrition in patient with liver cirrhosis and its association with dietary

- Lian, D., Wang, W., Liu, L., Wang, J., Rao, S., & Zhou, J. (2022). CT volumetry helps predict prognosis of large hepatocellular carcinoma after resection. *Clin. Radiol.* 77 : e599–e605.
- Ling, Y., Ying, S., Xu, L., Peng, Z., Mao, X., Chen, Z., *et al.* (2022). Automatic volumetric diagnosis of hepatocellular carcinoma based on four-phase CT scans with minimum extra information. *Front. Oncol.* 12 : 1–12.
- Liu, Y., Ran, L., Zhang, H., Ren, H., Jiang, X., Liao, P., *et al.* (2023). Comparison of Child-Pugh , MELD , MELD-Na , and ALBI Scores in Predicting In-Hospital Mortality in Patients with HCC 148–162.
- Llovet, J.M., Kelley, R.K., Villanueva, A., Singal, A.G., Pikarsky, E., Roayaie, S., *et al.* (2021). Hepatocellular carcinoma 0123456789.
- Llovet, J.M., Zucman-rossi, J., Pikarsky, E., Sangro, B., Sherman, M., & Gores, G. (2016). Hepatocellular carcinoma.
- Loho, I.M., Siregar, L., Waspodo, A.S., & Hasan, I. (2018). Current Practice of Hepatocellular Carcinoma Surveillance. *Acta Med. Indones.* 50 : 353–360.
- Mahadevan, V. (2020). Anatomy of the liver 427–431.
- Mcglynn, K.A., Petrick, J.L., & El-serag, H.B. (2021). Epidemiology of Hepatocellular Carcinoma 73 : 4–13.
- Mickey, R.M., & Greenland, S. (1989). The impact of confounder selection criteria on effect estimation. *Am. J. Epidemiol.* 129 : 125–137.
- Montenegro, T.D.E., Ramirez, T.C.A., Mateo C, Y.S., Palma, Govea J. Quiñones, J.C., & Vazquez, O.J.S. (2023). CT-Based Hepatic Residual Volume and Predictors of Outcomes of Patients with Hepatocellular Carcinoma Unsuitable for Surgical Therapy Undergoing Transarterial Chemoembolization.

derived extracellular volume and liver volumetry can predict posthepatectomy liver failure in hepatocellular carcinoma. *Insights Imaging*.

Perhimpunan Peneliti Hati Indonesia (2017). Penatalaksanaan Karsinoma Sel Hati.

Prasad, S.R., Jhaveri, K.S., Saini, S., Hahn, P.F., Halpern, E.F., & Sumner, J.E. (2002). CT tumor measurement for therapeutic response assessment: Comparison of unidimensional, bidimensional, and volumetric techniques - Initial observations. *Radiology* 225 : 416–419.

Reig, M., Forner, A., Rimola, J., Ferrer-Fàbrega, J., Burrel, M., Garcia-Criado, Á., *et al.* (2022). BCLC strategy for prognosis prediction and treatment recommendation: The 2022 update. *J. Hepatol.* 76 : 681–693.

Schuppan, D., & Afdhal, N.H. (2009). Liver Cirrhosis 371 : 838–851.

Shouval, D. (2014). Focus The pros and cons of lamivudine vs . entecavir in decompensated or severe acute exacerbation of chronic hepatitis B. *J. Hepatol.* 60 : 1108–1109.

Siswosudarmo, R.H. (2015). Pendekatan Praktis Penelitian Epidemiologi Klinis dan Aplikasi SPSS untuk Analisis Statistika. Yogyakarta : Departemen Obstetika dan Ginekologi Fakultas Kedokteran UGM-RS Sardjito Yogyakarta.

et al. (2003). Model for End-Stage Liver Disease (MELD) and Allocation of Donor Livers 91–96.

Wu, E.M., Wong, L.L., Hernandez, B.Y., Ji, J., Jia, W., & Kwee, S.A. (2018). Gender differences in hepatocellular cancer : disparities in nonalcoholic fatty liver disease / steatohepatitis and liver transplantation.

Zakaria, H.M., Macshut, M., Gaballa, N.K., Sherif, A.E., Abdel-Samea, M.E., Abdel-Samiee, M., *et al.* (2020). Total tumor volume as a prognostic value for survival following liver resection in patients with hepatocellular carcinoma. Retrospective cohort study. *Ann. Med. Surg.* 54 : 47–53.

Zhang, J.W., Feng, X.Y., Liu, H.Q., Yao, Z.W., Yang, Y.M., Liu, B., *et al.* (2010). CT volume measurement for prognostic evaluation of unresectable hepatocellular carcinoma after TACE. *World J. Gastroenterol.* 16 : 2038–2045.