UNIVERSITAS
GADJAH MADA
Performa Microwave-Assisted Extraction untuk Ekstraksi Senyawa Fungsional pada Daun Kelor (Moringa oleifera Lam.) dengan Green Solvent
Ana Nur Fauziah, Dr. Ir. Devi Yuni Susanti, S.T.P., M. Sc., IPU, ASEAN Eng.; Bayu Nugraha, S.T.P., M.Sc., Ph.D.
Universitas Gadjah Mada, 2026 | Diunduh dari http://etd.repository.ugm.ac.id/

DAFTAR PUSTAKA

Adjali, A., Clarot, I., Chen, Z., Marchioni, E., & Boudier, A. (2022). Physicochemical degradation of phycocyanin and means to improve its stability: A short review. *Journal of Pharmaceutical Analysis*, 12(3), 406–414.

Alara, O. R., Abdurahman, N. H., & Olalere, O. A. (2018). Optimization of microwave-assisted extraction of flavonoids and antioxidants from Vernonia amygdalina leaf using response surface methodology. *Food and bioproducts processing*, *107*, 36-48.

Alara, O. R., Nour, A. H., & Mudalip, S. K. A. (2019). Screening of microwave-assisted-batch extraction parameters for recovering total phenolic and flavonoid contents from Chromolaena odorata leaves through two-level factorial design. *Indones J Chem*, *19*(2), 511-21.

Ali, A., Chong, C. H., Mah, S. H., Abdullah, L. C., Choong, T. S. Y., & Chua, B. L. (2018). Impact of storage conditions on the stability of predominant phenolic constituents and antioxidant activity of dried *Piper betle* extracts. *Molecules*, 23(2), 484.

Alshammari, O. A., Almulgabsagher, G. A., Ryder, K. S., & Abbott, A. P. (2021). Effect of solute polarity on extraction efficiency using deep eutectic solvents. Green Chemistry, 23(14), 5097-5105.

Aminah, S., Ramdhan, T., & Yanis, M. (2015). Kandungan nutrisi dan sifat fungsional tanaman kelor (*Moringa oleifera*). *Buletin Pertanian Perkotaan*, 5(2), 35–44.

Anwar, F., Naseer, R., Bhanger, M. I., & Ashraf, S. (2021). Citric acid as a natural antioxidant and pH regulator in food systems. *Food Chemistry*, 365, 130–141.

Apriantini, A., Putra, R. G., & Suryati, T. (2022). Aplikasi ekstrak daun kelor (*Moringa oleifera*) pada berbagai produk olahan daging. *Jurnal Ilmu Produksi dan Teknologi Hasil Peternakan*, 10(3), 132–143.

Arimboor, R., Natarajan, R. B., & Menon, K. R. (2021). Carotenoids and their health benefits. In R. Carle & R. Schweiggert (Eds.), *Handbook on Natural Pigments in Food and Beverages* (pp. 1–25). Woodhead Publishing.

Avilés-Gaxiola, S., León-Félix, J., Jiménez-Nevárez, Y. B., Angulo-Escalante, M. A., Ramos-Payán, R., Colado-Velázquez III, J., & Heredia, J. B. (2021). Antioxidant and anti-inflammatory properties of novel peptides from *Moringa oleifera* Lam. leaves. *South African Journal of Botany*, 141, 466–473.

Batool, S., Khan, S., & Basra, S. M. A. (2020). Foliar application of moringa leaf extract improves the growth of moringa seedlings in winter. *South African Journal of Botany*, 129, 347–353.

Bhadange, Y. A., Carpenter, J., & Saharan, V. K. (2024). A comprehensive review on advanced extraction techniques for retrieving bioactive components from natural sources. *ACS Omega*, 9(29), 31274–31297.

Cavalloro, V., Martino, E., Linciano, P., & Collina, S. (2021). Microwave-assisted solid extraction from natural matrices. Microwave heating-electromagnetic fields causing thermal and non-thermal effects, 35-56.

Cerro, D., González, C., Armijo, C., & Sepúlveda, L. (2026). Recovery of polyphenolic compounds from Brassica oleracea leaves biowaste by ultrasound-assisted and maceration extraction: Study of variables effect and kinetics using Peleg model. *Preparative Biochemistry & Biotechnology*, *56*(2), 274-286.

UNIVERSITAS GADJAH MADA
Performa Microwave-Assisted Extraction untuk Ekstraksi Senyawa Fungsional pada Daun Kelor (Moringa oleifera Lam.) dengan Green Solvent
Ana Nur Fauziah, Dr. Ir. Devi Yuni Susanti, S.T.P., M. Sc., IPU, ASEAN Eng.; Bayu Nugraha, S.T.P., M.Sc., Ph.D.
Universitas Gadjah Mada, 2026 | Diunduh dari http://etd.repository.ugm.ac.id/

Chaves, J. O., De Souza, M. C., Da Silva, L. C., Lachos-Perez, D., Torres-Mayanga, P. C., Machado, A. P. D. F., ... & Rostagno, M. A. (2020). Extraction of flavonoids from natural sources using modern techniques. *Frontiers in chemistry*, *8*, 507887.

Chemat, F., & Cravotto, G. (Eds.). (2012). *Microwave-assisted extraction for bioactive compounds: theory and practice* (Vol. 4). Springer Science & Business Media.

Cheng, H. P., Dai, J., Nemes, S., & Vijaya Raghavan, G. S. (2006). Comparison of conventional extraction under reflux conditions and microwave-assisted extraction of oil from popcorn. Journal of Microwave Power and Electromagnetic Energy, 41(1), 36-44.

Chuyen, H. V., Nguyen, M. H., Roach, P. D., Golding, J. B., & Parks, S. E. (2018). Microwave-assisted extraction and ultrasound-assisted extraction for recovering carotenoids from Gac peel and their effects on antioxidant capacity of the extracts. Food Science & Nutrition, 6(1), 189-196.

Coelho, J. P., Barreira, J. C. M., & Ferreira, I. C. F. R. (2021). Optimization of microwave-assisted extraction of phenolic compounds from food by-products using RSM. *Molecules*, 26(15), 4578.

Cueto Covarrubias, L. A., Valdez Solana, M. A., Avitia Domínguez, C., Téllez Valencia, A., Meza Velázquez, J. A., & Sierra Campos, E. (2023). Characterization of *Moringa oleifera* seed oil for the development of a biopackage applied to maintain the quality of turkey ham. *Polymers*, 16(1), 132.

Dahmoune, F., Nayak, B., Moussi, K., Remini, H., & Madani, K. (2015). Optimization of microwave-assisted extraction of polyphenols from Myrtus communis L. leaves. *Food chemistry*, *166*, 585-595.

Djiobie Tchienou, G. E., Tsatsop Tsague, R. K., Dongmo Sokeng, S., & Ngassoum, M. B. (2022). Kinetic study in the extraction of xanthones from Securidaca longepedunculata roots by microwave-assisted-extraction. Beni-Suef University Journal of Basic and Applied Sciences, 11(1), 116.

Falowo, A. B., Mukumbo, F. E., Idamokoro, E. M., Lorenzo, J. M., Afolayan, A. J., & Muchenje, V. (2018). Multi-functional application of *Moringa oleifera* Lam. in nutrition and animal food products: A review. *Food Research International*, 106, 317–334.

Fraterrigo Garofalo, S., Demichelis, F., Peletti, V., Picco, L., Tommasi, T., & Fino, D. (2025). Comparative study of polyphenol extraction using physical techniques and water as a solvent: A sustainable approach for the valorization of apple pomace. *Environmental Science and Pollution Research*, *32*(48), 27915-27928.

Gandji, K., Chadare, F. J., Idohou, R., Salako, V. K., Assogbadjo, A. E., & Glele Kakai, R. I. (2018). Status and utilisation of *Moringa oleifera* Lam: A review. *African Crop Science Journal*, 26(1), 137–156.

García-Beleño, J., & Rodríguez de San Miguel, E. (2021). Integration of response surface methodology (RSM) and principal component analysis (PCA) as an optimization tool for polymer inclusion membrane based-optodes designed for Hg (II), Cd (II), and Pb (II). *Membranes*, 11(4), 288.

García-Martín, J. F., et al. (2023). Microwave-assisted extraction of polyphenols from citrus by-products: process optimization and kinetic modeling. *Food Chemistry*, 404, 134586.

UNIVERSITAS
GADJAH MADA
Performa Microwave-Assisted Extraction untuk Ekstraksi Senyawa Fungsional pada Daun Kelor (Moringa oleifera Lam.) dengan Green Solvent
Ana Nur Fauziah, Dr. Ir. Devi Yuni Susanti, S.T.P., M. Sc., IPU, ASEAN Eng.; Bayu Nugraha, S.T.P., M.Sc., Ph.D.
Universitas Gadjah Mada, 2026 | Diunduh dari http://etd.repository.ugm.ac.id/

Granato, D., Shahidi, F., Wrolstad, R., Kilmartin, P., Melton, L. D., Hidalgo, F. J., ... & Finglas, P. (2018). Antioxidant activity, total phenolics and flavonoids contents: Should we ban in vitro screening methods?. *Food chemistry*, *264*, 471-475.

Hardiyanthi, F. (2015). *Pemanfaatan aktivitas antioksidan ekstrak daun kelor (Moringa oleifera) dalam sediaan hand and body cream*. Skripsi. UIN Syarif Hidayatullah Jakarta.

Hiranvarachat, B., Devahastin, S., Chiewchan, N., & Raghavan, G. V. (2013). Structural modification by different pretreatment methods to enhance microwave-assisted extraction of β-carotene from carrots. *Journal of Food Engineering*, *115*(2), 190-197.

Hosseini, S., Gharachorloo, M., Ghiassi-Tarzi, B., & Ghavami, M. (2016). Evaluation of the organic acids ability for extraction of anthocyanins and phenolic compounds from different sources and their degradation kinetics during cold storage. *Polish Journal of Food and Nutrition Sciences*, 66(4), 261–269.

Huang, Y., Yin, Z., Guo, J., Wang, F., & Zhang, J. (2019). Oil extraction and evaluation from yellow horn using a microwave-assisted aqueous saline process. *Molecules*, 24(14), 2598.

Ibrahim, N., Hashim, S. N., & Zawawi, R. (2019). Application of organic acids in green extraction of phenolic compounds from plant matrices. *Journal of Cleaner Production*, 239, 118–135.

Integrated Taxonomic Information System. (2013). *Moringa oleifera* (Drumstick Tree): Biological Classification and Name. *Encyclopedia of Life Newsletter.* Tanggal akses 6 November 2021. http://hy_entries/46214757/overview/moringa-oleifera

Jaeschke, D. P., Teixeira, I. R., Marczak, L. D. F., & Mercali, G. D. (2021). Phycocyanin from *Spirulina*: A review of extraction methods and stability. *Food Research International*, 143, 110314.

Jolliffe, I. T., & Cadima, J. (2016). Principal component analysis: A review and recent developments. *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences*, 374(2065), 20150202.

Karam, E. A., Abou-Arab, A. A., & Salama, M. F. (2020). Effect of citric acid-assisted extraction on phenolic and antioxidant content of plant materials. *Industrial Crops and Products*, 150, 112–119.

Kaur, P., Gupta, R. C., Dey, A., Malik, T., & Pandey, D. K. (2021). Optimization of harvest and extraction factors by full factorial design for the improved yield of C-glucosyl xanthone mangiferin from Swertia chirata. *Scientific Reports*, *11*(1), 16346

Khalishah, A. N., Mawarni, B., Hidayati, L., & Dhiya, S. S. (2023). Pemanfaatan daun kelor (*Moringa oleifera* L.) untuk mengatasi anemia. *Jurnal Pengemas Kesehatan*, 2(1), 18–25.

Kortei, N. K., Odamtten, G. T., Obodai, M., Appiah, V., & Akonor, P. T. (2015). Determination of color parameters of gamma irradiated fresh and dried mushrooms during storage. *Croatian Journal of Food Technology, Biotechnology and Nutrition*, 10(1–2), 66–71.

Koubaa, M., et al. (2020). Modelling of polyphenols extraction kinetics from food matrices: Comparison between Peleg, second-order and Fick's diffusion models. *Separation and Purification Technology*, 248, 117–118.

UNIVERSITAS GADJAH MADA
Performa Microwave-Assisted Extraction untuk Ekstraksi Senyawa Fungsional pada Daun Kelor (Moringa oleifera Lam.) dengan Green Solvent
Ana Nur Fauziah, Dr. Ir. Devi Yuni Susanti, S.T.P., M. Sc., IPU, ASEAN Eng.; Bayu Nugraha, S.T.P., M.Sc., Ph.D.
Universitas Gadjah Mada, 2026 | Diunduh dari http://etd.repository.ugm.ac.id/

Kumar, V., Sharma, R., & Singh, S. (2020). Citric acid: Properties and applications in food, pharmaceuticals and bioprocessing industries. *Biotechnology Reports*, 27, e00532.

Kuna-Broniowska, I., Blicharz-Kania, A., Andrejko, D., Kubik-Komar, A., Kobus, Z., Pecyna, A., ... & Rydzak, L. (2019). Modelling water absorption in micronized lentil seeds with the use of Peleg's Equation. *Sustainability*, 12(1), 261.

Lim, Y. Y., & Murtijaya, J. (2007). Antioxidant properties of Phyllanthus amarus extracts as affected by different drying methods. *LWT-Food Science and Technology*, *40*(9), 1664-1669.

López-Salazar, H., Camacho-Díaz, B. H., Ocampo, M. A., & Jiménez-Aparicio, A. R. (2023). Microwave-assisted extraction of functional compounds from plants: A Review. BioResources, 18(3), 6614.

Maity, T., Bawa, A. S., & Raju, P. S. (2018). Effect of preconditioning on physicochemical, microstructural, and sensory quality of vacuum-fried jackfruit chips. *Drying Technology*, 36(1), 63–71.

Mandal, V., Mohan, Y., & Hemalatha, S. J. P. R. (2007). Microwave assisted extraction—an innovative and promising extraction tool for medicinal plant research. *Pharmacognosy reviews*, *1*(1), 7-18.

Melo, N. V., Vargas, T. Q., & Calvo, C. M. C. (2013). *Moringa oleifera* L.: An underutilized tree with macronutrients for human health. *Emirates Journal of Food and Agriculture*, 25(10), 785–789.

Muteeb, G., Aatif, M., Farhan, M., Alsultan, A., Alshoaibi, A., & Alam, M. W. (2023). Leaves of *Moringa oleifera* are potential source of bioactive compound β-carotene: Evidence from in silico and quantitative gene expression analysis. *Molecules*, 28(4), 1578.

Nebolisa, N. M., Umeyor, C. E., Ekpunobi, U. E., Umeyor, I. C., & Okoye, F. B. (2023). Profiling the effects of microwave-assisted and Soxhlet extraction techniques on the physicochemical attributes of *Moringa oleifera* seed oil and proteins. *Oil Crop Science*, 8(1), 16–26.

Ngcobo, N., Manzi, M., Nwaila, G., Bourdeau, J., & Zhang, S. (2024, September). Comparing Modern Deep Learning with Traditional Methods for Seismic Fault Interpretation. In *NSG 2024 5th Conference on Geophysics for Mineral Exploration and Mining* (Vol. 2024, No. 1, pp. 1-5). European Association of Geoscientists & Engineers.

Nimse, S. B., Pal, D., & Sonawane, S. K. (2023). Green extraction of carotenoids: Recent trends and applications. *Trends in Food Science & Technology*, 139, 50–65.

Nobosse, P., Fombang, E. N., & Mbofung, C. M. (2018). Effects of age and extraction solvent on phytochemical content and antioxidant activity of fresh Moringa oleifera L. leaves. BMC Research Notes / Journal of Food Science and Technology.

Oktaviani, D. J., Widiyastuti, S., Maharani, D. A., Amalia, A. N., Ishak, A. M., & Zuhrotun, A. (2019). Review: Bahan alami penyembuh luka. *Farmasetika.com*, 4(3), 44.

Oyeyinka, A. T., & Oyeyinka, S. A. (2018). *Moringa oleifera* as a food fortificant: Recent trends and prospects. *Journal of the Saudi Society of Agricultural Sciences*, 17(2), 127–136.

UNIVERSITAS GADJAH MADA
Performa Microwave-Assisted Extraction untuk Ekstraksi Senyawa Fungsional pada Daun Kelor (Moringa oleifera Lam.) dengan Green Solvent
Ana Nur Fauziah, Dr. Ir. Devi Yuni Susanti, S.T.P., M. Sc., IPU, ASEAN Eng.; Bayu Nugraha, S.T.P., M.Sc., Ph.D.
Universitas Gadjah Mada, 2026 | Diunduh dari http://etd.repository.ugm.ac.id/

Palos-Hernandez, A., Fernández, M. Y. G., Burrieza, J. E., Pérez-Iglesias, J. L., & González-Paramás, A. M. (2022). Obtaining green extracts rich in phenolic compounds from underexploited food by-products using natural deep eutectic solvents. *Sustainable Chemistry and Pharmacy*, 29, 100773.

Patil, S. S., Deshannavar, U. B., Ramasamy, M., Emani, S., Khalilpoor, N., & Issakhov, A. (2021). Study of extraction kinetics of Total polyphenols from curry leaves. International journal of chemical engineering, 2021(1), 9988684.

Patil, D., Vaknin, Y., Rytwo, G., Lakemond, C., & Benjamin, O. (2022). Characterization of *Moringa oleifera* leaf and seed protein extract functionality in emulsion model system. *Innovative Food Science & Emerging Technologies*, 75, 102903.

Pitarokoili, K., et al. (2022). Optimization of microwave-assisted extraction of carotenoids from tomato waste using response surface methodology. *LWT–Food Science and Technology*, 155, 112968.

Putri, W. D. R., Riyanto, E. I., Heliana, A., Wellang, A., & Zubaidah, E. (2021). Optimization of microwave-assisted extraction (mae) time and material to solvent ratio of gembili (Dioscorea esculenta) water-soluble polysaccharides (WSP). Industria: Jurnal Teknologi dan Manajemen Agroindustri, 10(1), 78-89.

Rodríguez-Pérez, C., Quirantes-Piné, R., Fernández-Gutiérrez, A., & Segura-Carretero, A. (2015). Optimization of extraction method to obtain a phenolic compounds-rich extract from Moringa oleifera Lam leaves. *Industrial Crops and Products*, *66*, 246-254.

Rodríguez-Pérez, C., Gilbert-López, B., Mendiola, J. A., Quirantes-Piné, R., Segura-Carretero, A., & Ibáñez, E. (2016). Optimization of microwave-assisted extraction and pressurized liquid extraction of phenolic compounds from Moringa oleifera leaves by multiresponse surface methodology. *Electrophoresis*, *37*(13), 1938-1946.

Rodríguez-Concepción, M., et al. (2018). A global perspective on carotenoids: metabolism, biotechnology, and benefits for nutrition and health. *Progress in Lipid Research*, 70, 62–93.

Routray, W., & Orsat, V. (2012). Microwave-assisted extraction of flavonoids: a review. *Food and Bioprocess Technology*, *5*(2), 409-424.

Saini, R. K., Sivanesan, I., & Keum, Y. S. (2016). Phytochemicals of *Moringa oleifera*: A review of their nutritional, therapeutic and industrial significance. *3 Biotech*, 6, 203. https://doi.org/10.1007/s13205-016-0526-3

Saini, R. K., & Keum, Y. S. (2018). Carotenoid extraction methods: A review of recent developments. *Food chemistry*, *240*, 90-103.

Satriyani, D. P. P. (2021). Review artikel: Aktivitas antioksidan ekstrak daun kelor (*Moringa oleifera* Lam.). *Jurnal Farmasi Malahayati*, 4(1), 31–43.

Shiriki, D., Igyor, M. A., & Gernah, D. I. (2015). Nutritional evaluation of complementary food formulations from maize, soybean and peanut fortified with *Moringa oleifera* leaf powder. *Food and Nutrition Sciences*, 6, 494–500.

Souza, D. E. S., Melo, J. J. C. D., Santos, F. F. D., Vasconcelos, A. L. D. S., Jesus, A. D. S. D., Freitas, L. D. S., ... & Soares, C. M. F. (2024). Microwave-assisted vs. conventional extraction of *Moringa oleifera* seed oil: Process optimization and efficiency comparison. *Foods*, 13(19), 3141.

UNIVERSITAS
GADJAH MADA
Performa Microwave-Assisted Extraction untuk Ekstraksi Senyawa Fungsional pada Daun Kelor (Moringa oleifera Lam.) dengan Green Solvent
Ana Nur Fauziah, Dr. Ir. Devi Yuni Susanti, S.T.P., M. Sc., IPU, ASEAN Eng.; Bayu Nugraha, S.T.P., M.Sc., Ph.D.
Universitas Gadjah Mada, 2026 | Diunduh dari http://etd.repository.ugm.ac.id/

Srivastava, S., Pandey, V. K., Dash, K. K., Dayal, D., Wal, P., Debnath, B., ... & Dar, A. H. (2023). Dynamic bioactive properties of nutritional superfood *Moringa oleifera*: A comprehensive review. *Journal of Agriculture and Food Research*, 14, 100860.

Stahl, W., & Sies, H. (2020). Antioxidant activity of carotenoids. *Molecular Aspects of Medicine*, 70, 100–115.

Susanti, D. Y., Sediawan, W. B., Fahrurrozi, M., & Hidayat, M. (2021). Foam-mat drying in the encapsulation of red sorghum extract: Effects of xanthan gum addition on foam properties and drying kinetics. *Journal of the Saudi Society of Agricultural Sciences*, 20(4), 270–279.

Susanti, D. Y., Kalyani, N. F., Astuti, F. P., Purwandari, F. A., Kurniawan, M. P., Kistanti, A., ... & Setyaningsih, W. (2025). Characterization and Performance Evaluation of NADES-CAF in Ultrasound and Microwave Extraction of Phycocyanin from Arthrospira platensis. International Journal of Design & Nature and Ecodynamics, 20(1), 21-30.

Teixeira, E. M., Carvalho, M. R., Neves, V. A., Silva, M. A., & Arantes-Pereira, L. (2014). Chemical characteristics and fractionation of proteins from *Moringa oleifera* Lam. leaves. *Food Chemistry*, 147, 51–54.

Torun, M., Dincer, C., Topuz, A., Sahin–Nadeem, H., & Ozdemir, F. (2015). Aqueous extraction kinetics of soluble solids, phenolics and flavonoids from sage (Salvia fruticosa Miller) leaves. *Journal of Food Science and Technology*, *52*(5), 2797-2805.

Toscano, S., Gómez-Bellot, M. J., Romano, D., & Sánchez-Blanco, M. J. (2023). Physiological and biochemical changes in response to *Moringa oleifera* biostimulant in petunia plants under water deficit. *Scientia Horticulturae*, 319, 112187.

Villegas-Vazquez, E. Y., Gómez-Cansino, R., Marcelino-Pérez, G., Jiménez-López, D., & Quintas-Granados, L. I. (2025). Unveiling the miracle tree: Therapeutic potential of *Moringa oleifera* in chronic disease management and beyond. *Biomedicines*, 13(3), 634.

Vongsak, B., Sithisarn, P., & Gritsanapan, W. (2013). Bioactive content and antioxidant activity of *Moringa oleifera* leaf extracts. *Industrial Crops and Products*, 49, 723–728.

Waterman, C., Rojas-Silva, P., Tumer, T. B., Kuhn, P., Richard, A. J., & Raskin, I. (2014). Isothiocyanate-rich *Moringa oleifera* extract reduces weight gain, insulin resistance, and hepatic gluconeogenesis in mice. *Molecular Nutrition & Food Research*, 58(8), 1615–1627.

Zhang, Q., Wang, H., & Li, X. (2021). Microwave-assisted extraction of phenolic compounds using citric acid-based green solvents. *Separation and Purification Technology*, 276, 119–218.

Zhi Ling, M., & Chang Bin, W. (2018, June). The effect of different bags on chlorophyll and carotenoids in the peel of Chinhuang mango. In *E3S Web of Conferences* (Vol. 38, p. 02026). EDP Sciences.