

## DAFTAR PUSTAKA

- Alfarra, H. Y., & Omar, M. N., 2013. *Centella asiatica* : from folk remedy to the medicinal biotechnology - a state revision. *Int Journal Biosci*; 6655: 49–67.
- Benzing, T. 2004. Signaling at the Slit Diaphragm. *J Am Soc Nephro*; 34: 1382–1391.
- Bidani, Anil K., Polichnowski, Aaron J, Loutzenhiser, R, and Griffin, Karen A., 2013. Renal microvascular dysfunction, hypertension and CKD progression. *Curr Opin Nephrol Hypertens*: 1–9.
- Breyer, M D., and Susztak, K., 2016. The next generation of therapeutics for chronic kidney disease. *Nat Rev Drug Discov*: 568-588.
- Budiman, L., 2014. Inhibisi ekstrak pegagan (*Centella asiatica* L.) dan tempuyung (*Sonchus arvensis* L.) terhadap aktivitas ACE (Angiotensin Converting Enzyme) [skripsi]. Bogor (ID): Institut Pertanian Bogor.
- Cahyawati, P. N., Arfian, N., Ngatidjan, Sari, D. C. R., Romi, M. M., 2017. Simvastatin attenuates renal failure in mice with A 5/6 subtotal nephrectomy. *Int J Pharm Sci*; 9(5):12-17.
- Drawz, P.E. and Rosenberg, M.E., 2013. Slowing progression of chronic kidney disease. *Kidney Int Sup*; 3(4): p.372.
- Fogo, A B., 2000. Glomerular hypertension, abnormal glomerular growth, and progression of renal diseases. *Kidney Int*: S15-S21.
- Fogo, A. B., 2003. Animal Models of FSGS : Lessons for Pathogenesis and Treatment. *Semin Nephrol*; 23(2): 161–171.
- Fogo, A. B., 2007. Mechanisms of progression of chronic kidney disease. *Pediatr Nephrol*: 22(12): 2011–2022.
- Gava A. L., Freitas F. P., Balarini C. M., Vasquez E. C., Meyrelles S. S., 2012. Effects of 5/6 nephrectomy on renal function and blood pressure in mice. *Int J Physiol Pathophysiol Pharmacol*: 4(3): 167-173.
- Gribouval, O., Roselli, S., Boute, N., Sich, M., Benessy, F., Attié, T., Antignac, C., 2002. Podocin Localizes in the Kidney to the Slit Diaphragm Area. *Am J Pathol*, 160(1), 131–139.
- Hill, N. R., Fatoba, S. T., Oke, J. L., Hirst, J. A., O’Callaghan, C. A., Lasserson, D. S., & Hobbs, F. D. R., 2016. Global Prevalence of Chronic Kidney Disease – A Systematic Review and Meta-Analysis. *PLoS ONE*; 11(7).
- Ikezumi, Y., Suzuki, T., Karasawa, T., Kawachi, H., Nikolic-Paterson, D J., Uchiyama, M., 2008. Activated macrophages down-regulate podocyte nephryn and podocin expression via stress-activated protein kinases. *Biochem Biophys Res Commun*: 706–711.
- Jefferson, J A., and Shankland, S J., 2014. The Pathogenesis of Focal Segmental Glomerulosclerosis. *Adv Chronic Kidney Dis*: 408–416.
- Just, A., 2006. Mechanisms of renal blood flow autoregulation: dynamics and contributions. *Am J Physiol Regul Integr Comp Physiol*: R1-R17.
- Kren S., Hostetter T.H., 1999. The course of the remnant kidney model in mice. *Kidney Int*; 56: 333–337.

- Kriz, W., Hackenthal, E., Nobiling R., Sakai, T., Elger, M., and Hahnel, B., 1994. A role for podocytes to counteract capillary wall distension. *Kidney Int*: 369–376.
- Kriz, W., and Lemley, K.V., 2015. A potential role formechanical forces in the detachment of podocytes and the progression of CKD, *J Am Soc Neph* : 258–269.
- Kumar, V., Abbas, A.K., Fausto, N. and Aster, J.C., 2014. *Robbins and Cotran Pathologic Basis of Disease, Professional Edition E-Book*. Elsevier Health Sciences.
- Ljungberg, I. V. I. L., Artiovaara, J. O. W., Enkkeri, U. L. L. A. L., & Estila, M. A. K., 1999. Neph rin is specifically located at the slit diaphragm of glomerular podocytes. *Proc Natl Acad Sci*; 96(July): 7962–7967.
- Mao, J., Zhang, Y., Du, L., Dai, Y., Yang, C., Liang, L., 2006. Expression profile of neph rin, podocin, and CD2AP in Chinese children with MCNS and IgA nephropathy. *Pediatr Nephrol*; 21: 1666-1675.
- Matovinovic, M. S., 2009. Pathophysiology and classification of kidney. *eJIFCC*; 20(1): 1–10.
- Matsubara T<sup>1</sup>, Abe H, Arai H, Nagai K, Mima A, Kanamori H, Sumi E, Takahashi T, Matsuura M, Iehara N, Fukatsu A, Kita T, Doi T., 2006. Expression of Smad1 is directly associated with mesangial matrix expansion in rat diabetic nephropathy. *Lab Invest*;86(4):357-68
- Mescher, A. L., & Junqueira, L. C. U., 2013. Junqueira's basic histology: Text and atlas (Thirteenth edition.). New York: *McGraw Hill Medical*.
- Menne, J., Meier, M., Park, J K., Boehne, M., Kirsch, T., Lindschau, C., *et al.*, 2006. Neph rin loss in experimental diabetic nephropathy is prevented by deletion of protein kinase C alpha signaling *in-vivo*. *Kidney Int*: 1456-1462.
- Moore, K L., Arthur, F. Dalley, II, and Agur, M R A., 2014. Clinically Oriented Anatomy 7th Edition. Philadelphia: *Lippincott Williams & Wilkins*.
- Miesen, L., Steenbergen, E., and Smeets, B., 2017. Parietal cells—new perspectives in glomerular disease. *Cell Tissue Res*: 237–244.
- Miner, J H., 1999. Neph rin is specifically located at the slit diaphragm of glomerular podocytes. *Proc Natl Acad Sci*: 7962-7967
- Mitch, W E., Walser, M., Buffington, G A., Lemann, J Jr., 1976. A simple method of stimating progression of chronic renal failure. *Lancet*. 2:1326–1328.
- Mundel, Peter and Shankland, Stuart J., 2002. Podocyte Biology and Response to Injury. *J Am Soc Nephrol*. 13: 3005–3015
- Nagata, Michio., 2016. Podocyte Injury and Its Consequences. *Kidey Int*. 89: 1221–1230.
- Ojo, A., 2014. Addressing the Global Burden of Chronic Kidney Disease Through Clinical and Translational Research. *Trans Am Clin Climatol Assoc*; 125: 229–246.
- Leelahavanichkul, A., Yan, Q., Hu, X., Eisner, C., Huang, Y., Chen, R., 2010. Rapid CKD progression in a new mouse kidney remnant model: strain-dependent resistance is overcome by angiotensin II. *Kidney Int*, 78(11), 1136–1153.

- Persatuan Nefrologi Indonesia. 2014. 7th Report Of Indonesian Renal Registry .
- Rao, K. G. M., Rao, S. M., & Rao, S. G., 2006. *Centella asiatica* ( L .) Leaf Extract Treatment During the Growth Spurt Period Enhances Hippocampal CA3 Neuronal Dendritic Arborization in Rats, *Evid Based Complement Alternat Med.* 3(June), 349–357.
- Schlondorff, D.O., 2008. Overview of factors contributing to the pathophysiology of progressive renal disease. *Kidney Int*, 74(7); 860-866.
- Song J, Xu H, Lu Q, Xu Z, Bian D, Xia Y, Wei Z, Gong Z, Dai Y., 2012. Madecassoside suppresses migration of fibroblasts from keloids: involvement of p38 kinase and PI3K signaling pathways. *J Burns* 38, 677-684.
- Tang, L., He, R., Yang, G., Tan, J., Zhou, L., Meng, X., & Huang, X. R., 2012. Asiatic Acid Inhibits Liver Fibrosis by Blocking TGF-beta / Smad Signaling In Vivo and In Vitro. *Plos ONE*: 7(2).
- Van de Lest, N A., Zandbergen, M., Ijpelaar, D H T., Wolterbeek, R., Brujin, J A., Bajema, I M., Scharpfenecker, M., 2017. Neph rin Loss Can Be Used to Predict Remission and Long-Term Renal Outcome in Patients With Minimal Change Disease. *Kidney Int Rep*:168 – 177.
- Xu MF, Xiong YY, Liu JK, Qian JJ, Zhu L, Gao J., 2012. Asiatic acid, a pentacyclic triterpene in *Centella asiatica*, attenuates glutamate-induced cognitive deficits in mice and apoptosis in SH-SY5Y cells. *Acta Pharmacol Sin* 33(5), 578-587
- Wang, Zhu, Juntian Liu, Wansen Sun., 2013. Effects of asiaticoside on levels of podocyte cytoskeletal proteins and renal slit diaphragm proteins in adriamycin-induced rat nephropathy. (93)8: 352-358,
- Wiener, C., Fauci, A., Braunwald, E., Kasper, D., Hauser, S., Longo, D., Jameson, J., Loscalzo, J. and Brown., 2012. Harrison's Principles of Internal Medicine 18th Edition. New York: *The McGraw Hill*, 2308-2309.
- Wühl, E., & Schaefer, F., 2008. Therapeutic strategies to slow chronic kidney disease progression, 705–716.
- Yang, H. C., Zuo, Y., & Fogo, A. B., 2010. Models of chronic kidney disease. *Drug Discovery Today: Disease Models.* 7(1): 13-19.
- Yuan, H., Takeuchi, E., and Salant, D J., 2002. Podocyte slit-diaphragm protein neph rin is linked to the actin cytoskeleton. *Am J Physiol Renal Physiol*: F585–F591.
- Zheng J, Zhang L, Wu M, Li X, Wan J., 2010. Protective effects of asiaticoside on sepsis-induced acute kidney injury in mice. *Zhongguo Zhong Yao Za Zhi*: 1482-1485.