

DAFTAR PUSTAKA

- Adame, M. F., Connolly, R. M., Turschwell, M. P., Lovelock, C. E., Fatoyinbo, T., Lagomasino, D., ... & Brown, C. J. (2021). Future carbon emissions from global mangrove forest loss. *Global change biology*, 27(12), 2856-2866.
- Adidharma, G. B. (2025). *Perbandingan model regresi linear dan machine learning untuk estimasi stok karbon atas permukaan mangrove di Teluk Pangpang, Kabupaten Banyuwangi menggunakan citra Sentinel-2A MSI*. Skripsi. Program Studi Kartografi dan Penginderaan Jauh, Universitas Gadjah Mada, Yogyakarta.
- Akbar, C., Arsepta, Y., Dewiyanti, I., & Bahri, S. (2019). Dugaan serapan karbon pada vegetasi mangrove, di kawasan mangrove Desa Beureunut, Kecamatan Seulimum, Kabupaten Aceh Besar. *Jurnal Laot Ilmu Kelautan*, 1(2), 63-70.
- Al Latifa, R., Sari, K. E., & Meidiana, C. (2022). Faktor rumah tangga yang mempengaruhi emisi CO₂ di Kelurahan Jodipan, Kota Malang. *Planning for Urban Region and Environment Journal (PURE)*, 11(3), 89-100.
- Alongi, D. M. (2008). Mangrove forests: resilience, protection from tsunamis, and responses to global climate change. *Estuarine, coastal and shelf science*, 76(1), 1-13.
- Althouse, L.A., Ware, W.B. & Ferron, J.M. (1998). Detecting departures from normality: A monte carlo simulation of a new omnibus test based on moments. *Paper presented at the Annual Meeting of the American Educational Research Association*, San Diego, CA.
- Astaman, I. D. M. K. P., Karang, I. W. G. A., & Gede, I. (2021). Pemetaan habitat dasar perairan dangkal menggunakan Citra Satelit SPOT-7 di Pulau Nusa Lembongan, Bali. *Journal of Marine and Aquatic Sciences*, 7(2), 184-195.
- Baloloy, A. B., Blanco, A. C., Candido, C. G., Argamosa, R. J. L., Dumalag, J. B. L. C., Dimapilis, L. L. C., & Paringit, E. C. (2018). Estimation of mangrove forest aboveground biomass using multispectral bands, vegetation indices and biophysical variables derived from optical satellite imageries: Rapideye, planetscope and sentinel-2. *ISPRS annals of the photogrammetry, remote sensing and spatial information sciences*, 4, 29-36.

- Bengen, D.G. (2002). *Pengenalan dan pengelolaan ekosistem mangrove*. Pusat Kajian Sumber Daya Pesisir dan Kelautan Institut Pertanian Bogor. Bogor.
- Bengen, D. G., Yonvitner, & Rahman. (2022). *Pedoman teknis pengenalan dan pengelolaan ekosistem mangrove*. IPB Press.
- Bouillon, S. (2011). Storage beneath mangroves. *Nature Geoscience*, 4(5), 282-283.
- Breiman, L. (2001). Random forests. *Machine learning*, 45(1), 5-32.
- Cahyo, W. E. (2010). Pengaruh pemanasan global terhadap lingkungan bumi. *Berita Dirgantara*, 8(2).
- Campbell, N. A., Reece, J. B., & Mitchell, L. G. (2005). *Biology* (7th ed.). San Francisco, CA: Pearson Benjamin Cummings.
- Candiago, S., Remondino, F., De Giglio, M., Dubbini, M., & Gattelli, M. (2015). Evaluating multispectral images and vegetation indices for precision farming applications from UAV images. *Remote sensing*, 7(4), 4026-4047.
- Candra, E. D. (2016). *Estimasi stok karbon hutan mangrove menggunakan Citra WorldView-2 di Teluk Benoa Bali*. Skripsi. Fakultas Geografi, UGM Yogyakarta.
- Chai, T., & Draxler, R. R. (2014). Root mean square error (RMSE) or mean absolute error (MAE) Arguments against avoiding RMSE in the literature. *Geoscientific model development*, 7(3), 1247-1250.
- Chave, J., Réjou-Méchain, M., Búrquez, A., Chidumayo, E., Colgan, M. S., Delitti, W. B., ... & Vieilledent, G. (2014). Improved allometric models to estimate the aboveground biomass of tropical trees. *Global change biology*, 20(10), 3177-3190.
- Ciais, P., Sabine, C., Bala, G., Bopp, L., Brovkin, V., Canadell, J., Chhabra, A., DeFries, R., Galloway, J., Heimann, M., Jones, C., Le Quéré, C., Myneni, R. B., Piao, S., & Thornton, P. (2013). Carbon and other biogeochemical cycles. In T. F. Stocker, D. Qin, G.-K. Plattner, M. Tignor, S. K. Allen, J. Boschung, A. Nauels, Y. Xia, V. Bex, & P. M. Midgley (Eds.), *Climate change 2013: The physical science basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change* (pp. 465–570). Cambridge University Press.

- Clough, B.F., Scott, K. (1989). Allometric relationships for estimating above ground biomass in six mangrove species. *Forest Ecol. Manage.* 27, 117-127.
- Comley, B.W.T., & McGuinness, K.A. (2005). Above- and below-ground biomass, and allometry of four common northern Australian mangroves. *Aust. J. Bot.* 53, 431–436.
- Danoedoro, P. (1996). *Pengolahan citra digital, teori dan aplikasinya dalam penginderaan jauh*. Fakultas Geografi Universitas Gadjah Mada, Yogyakarta.
- Danoedoro, P. (2009). Penginderaan jauh untuk inventarisasi mangrove: Potensi, keterbatasan dan kebutuhan data. *Proceedings Sinergi Survei dan Pemetaan Nasional dalam Mendukung Pengelolaan Mangrove Berkelanjutan*, 98-113.
- Donato, D. C., Kauffman, J. B., Murdiyarso, D., Kurnianto, S., Stidham, M., & Kanninen, M. (2011). Mangroves among the most carbon-rich forests in the tropics. *Nature geoscience*, 4(5), 293-297.
- English, S., Wilkinson, C., & Baker, V. (1997). *Survey Manual for Tropical Marine Resources* (2nd ed.). Townsville: Australian Institute of Marine Science.
- ESA. (2015). *ESA's Optical High-Resolution Mission for GMES Operational Services*. Paris, France: ESA.
- Falkowski, P. G., Scholes, R. J., Boyle, E., Canadell, J., Canfield, D., Elser, J., Gruber, N., Hibbard, K., Höglberg, P., Linder, S., Mackenzie, F. T., Moore III, B., Pedersen, T., Rosenthal, Y., Seitzinger, S., Smetacek, V., & Steffen, W. (2000). The global carbon cycle: A test of our knowledge of Earth as a system. *Science*, 290(5490), 291–296.
- Fariz, T. R., Ihsan, H. M., Lutfiananda, F., Sartohadi, J., Darmajati, Y., & Syahputra, A. (2023). Perbandingan pengukuran kerapatan kanopi dari hemispherical photography dan UAV untuk pemetaan menggunakan citra sentinel-2. *Jurnal Hutan Tropis Volume*, 11(1), 123-132.
- Firdaus, M. R., & Wijayanti, L. A. S. (2019). Fitoplankton dan siklus karbon global. *Oseana*, 44(2), 35-48.
- Frampton, W. J., Dash, J., Watmough, G., & Milton, E. J. (2013). Evaluating the capabilities of Sentinel-2 for quantitative estimation of biophysical variables

- in vegetation. *ISPRS journal of photogrammetry and remote sensing*, 82, 83-92.
- Fromard, F., Puig, H., Mougin, E., Marty, G., Betoulle, J. L., dan Cadamuro, L. (1998). Structure, above-ground biomass and dynamics of mangrove ecosystems: new data from French Guiana. *Oecologia*, 115(1-2), 39-53.
- Forestriko, H. F. (2016). Pemanfaatan Citra Landsat 8 untuk estimasi stok karbon hutan mangrove di Kawasan Segara Anakan Cilacap Jawa Tengah. *Jurnal Bumi Indonesia*, 5(1).
- Gitelson, A. A., Kaufman, Y. J., & Merzlyak, M. N. (1996). Use of a green channel in remote sensing of global vegetation from EOS-MODIS. *Remote Sensing of Environment*, 58(3), 289-298.
- Hadi, A. R. (2018). *Pendugaan cadangan karbon berdasarkan indeks vegetasi* (Doctoral dissertation, Universitas Brawijaya).
- Hadidi, A. F., Nana, K. T. M., & Boedijantoro, P. M. H. (2022). Estimasi stok karbon mangrove strata pohon di Kelurahan Trimulyo Kota Semarang sebagai upaya konservasi mangrove. *BIOMA*, 18(1).
- Hairiah, K., & Rahayu, S. (2007). Pengukuran karbon tersimpan di berbagai macam penggunaan lahan. *World agroforestry centre*. Bogor, 77.
- Hakim, M. A., Kamal, M., & Arjasakusuma, S. (2022). Mapping mangrove surface carbon stocks using multisensor imagery in Clungup Mangrove Conservation (CMC) Malang Regency. *Jurnal Geografi*, (2).
- Heriyanto, N.M., & Subiandono, E. (2016). Peran biomasa mangrove dalam menyimpan karbon di Kubu Raya, Kalimantan Barat. *Jurnal Analisis Kebijakan*, 13(1): 1-12
- Hermon, D., Ganefri, P. A., & Oktorie, O. (2018). The model of mangrove land cover change for the estimation of blue carbon stock change in Belitung Island-Indonesia. *International Journal of Applied Environmental Sciences*, 13(2), 191-202.
- Hidayat, Y. N., & Khakhim, N. (2017). Pemantauan distribusi muatan padatan tersuspensi menggunakan Citra Landsat 8 Oli di Muara Ci Tarum, Jawa Barat. *Jurnal Bumi Indonesia*, 6(1).

- Hidayatullah, M. F., Kamal, M., & Wicaksono, P. (2023). Species-based aboveground mangrove carbon stock estimation using WorldView-2 image data. *Remote Sensing Applications: Society and Environment*, 30, 100959.
- Howard, J., Hoyt, S., Isensee, K., Pidgeon, E., & Telszewski, M. (2014). *Coastal Blue Carbon: Methods for assessing carbon stocks and emissions factors in mangroves, tidal salt marshes, and seagrass meadows*.
- Huang, S., Tang, L., Hupy, J. P., Wang, Y., & Shao, G. (2021). A commentary review on the use of normalized difference vegetation index (NDVI) in the era of popular remote sensing. *Journal of forestry research*, 32(1), 1-6.
- Huete, A., Didan, K., Miura, T., Rodriguez, E. P., Gao, X., & Ferreira, L. G. (2002). Overview of the radiometric and biophysical performance of the MODIS vegetation indices. *Remote Sensing of Environment*, 83(1–2)
- Huete, A., & Glenn, E.P. (2011). *Remote sensing of ecosystem structure and function, advance in environment remote sensing*. Boca Raton: CRC Press.
- Huete, A. R., Liu, H. Q., Batchily, K. V., & Van Leeuwen, W. J. D. A. (1997). A comparison of vegetation indices over a global set of TM images for EOS-MODIS. *Remote sensing of environment*, 59(3), 440-451.
- Hutami, W. F. (2024). Populasi dan sampel dalam penelitian. *Jurnal Public Relations Mercu Buana, Query date*, 21, 23.
- Imbert, D., & Rollet, B. (1989). Phytomassierenne et production primaire dans la mangrove du Grand Cul-de-sac Marine (Guadeloupe, Antilles francaises). *Bull. Ecol.* 20, 27–39.
- Jensen, J. R. (2007). *Remote sensing of the environment: an earth resource perspective*. 2nd Edition. New Jersey: Pearson Prentice Hall.
- Jiang, Z., Huete, A. R., Didan, K., & Miura, T. (2008). Development of a two-band enhanced vegetation index without a blue band. *Remote sensing of Environment*, 112(10), 3833-3845.
- Jonathan Sarwono. (2012). *Metode penelitian kuantitatif dan kualitatif*. Yogyakarta: Graha Ilmu
- Jordan, C. F. (1969). Derivation of leaf-area index from quality of light on the forest floor. *Ecology*, 50(4), 663–666.

- Justice, C. O., & Townshend, J. R. G. (1981). Integrating ground data with remote sensing. In J. R. G. Townshend (Ed.), *Terrain Analysis and Remote Sensing* (pp. 38–58). London: Allen & Unwin.
- Kangkuso, A., Sharma, S., Jamili, J., Septiana, A., Sahidin, I., Rianse, U., & Nadaoka, K. (2018). Trends in allometric models and aboveground biomass of family Rhizophoraceae mangroves in the Coral Triangle ecoregion, Southeast Sulawesi, Indonesia. *Journal of Sustainable Forestry*, 37(7), 691–711.
- Karnieli, A., Agam, N., Pinker, R.T., Anderson, M., Imhoff, M. L., Gutman, G.G., Panov, N., & Goldberg, A. 2010. Use of NDVI and land surface temperature for drought assessment: merits and limitations. *Journal of Climate*, 23(3): 618-633.
- Kathiresan, K., & Bingham, B. L. (2001). Biology of mangroves and mangrove ecosystems.
- Kauffman, J. B., & Donato, D. C. (2012). *Protocols for the measurement, monitoring and reporting of structure, biomass and carbon stocks in mangrove forests* (Vol. 86, p. 7). Bogor, Indonesia: Cifor.
- Khairuddin, M. Y., & Syukur, A. (2018). Analisis kandungan logam berat pada tumbuhan mangrove. *Jurnal Biologi Tropis*, 18(1), 69-79.
- Kishwan, J., Pandey, R., & Dadhwal, V. K. (2009). India's forest and tree cover.
- Komiyama, A., Ong, J. E., & Pongparn, S. (2008). Allometry, biomass, and productivity of mangrove forests: A review. *Aquatic botany*, 89(2), 128-137.
- Komiyama A, Pongparn S, Kato S. (2005). Common allometric equations for estimating the tree weight of mangroves. *Journal of Tropical Ecology*. 21:471—47.
- Kuhn, M., & Johnson, K. (2013). *Applied predictive modeling* (Vol. 26, p. 13). Springer.
- Lillesand. T.M., W. Kiefer., Chipman, J.W. (2004). *Remote sensing and image interpretation (fifth edition)*. John Wiley & Sons, Inc., New York.
- Mahmudi, I. (2017). *Komparasi indeks vegetasi untuk estimasi stok karbon hutan mangrove di pesisir Muara Kali Porong, Sidoarjo*. Dalam *Seminar Nasional Geografi Universitas Negeri Malang* (hlm. 69). Universitas Negeri Malang.

- Mahyatar, P., & Kamal, M. (2021). Aboveground mangrove carbon stock mapping using WorldView-2 imagery. In *Seventh Geoinformation Science Symposium 2021* (Vol. 12082, pp. 269-278). SPIE.
- Maulidya, M., Novita, M., Hafsah, N., & Amin, N. (2019). Estimasi biomassa karbon serasah di Kawasan Hutan Gampong Deudap Pulau Nasi, Kecamatan Pulo Aceh, Kabupaten Aceh Besar. In *Prosiding Seminar Nasional Biologi, Teknologi dan Kependidikan* (Vol. 6, No. 1).
- Meyer, H., Reudenbach, C., Wöllauer, S., & Nauss, T. (2019). Importance of spatial predictor variable selection in machine learning applications—Moving from data reproduction to spatial prediction. *Ecological Modelling*, 411, 108815.
- Montgomery, D. C., Peck, E. A., & Vining, G. G. (2012). *Introduction to linear regression analysis* (5th ed.). Wiley.
- Muhsoni, F. F., Sambah, A. B., Mahmudi, M., & Wiadnya, D. G. R. (2018). Comparison of different vegetation indices for assessing mangrove density using sentinel-2 imagery. *GEOMATE Journal*, 14(45), 42-51.
- Mulyadi, E., Hendriyanto, O., & Fitriani, N. (2010). Konservasi hutan mangrove sebagai ekowisata. *Jurnal Ilmiah Teknik Lingkungan*, 2(1), 11-18.
- Mukhoriyah, M., Yudhatama, D., & Aprili, R. P. (2016). Pemantauan pemanfaatan ruang terbuka hijau kota menggunakan data SPOT 6. *Prosiding Seminar Nasional Penginderaan Jauh 2016*, 528–585. Depok: Sinas Inderaja.
- Noor, Y. R., Khazali, M., & Suryadiputra, I. N. N. (1999). *Panduan Pengenalan Mangrove di Indonesia*. PHKA/WI-IP.
- Noor, Y. R., Khazali, M., dan Suryadiputra, I. N. N. (2006). *Panduan pengenalan mangrove di Indonesia*. Wetlands International-Indonesia Programme). 2nd ed., Issue May.
- Novanda, I. G. A., Setiawati, M. D., Sugiana, I. P., Dewi, I. G. A. I. P., Andiani, A. A. E., Kamasan, M. W., ... & As-syakur, A. R. (2025). Vegetation index comparison for estimating above-ground carbon (cag) in Mangrove Forests Using Sentinel-2 Imagery: Case study from West Bali, Indonesia. *Coasts*, 5(3), 33.
- Ong, J.E., Gong, W.K., Wong, C.H. (2004). Allometry and partitioning of the mangrove, *Rhizophora apiculata*. *Forest Ecol. Manage.* 188, 395–408.

- Pahleviannur, M. R. (2019). Pemanfaatan informasi geospasial melalui interpretasi citra digital penginderaan jauh untuk monitoring perubahan penggunaan lahan. *JPIG (Jurnal Pendidikan Dan Ilmu Geografi)*, 4(2), 18-26.
- Pan, Y., Birdsey, R. A., Fang, J., Houghton, R., Kauppi, P. E., Kurz, W. A., Phillips, O. L., Shvidenko, A., Lewis, S. L., Canadell, J. G., Ciais, P., Jackson, R. B., Pacala, S. W., McGuire, A. D., Piao, S., Rautiainen, A., Sitch, S., & Hayes, D. (2011). *A large and persistent carbon sink in the world's forests. Science*, 333(6045), 988–993.
- Pham, T. D., Yokoya, N., Bui, D. T., Yoshino, K., & Friess, D. A. (2019). Remote sensing approaches for monitoring mangrove species, structure, and biomass: Opportunities and challenges. *Remote Sensing*, 11(3), 230.
- Pierson, H.O. (1993). *Handbook of carbon, graphite, diamond and fullerenes properties, processing and applications*. William Andrew Publishing, Noyes. USA: 419pp.
- Probst, P., Wright, M. N., & Boulesteix, A. L. (2019). Hyperparameters and tuning strategies for random forest. *Wiley Interdisciplinary Reviews: data mining and knowledge discovery*, 9(3), e1301.
- Purnamasari, E., Kamal, M., & Wicaksono, P. (2021). Comparison of vegetation indices for estimating above-ground mangrove carbon stocks using PlanetScope image. *Regional Studies in Marine Science*, 44, 101730.
- Putri, E. A. W., Lestariningsih, S. P., Riyono, J. N., & Prihantarto, W. J. (2023). Pemetaan kerapatan mangrove berbasis NDVI (Normalized Difference Vegetation Index) menggunakan Citra PlanetScope pada sebagian wilayah konsesi PT. Kandelial Alam. *Jurnal Laot Ilmu Kelautan*, 5(2), 142-151.
- Rachmawati, D., Setyobudiandi, I., & Hilmi, E. (2014). Potensi estimasi karbon tersimpan pada vegetasi mangrove di wilayah pesisir muara gembong Kabupaten Bekasi. *Omni-Akuatika*, 10(2)85-91.
- Richardson, A. J., & Wiegand, C. L. (1977). Distinguishing vegetation from soil background information. *Photogrammetric Engineering and Remote Sensing*, 43(12), 1541–1552.

- Risyanda, A. M. (2019). *Pemanfaatan Citra Sentinel-2A untuk estimasi kerapatan mangrove di Desa Pangkah Wetan dengan metode indeks vegetasi (NDVI, SAVI dan DVI)* (Doctoral dissertation, Universitas Brawijaya).
- Rodríguez-Sobreyra, R., Valderrama-Landeros, L., Flores-Verdugo, F., Green-Ruiz, C., & Flores-de-Santiago, F. (2025). The influence of hydrological variability, geomorphology, and salinity determines the allocation of two dominant fringe mangrove species in semiarid coastal systems. *Discover Conservation*, 2(1), 10.
- Rouse, J. W. (1974). *Monitoring the vernal advancement and retrogradation (green wave effect) of natural vegetation*.
- Setyawan, A.D. dan Winarno, K. (2006). Pemanfaatan langsung ekosistem mangrove di Jawa Tengah dan penggunaan lahan di sekitarnya; kerusakan dan upaya restorasinya. *Biodiversitas*7(3): 282-291.
- Sianturi, R. (2025). Uji normalitas sebagai syarat pengujian hipotesis. *Jurnal Pembelajaran dan Matematika Sigma*, 11(1), 1-14.
- Siddiq, A., Dimiyati, M., & Damayanti, A. (2020). Analysis of carbon stock distribution of mangrove forests in the coastal city of Benoa, Bali with combination vegetation index, and statistics approach. *International Journal on Advanced Science, Engineering and Information Technology*, 10(6), 2386-2393.
- Simamora, H. P., Khairijon, K., & Isda, M. N. (2014). *Analisis vegetasi mangrove di ekosistem mangrove Desa Tapan Nauli I Kecamatan Tapan Nauli Kabupaten Tapanuli Tengah Provinsi Sumatera Utara* (Doctoral dissertation, Riau University).
- Siregar, A. Z. (2021). Implementasi metode regresi linier berganda dalam estimasi tingkat pendaftaran mahasiswa baru. *Kesatria: Jurnal Penerapan Sistem Informasi (Komputer dan Manajemen)*, 2(3), 133-137.
- SNI (2011) 'Pengukuran dan penghitungan cadangan karbon–Pengukuran lapangan untuk penaksiran cadangan karbon hutan (ground based forest carbon accounting)', *Standar Nasional Indonesia*, 7724.
- Somantri, L. (2008). Pemanfaatan teknik penginderaan jauh untuk mengidentifikasi kerentanan dan risiko banjir. *Jurnal Geografi Gea*, 8(2).

- Stefano, A. (2019). Pemanfaatan gis (geographic information system) untuk memonitor kesehatan tanaman kelapa sawit. *Buletin LOUPE* Vol, 15(02), 8.
- Sugiyono. (2007). *Statistika untuk penelitian*. Bandung: Penerbit Alfabeta.
- Suharini, E. (2007). Menemukenali agihan permukiman kumuh di perkotaan melalui interpretasi citra penginderaan jauh. *Jurnal Geografi: Media Informasi Pengembangan dan Profesi Kegeografian*, 4(2).
- Susiana. (2011). *Diversitas dan kerapatan mangrove, gastropoda dan bivalvia di Estuari Perancak, Bali*. Skripsi. Program Studi Manajemen Suberdaya Perairan. Fakultas Ilmu Kelautan dan Perikanan. Universitas Hasanuddin: Makassar
- Sutanto. (1986). *Penginderaan Jauh Jilid 1*. Gadjah Mada University Press. Yogyakarta
- Sutaryo, D. (2009). Penghitungan biomassa sebuah pengantar untuk studi karbon dan perdagangan karbon. *Wetlands Internasional Indonesia Programme. Bogor*, 39.
- Syah, A. F. (2020). Penanaman mangrove sebagai upaya pencegahan abrasi di desa socah. *Jurnal Ilmiah Pangabdhi*, 6(1), 13-16.
- Tran, T. V., Reef, R., & Zhu, X. (2022). A review of spectral indices for mangrove remote sensing. *Remote Sensing*, 14(19), 4868.
- Wai, P., Su, H., & Li, M. (2022). Estimating aboveground biomass of two different forest types in myanmar from sentinel-2 data with machine learning and geostatistical algorithms. *Remote Sensing*, 14(9), 2146.
- White, A. T., Martosubroto, P., & Sadorra, M. S. M. (Eds.). (1989). The coastal environmental profile of Segara Anakan-Cilacap, South Java, Indonesia (Vol. 522). *WorldFish*.
- Wicaksono, P., Danoedoro, P., Hartono, & Nahren, U. (2016). Mangrove biomass carbon stock mapping of the Karimunjawa Islands using multispectral remote sensing. *International Journal of Remote Sensing*, 37(1): 26-52.
- Widiyanto, W. (2022). *Analisis multitemporal stok karbon atas permukaan hutan mangrove di sebagian Taman Nasional Karimunjawa tahun 2018-2022*. Skripsi. Fakultas Geografi, UGM Yogyakarta.

- Willmott, C. J., & Matsuura, K. (2005). Advantages of the mean absolute error (MAE) over the root mean square error (RMSE) in assessing average model performance. *Climate research*, 30(1), 79-82.
- World Resources Institute (WRI) Indonesia. (2016). Menginterpretasikan Indc: Menilai Transparansi Target Gas Rumah Kaca Pasca2020 Dari 8 Negara Penyumbang Emisi Terbesar. Washington, DC: Climate Network (OCN).
- Xue, J., & Su, B. (2017). Significant remote sensing vegetation indices: A review of developments and applications. *Journal of sensors*, 2017(1).
- Yudistira, R., Meha, A. I., & Prasetyo, S. Y. J. (2019). Perubahan konversi lahan menggunakan NDVI, EVI, SAVI dan PCA pada Citra Landsat 8 (studi kasus: Kota Salatiga). *Indonesian Journal of Computing and Modeling*, 2(1), 25-30.
- Zanne, A.E., Lopez-Gonzalez, G., Coomes, D.A., Ilic, J., Jansen, S., Lewis, S.L., Miller, R.B., Swenson, N.G., Wiemann, M.C., and Chave, J. (2009). *Global wood density database*. Dryad Digital Repository.
- Zhen, J., Liao, J., & Shen, G. (2018). Mapping mangrove forests of Dongzhaigang nature reserve in China using Landsat 8 and Radarsat-2 polarimetric SAR data. *Sensors*, 18(11), 4012.
- Zhu, Y., Liu, K., Liu, L., Wang, S., & Liu, H. (2015). Retrieval of mangrove aboveground biomass at the individual species level with WorldView-2 Images. *Remote Sensing*, 7, 12192-12214.
- Zhu, Z., & Woodcock, C. E. (2012). Object-based cloud and cloud shadow detection in Landsat imagery. *Remote Sensing of Environment*, 118, 83–94.