UNIVERSITAS
GADJAH MADA
**Uji Virulensi Nematoda Puru Akar Padi (Meloidogyne graminicola) Asal Berbagai Lokasi terhadap Tiga
Jenis Varietas Padi**
Ahmad Jodi Radityo, Prof. Dr. Ir. Siwi Indarti, M.P.
Universitas Gadjah Mada, 2026 | Diunduh dari http://etd.repository.ugm.ac.id/

DAFTAR PUSTAKA

Almaysuri, M. dan Novalina. 2025. Pengaruh Pemberian Mutagen EMS (*Ethyl Methane Sulphonate*) pada Tanaman Padi (*Oryza sativa* L) Varietas IR 64 dan Inpari 32 terhadap Karakter Kualitatif dan Kuantitatif (Doctoral dissertation, Universitas Sumatera Utara). Jurnal Agroteknologi, 13(02): 69-76

Anantha, M.S., Patel, D., Quintana, M., Swain, P., Dwivedi, J.L., Torres, R.O., Verulkar, S.B., Variar, M., Mandal, N.P., Kumar, A., Henry, A. 2016. Trait combinations that improve rice yield under drought: Sahbhagi Dhan and new drought-tolerant varieties in South Asia. Crop Science, 56 :408–421

Anindita, S., Sandrawati, A., Arifin, M., and Devnita, R. 2024. Evaluating Soil Particle Analysis Methods for Tropical Andosols in Indonesia. soilrens, 22(2): 83-90.

Arun, A., Shanthi, A., Raveendran, M., Seenivasan, N., Pushpam, R., dan Shandeep, G. 2023. An insight into occurrence, biology, and pathogenesis of rice root-knot nematode *Meloidogyne graminicola*. Biology, 12(7): 984-987.

Bellafiore, S., Jougla, C., Chapuis, E., Besnard, G., Suong, M., Vu, P. N., and Thi, X. N. 2015. Intraspecific variability of the facultative meiotic parthenogenetic root-knot nematode (Meloidogyne *graminicola*) from rice fields in Vietnam. Comptes Rendus Biologies, 338(7): 471-483.

Beretta, A. N., Silbermann, A. V., Paladino, L., Torres, D., Bassahun, D., Musselli, R., and García-Lamohte, A. 2014. Soil texture analyses using a hydrometer: modification of the Bouyoucos method. International Journal of Agriculture and Natural Resources, 41(2): 263-271.

BPS. 2025. Pada 2024, luas panen padi mencapai sekitar 10,05 juta hektare dengan produksi padi sebanyak 53,14 juta ton gabah kering giling (GKG). <https://www.bps.go.id/id/pressrelease/2025/02/03/2414/pada-2024--luas-panen-padi-mencapai-sekitar-10-05-juta-hektare-dengan-produksi-padi-sebanyak-53-14-juta-ton-gabah-kering-giling--gkg--.html>. Diakses pada 12 Desember 2025.

Bui, H. X., Oliveira, C. J., Desaeger, J. A., and Schroeder, N. E. 2022. Rice Root-Knot Nematode *Meloidogyne graminicola* (Nematoda: Chromadorea: Tylenchida: Meloidogynidae: Meloidogyne): EENY-776.

EPPO. 2016. PM 7/103 (2) *Meloidogyne enterolobii*, EPPO Bulletin, 46(2): 190–201

Cabasan, M. T. N., Kumar, A., and De Waele, D. 2012. Comparison of migration, penetration, development and reproduction of Meloidogyne graminicola on susceptible and resistant rice genotypes. Nematology, 14(4): 405-415.

Chaerani, C. 2022. Plant parasitic nematodes in agricultural ecosystem of Indonesia. Jurnal Perlindungan Tanaman Indonesia, 26(1): 1-12.

Chavan, Satish, N., and Tina, K. 2023. "*Meloidogyne graminicola*." Trends in Parasitology, 39(12):1114-1115.

Chen, B., Sikandar, A., Ahmad, S., Luo, M., and Wu, H. 2022. *Meloidogyne graminicola's* effect on growth performance of rice under low population density. Agronomy, 12(3): 587-589.

Cuc, T.T., and Prot, J.C. 1994. Nematode parasites of deepwater and irrigated rice

Uji Virulensi Nematoda Puru Akar Padi (Meloidogyne graminicola) Asal Berbagai Lokasi terhadap Tiga
Jenis Varietas Padi
Ahmad Jodi Radityo, Prof. Dr. Ir. Siwi Indarti, M.P.
Universitas Gadjah Mada, 2026 | Diunduh dari http://etd.repository.ugm.ac.id/

in the Mekong River Delta. In Proceedings of the conference held by Vietnam and IRRI: A Partnership in Rice Research, Hanoi, Vietnam, 4(7): 51–260.

EPPO. 2025. Datasheets on pests recommended for regulation. *Meloidogyne graminicola*. EPPO Bulletin, 55(2): 242-249.

Devaraja, K. P., Ellur, R. K., Pankaj, Singh, A. K., Abhishek Gowda, A. P., and Sirohi, A. 2023. Response of rice genotypes to rice root-knot nematode (*Meloidogyne graminicola*) infection under varying temperature regimes. Plant Pathology, 72(1): 179-194.

Distan Luwu. 2021. Kadis Pertanian Kunjungi Lokasi Pertanaman Padi HMS 700. <https://distan.luwukab.go.id/detailpost/kadis-pertanian-kunjungi-lokasi-pertanaman-padi-hms-700?>, Diakses pada 1 Juni 2026.

Durahman, D., Tarno, H., dan Rahardjo, B. T. 2014. Eksplorasi nematoda parasit tumbuhan pada tanaman nilam (*Pogostemon cablin* Benth) di Kecamatan Kesamben Kabupaten Blitar. Jurnal HPT (Hama Penyakit Tumbuhan), 2(4): 1-10.

Durner, W., and Iden, S. C. 2021. The improved integral suspension pressure method (ISP+) for precise particle size analysis of soil and sedimentary materials. Soil and Tillage Research, 213: 105086.

Dutta, S., Mondal, S., Hazra, A., Ghosh, S., Panja, B., Chakrabarti, M., and Mukherjee, A. 2023. Evaluation of root-gall associated fungal endophytes for the control of Meloidogyne graminicola infecting rice. Biological Control, 186: 105341.

Dutta, T. K., Powers, S. J., Kerry, B. R., Gaur, H. S., and Curtis, R. H. 2011. Comparison of host recognition, invasion, development and reproduction of *Meloidogyne graminicola* and *M. incognita* on rice and tomato. Nematology, 13(5): 509-520.

Dwiningsih, Y., and Alkahtani, J. 2022. Agronomics, genomics, breeding and intensive cultivation of Ciherang rice variety.

Eisenback, J.D. and Triantaphyllou, H.H. 1991. Root Knot Nematode: Meloidogyne Species and Races. In. Manual of Agricultural Nematology (Ed. Nickle, W.R): 191-274.

Eisenback, J.D., Hirschmann, H. and Triantaphyllou, A.C. 1980. Morphological comparison of Meloidogyne female head structures, perineal patterns and stylets. J. Nematol., 12: 300-313.

Golden, A.M. and Birchfield, W. 1965. *Meloidogyne graminicola* (Heteroderidae) a new species of rootknot nematode from grass. Proceedings of the Helminthology Society of Washington, 32: 228-231.

EURL. 2024. EURL diagnostic protocol for Meloidogyne graminicola: Morphological & molecular methods (Version 1). Zenodo: 1-48.

Fanelli, E., Cotroneo, A., Carisio, L., Troccoli, A., Grosso, S., Boero, C., Capriglia, F., and De Luca., F. 2017. Detection and molecular characterization of the rice root-knot nematode *Meloidogyne graminicola* in Italy. European Journal of Plant Pathology 149: 467–476.

Fanelli, E., Gaffuri, F., Troccoli, A., Sacchi, S., and De Luca F. 2022. New occurrence of *Meloidogyne graminicola* (Nematoda: Meloidogyninae) from rice fields in Italy: Variability and phylogenetic relationships. Ecology and Evolution 12(9): 1–14.

Favorgen. FavorPrep™ Tissue Genomic DNA Extraction HE Mini Kit.

UNIVERSITAS GADJAH MADA
Uji Virulensi Nematoda Puru Akar Padi (Meloidogyne graminicola) Asal Berbagai Lokasi terhadap Tiga Jenis Varietas Padi
Ahmad Jodi Radityo, Prof. Dr. Ir. Siwi Indarti, M.P.
Universitas Gadjah Mada, 2026 | Diunduh dari http://etd.repository.ugm.ac.id/

<https://www.favorgen.com/en/product.php?act=viewdancid=163danid=61>. Diakses pada 10 Januari 2026.

Galeng-Lawilao, J., Kumar, A., and De Waele, D. 2021. High initial population densities of Meloidogyne graminicola result in loss of resistance and tolerance of recombinant inbred lines (RIL) of Asian rice (*Oryza sativa*). Russian Journal of Nematology, 29(2): 111-126.

Golden, A.M. and W. Birchfield. 1965. *Meloidogyne graminicola* (Heteroderidae), a New Species of Root Knot Nematode from Grasses. Proceeding Helminthological Society of Washington 32: 228–231

Harian Merapi. 2025. Pemdes Madurejo Prambanan (PL) tanam padi HMS 700 sistem organik, benih unggul produktivitas panen 12 ton per hektar <https://www.harianmerapi.com/news/4016291592/pemdes-madurejo-Prambanan (PL)-tanam-padi-hms-700-sistem-organik-benih-unggul-produktivitas-panen-12-ton-per-hektar?page=2>. Diakses pada 16 Februari 2026.

BPS. 2024. Luas Panen, Produksi, dan Produktivitas Padi di Kabupaten Klaten (Kuintal/Hektar)
(Kuintal/Hektar)<https://klatenkab.bps.go.id/id/statistics-table/2/MjkxIzI=/luas-panen--produksi--dan-produktivitas-padi-di-kabupaten-klaten--kuintal-hektar---kuintal-hektar-.html>. Diakses pada 16 Februari 2026.

IRRI. 1986. Annual report for 1985. International Rice Research Institute, Los Baños.

Jain, R. K., Khan, M. R., and Kumar, V. 2012. Rice root-knot nematode (*Meloidogyne graminicola*) infestation in rice. Archives of Phytopathology and Plant Protection, 45(6): 635-645.

Jesus, A., Pedrosa, E., Regis, M., Reis, J. B. R. D. S., and Porto, A. C. F. 2020. Morphological and morphometric characterization of Pratylenchus species in sugarcane cropping areas of Pernambuco State. Revista Caatinga, 33: 599-607.

Ji, H., Gheysen, G., Denil, S., Lindsey, K., Topping, J. F., Nahar, K., and Kyndt, T. 2013. Transcriptional analysis through RNA sequencing of giant cells induced by *Meloidogyne graminicola* in rice roots. Journal of experimental botany, 64(12), 3885-3898.

Khan, M.R., Jain, R.K., Ghule, T.M., Pal, S. 2014. Root-knot Nematodes in India-A Comprehensive Monograph. In All India Coordinated Research Project on Plant Parasitic Nematodes with Integrated Approach for their Control; Indian Agricultural Research Institute: New Delhi.

Khush, G.S., Virk, P.S. 2005. IR varieties and their impact. Int. Rice Res. Inst, Los Baños

King, W. L., Grandinette, E. M., Trase, O., Rolon, M. L., Salis, H. M., Wood, H., and Bell, T. H. 2024. Autoclaving is at least as effective as gamma irradiation for biotic clearing and intentional microbial recolonization of soil. Msphere, 9(7): e00476-24.

Kumar, S., Singh, V., and Kumar, A. 2024. Response of Planting Methods and Micronutrient on Growth and Yield of Rice (*Oryza sativa* L.) in North East Plain Zone Prayagraj. International Journal of Research in Agronomy, 7(8): 655-660.

Kumar, V., Verma, K. K., and Kumar, A. 2017. Effect of soil organic matter content

UNIVERSITAS
GADJAH MADA
**Uji Virulensi Nematoda Puru Akar Padi (Meloidogyne graminicola) Asal Berbagai Lokasi terhadap Tiga
Jenis Varietas Padi**
Ahmad Jodi Radityo, Prof. Dr. Ir. Siwi Indarti, M.P.
Universitas Gadjah Mada, 2026 | Diunduh dari http://etd.repository.ugm.ac.id/

and soil texture on reproduction of *Meloidogyne graminicola* on rice. Journal of entomology and zoology studies, 5(5): 1267-1270.

Lahari, Z., Ullah, C., Kyndt, T., Gershenzon, J., and Gheysen, G. 2019. Strigolactones enhance root-knot nematode (*Meloidogyne graminicola*) infection in rice by antagonizing the jasmonate pathway. New Phytologist, 224(1): 454-465.

Lee J.H., Muhsin, M., Atienza, G.A., Kwak, D.Y., Kim S.M., De Leon, T.B., Angeles, E.R., Coloquio, E., Kondoh, H., Satoh, K., Cabunagan, R.C., Cabauatan, P.Q., Kikuchi, S., Leung, H., Choi, I.R. 2010. Single nucleotide polymorphisms in a gene for translation initiation factor (eIF4G) of rice (*Oryza sativa*) associated with resistance to Rice tungro spherical virus. Mol Plant-Microbe Interact 23:29–38

Lesta, L., Ningsih, R. M., Badriyah, B., and Felisha, N. A. 2025. Diversity and Population of Non-Parasitic Nematodes in Various Accessions of Pepper (*Piper nigrum* L.) Bangka Island. Media Pertanian, 10(1): 39-45.

Li, Q., Yang, Y., Liu, F., Li, Y., Yao, H., Peng, D., and Hu, X. 2025. Morphological and Molecular Characterization and Life Cycle of *Meloidogyne graminicola* Infecting Allium cepa. Agronomy, 15(8): 19-94.

López, R. 1991. Primer hallazgo de *Meloidogyne salasi* en arroz en la provincia de Guanacaste. Agronomía Costarricense 15(2): 189–191.

Luc, M., Sikora, R. A., dan Bridge, J. 2005. Plant parasitic nematodes in subtropical and tropical agriculture 131): 1-25.

Luo, M., Li, B. X., and Wu, H. Y. 2020. Incidence of the rice root-knot nematode, meloidogyne graminicola, in guangxi, China. The plant pathology journal, 36(3): 290-297.

Mackill, D.J., and Kush, G.S. 2018. IR 64: A high quality and high-yielding mega variety rice. Rice, 11: 1-11.

Mantelin, S., Bellafiore, S., and Kyndt, T. 2016. *Meloidogyne graminicola*: a major threat to rice agriculture. Molecular plant pathology, 18(1): 3-9.

Mayra, R. C., Soares, Vanessa S. M., Raycenne R. L., Ana M. G., Cesar B. G. 2020. Integrative taxonomy of *Meloidogyne graminicola* populations with different esterase phenotypes parasitising rice in Brazil. Nematology, 23(6): 627-643.

Monareh, J., and Ogie, T. B. 2020. Disease Control Using Biopesticide on Rice Plants (*Oryza sativa* L.). Jurnal Agroekoteknologi Terapan, 1(1): 11-13.

Mondal, S., Purohit, A., Hazra, A., Das, S., Chakrabarti, M., Khan, M. R., and Mukherjee, A. 2024. Intraspecific variability of rice root knot nematodes across diverse agroecosystems for sustainable management. Scientific Reports, 14(1): 30032.

Muhamad, K., Ebana, K., Fukuoka, S., Okuno, K. 2017. Genetic relationships among improved varieties of rice (*Oryza sativa* L.) in Indonesia over the last 60 years as revealed by morphological traits and DNA markers. Genet Resour Crop Evol 64: 701–715.

Mulyadi, M. 1997. Pengaruh Populasi Nematoda Puru Akar (*Meloidogyne graminicola*) terhadap Pertumbuhan dan Hasil Padi. Jurnal Perlindungan Tanaman Indonesia, 3(1): 17-22.

Narasimhamurthy, H. B., Ravindra, H., Sehgal, M., Ekabote, S. D., and Ganapathi, G. 2017. Bio-management of rice root-knot nematode (*Meloidogyne graminicola*). J Entomol Zool Stud, 5(4): 1433-1439.

UNIVERSITAS
GADJAH MADA
**Uji Virulensi Nematoda Puru Akar Padi (Meloidogyne graminicola) Asal Berbagai Lokasi terhadap Tiga
Jenis Varietas Padi**
Ahmad Jodi Radityo, Prof. Dr. Ir. Siwi Indarti, M.P.
Universitas Gadjah Mada, 2026 | Diunduh dari http://etd.repository.ugm.ac.id/

Narasimhamurthy, H. B., Sehgal, M., and Naik, R. G. 2023. Rice Root-Knot Nematode (*Meloidogyne graminicola*): A Major Menace in Rice Production. In Sustainable Rice Production-Challenges, Strategies and Opportunities. IntechOpen, 1(3): 54-61.

Negretti, R. R. D., Manica-Berto, R., Agostinetto, D., Thürmer, L., and Gomes, C. B. 2014. Host suitability of weeds and forage species to root-knot nematode *Meloidogyne graminicola* as a funcion of irrigation management. Planta Daninha, 32: 555-561.

Nurjayadi, M. Y., Munif, A., and Suastika, G. 2015. Identifikasi nematoda puru akar, *Meloidogyne graminicola*, pada tanaman padi di Jawa Barat. Jurnal Fitopatologi Indonesia, 11(4): 113-113.

Pankaj, S. H., Sharma, H. K., dan Prasad, J. S. 2010. The rice root-knot nematode, Meloidogyne graminicola: an emerging problem in rice-wheat cropping system. Indian Journal of Nematology, 40(1): 1-9.

Peng S, Laza, R.C., Visperas, R.M., Sanico, A.L., Cassman, K.G., Khush, G.S. 2000. Grain yield of rice cultivars and lines developed in the Philippines since 1966. Crop Sci 40: 307–314.

Perry, R. N., Moens, M., dan Starr, J. L. 2009. *Root-knot nematodes*. Wallingford, UK: CABI.

Phule, A. S., Barbadikar, K. M., Madhav, M. S., Subrahmanyam, D., Senguttuvel, P., Babu, M. P., and Kumar, P. A. 2019. Studies on root anatomy, morphology and physiology of rice grown under aerobic and anaerobic conditions. Physiology and Molecular Biology of Plants, 25(1): 197-205.

EPPO. 2016. PM 7/103 (2) *Meloidogyne enterolobii* Bulletin OEPP/EPPO Bulletin, 46(2): 190–201.

Pokharel, R. R. 2009. Damage of root-knot nematode (*Meloidogyne graminicola*) to rice in fields with different soil types. Nematologia Mediterranea, 2(2): 41-46.

Pokharel, R. R., Duxbury, J. M., and Abawai, G. 2012. Evaluation of protocol for assessing the reaction of rice and wheat germplasm to infection by *Meloidogyne graminicola*. Journal of nematology, 44(3): 274.

Prasetiyono, J., Tasliah, T., Dadang, A., dan Fatimah, F. 2013. Perbaikan Padi (*Oryza Sativa* L.) Varietas Ciherang Untuk Sifat Umur Genjah Dan Produksi Tinggi Menggunakan Marka Molekuler. Berita Biologi, 12(1): 68094.

Prot, J. C., and Matias, D. M. 1995. Effects of water regime on the distribution of *Meloidogyne graminicola* and other root-parasitic nematodes in a rice field toposequence and pathogenicity of *M. graminicola* on rice cultivar UPL R15. Nematologica, 41(2): 219-228.

Putri, V. D. 2024. Identifikasi Morfologi dan Molekuler Spesies Nematoda Puru Akar di Lima Sentra Tanaman Padi di Kabupaten Klaten, Jawa Tengah (Skripsi, Universitas Gadjah Mada).

Ragil, C. A., Liestiany, E., and Soedijo, S. 2019. Uji Efektivitas Serbuk Putri Malu (*Mimosa pudica* L.) Terhadap Serangan Nematoda *Meloidogyne* spp. Pada Tanaman Tomat. Jurnal Proteksi Tanaman Tropika, 2(3): 143-150.

Rao, Y.S., Prasad, J.S., Yadava, C.P., Padalia, C.R. 1984. Influence of rotation crops in rice soils on the dynamics of parasitic nematodes. Biol. Agric. Hortic., 2: 69–78.

Roy, K., Samanta, P., Rathod, A., and Sarkar, S. 2018. Biology of different

UNIVERSITAS GADJAH MADA
Uji Virulensi Nematoda Puru Akar Padi (Meloidogyne graminicola) Asal Berbagai Lokasi terhadap Tiga Jenis Varietas Padi
Ahmad Jodi Radityo, Prof. Dr. Ir. Siwi Indarti, M.P.
Universitas Gadjah Mada, 2026 | Diunduh dari http://etd.repository.ugm.ac.id/

geographic isolates of *Meloidogyne graminicola* Golden and Birchfield in rice (*Oryza sativa* L.) cultivar IET-4786. Current Nematology, 29(1): 21–31,

Rusinque, L., Maleita, C., Abrantes, I., Palomares-Rius, J. E., and Inácio, M. L. 2021. *Meloidogyne graminicola* A threat to rice production: Review update on distribution, biology, identification, and management. Biology, 10(11): 1163-1167.

Ruszczak, B., and Boguszewska-Mańkowska, D. 2022. Soil moisture a posteriori measurements enhancement using ensemble learning. Sensors, 22(12): 4591.

Salalia, R., and Walia, R. K. 2017. Effect of soil texture and water regimes on the pathogenicity of rice root-knot nematode, *Meloidogyne graminicola* on rice. Indian Journal of Nematology, 47(1): 136-138.

Saragih, B., P. A. R. Utoro, R. A. Prasetyo, dan Q. Aini. 2021. Pertanian dan Masa Depan. Deepublish, Yogyakarta

Shivas, R. G., Kiss, L., Geering, A. D., and McTaggart, A. R. 2023. Management of plant pathogen collections. Department of Agriculture, Fisheries and Forestry.

Shrestha, R., Al-Shugeairy, Z., Al-Ogaidi, F., Munasinghe, M., Radermacher, M., Vandenhirtz, J., Price, A.H. 2014. Comparing simple root phenotyping methods on a core set of rice genotypes. Plant Biol 16: 632–642.

Sitaresmi, T., Hairmansis, A., Widyastuti, Y., Susanto, U., Wibowo, B. P., Widiastuti, M. L., dan Nugraha, Y. 2023. Advances in the development of rice varieties with better nutritional quality in Indonesia. Journal of Agriculture and Food Research, 12: 100602.

Sofian, S., S. Sopialena, S. Suyadi, dan H. Rudiyanto. 2022. Keragaman nematoda pada lahan reklamasi pasca tambang batu bara di Desa Bangun Rejo Kecamatan Tenggarong Seberang. Agrifor: Jurnal Ilmu Pertanian dan Kehutanan 21(1): 161- 174.

Song, Z. Q., Zhang, D. Y., Liu, Y. and Cheng, F. X. 2017. First report of Meloidogyne graminicola on rice (*Oryza sativa*) in Hunan province, China. Plant Dis. 101:2153.

Soriano, I.R.S., Prot, J.C., Matias, D.M. 2000. Expression of tolerance for *Meloidogyne graminicola* in rice cultivars as affected by soil type and flooding. J. Nematol., 32: 309–317.

Suprihatno, B., Daradjat, A. A., Baehaki, S. E., Widiarta, I. N., Setyono, A., Indrasari, S. D., dan Sembiring, H. 2009. Deskripsi varietas padi. Balai Besar Penelitian Tanaman Padi, Subang.

Talukder, B., Roy, K., Kumar, P. S., dan Saha, S. 2017. Comparative Morphometric Variability of Meloidogyne graminicola Golden and Birchfield Infesting Rice (*Oryza sativa* L.) in West Bengal, India. Int. J. Curr. Microbiol. App. Sci, 6(7): 2422-2431.

Tani Makmur Nusantara. 2021. Benih Padi HMS 700< https://www.tanimakmurnusantara.com/2021/08/benih-padi-hms700.html?>, Diakses pada 1 Juni 2026.

Taylor, A. L., and Sasser, J. N. 1978. Biology, identification and control of root-knot nematodes (*Meloidogyne species*). N.C. State University Graphics: Raleigh, N.C.

Tian, Z.-L., Maria, M., Barsalote, E. M., Castillo, P. and Zheng, J.-W. 2018.

UNIVERSITAS
GADJAH MADA
**Uji Virulensi Nematoda Puru Akar Padi (Meloidogyne graminicola) Asal Berbagai Lokasi terhadap Tiga
Jenis Varietas Padi**
Ahmad Jodi Radityo, Prof. Dr. Ir. Siwi Indarti, M.P.
Universitas Gadjah Mada, 2026 | Diunduh dari http://etd.repository.ugm.ac.id/

Morphological and molecular characterization of the rice root-knot nematode, Meloidogyne graminicola, Golden and Birchfeild, 1965 occurring in Zhejiang, China. J. Integr. Agr. 17:2724-2733.

Troccoli, A., de Sa, M. G., Viaene, N., and Damme, N. 2025. PM 7/158 (1) Meloidogyne graminicola. EPPO Bulletin, 55(1).

Vinod, K. K., Verma and Anil, K. 2017. Studies on Pathogenicity of *Meloidogyne graminicola* in Different Soil types on Scented and Non-Scented Rice Int. J. Pure App. Biosci. 5 (1): 687-693.

Wahyuni, T. I. 2024. Desain Primer Spesifik untuk Deteksi Nematoda Puru Akar *(Meloidogyne graminicola)* pada Padi (Doctoral dissertation, Universitas Gadjah Mada).

Wilandari, R., Swibawa, I. G., Aeny, T. N., dan Purnomo, P. 2022. Efikasi Bionematisida *Purpureocillium lilacinum* Terhadap Nematoda Puru Akar (*Meloidogyne* spp.) Dari Dua Inang Berbeda. Jurnal Agrotek Tropika, 10(2): 187-193.

Win, P. P., Kyi, P. P., Maung, Z. T. Z., and De Waele, D. 2014. Evaluation of the host response of lowland and upland rice varieties from Myanmar to the rice root-knot nematode *Meloidogyne graminicola*. Archives of Phytopathology and Plant Protection, 47(7): 869-891.

Win, P.P., Kyi, P.P., and DeWaele, D. 2011. Effect of agro-ecosystem on the occurrence of the rice root-knot nematode *Meloidogyne graminicola* on rice in Myanmar. Australas. Plant Pathology, 40:187–196.

Wondirad, M., Baujard, P., dan Decraemer, W. 2003. Effect of biotic and abiotic factors on the morphometric variability of *Paratrichodorus rhodesiensis* (Nematoda: Trichodoridae) from Senegal. Nematology, 5(3), 463-477.

Xie, J. L., Xu, X., Yang, F., Xue, Q., Peng, Y. L. and Ji, H. L. 2019. First report of root-knot nematode, *Meloidogyne graminicola*, on rice in Sichuan province, Southwest China. Plant Dis. 103: 2142-2142.

Yang, Z., Zhang, L., Li, X., Lin, Y., Ye, S., and Ding, Z. 2023. Population dynamics of *Meloidogyne graminicola* in soil in different types of direct-seeded rice agroecosystems in Hunan Province, China. Journal of nematology, 55(1): 20230040.

Zenna, N.S., Cruz, F.C.S., Javier, E.L., Duka, I.A., Barrion, A.A., Azzam, O. 2006. Genetic analysis of tolerance to rice tungro bacilliform virus in rice (*Oryza sativa* L.) through agroinoculation. J Phytopathol, 154:197–203

Zhang, J., Zhu, Y., Yu, L., Yang, M., Zou, X., Yin, C., and Lin, Y. 2022. Research Progress in The Absorption, Transport, and Resistance of Cadmium in Rice (*Oryza sativa* L.). Sel, 11(3): 569-574.