UNIVERSITAS
GADJAH MADA
Karakteristik Morfologi dan Hubungan Morfometrik Otolith dengan Panjang dan Berat Ikan Kakap (Lutjanus vitta Quoy & Gaimard, 1824) yang Didaratkan di Pelabuhan Perikanan Pantai Bajomulyo Kabupaten Pati
Nur Diana Rizqi Azizah, Prof. Dr. Ir. Djumanto, M.Sc.
Universitas Gadjah Mada, 2026 | Diunduh dari http://etd.repository.ugm.ac.id/

DAFTAR PUSTAKA

Aguera, A., & D. Brophy. 2011. Use of sagittal otolith shape analysis to discriminate Northeast Atlantic and Western Mediterranean stocks of Atlantic saury, *Scomberesox saurus saurus* (Walbaum). Fisheries Research, 110 : 465–471.

Anderson, W. D. Jr and G. R. Allen. 2001. *Lutjanidae.* Jobfishes. p. 28402918. In K.E. Carpenter and V. Niem (eds.) FAO species identification guide for fishery purposes. The living marine resources of the Western Central Pacific. Vol. 5. Bony fishes part 3 (Menidae to Pomacentridae). FAO, Rome. 2840 – 2853.

Arai, T and S. Kimura. 2022. Spatiotemporal Variability of Trace Elements Fingerprints in Otolith of Japanese Eel *(Aguilla japonica)* and Its Use in Tracing Geographic Origin. Biology, 11(12): 1733.

Athaa, F. F., Djumanto, S. Partosuwiryo and N. Probosunu. 2023. Correlation of otolith morphometrics with total length and weight of shortfin scad *(Decapterus* macrosoma Bleeker, 1851) in the Special Region of Yogyakarta. AACL Bioflux. 16(2) : 957-969.

Bani, A., S. Poursaeid and V. M. Tuset. 2013. Comparative morphology of the sagittal otolith in three species of South *Caspian gobies.* Journal of Fish Biology, 82 : 1321–1332.

Baweleng, S., F. B. Manginsela, dan J. R. R. Sangari. 2018. Studi otolit ikan layang, Decapterus akaadsi, Abe 1958 dari perairan Teluk Amurang. Jurnal Ilmiah Platax. 6(2) : 66-76.

Budiati, O. C., Djumanto, S., Partosuwiryo and N. Probosunu. 2024. Correlation of otolith morphometry with length and weight in barracuda *Sphyraena forsteri* (Cuvier, 1829) at the Coastal Fishing Port of Tasikagung Rembang, Central Java. AACL Bioflux. 17(6) : 2685-2695.

Campana, S. E and J. M. Casselman. 1993. Stock discrimination using otolith shape analysis. Canadian Journal of Fisheries and Aquatic Sciences, 50(5): 1062-1083.

Djumanto. 2020. Hubungan panjang ikan dan ukuran otolit pada ikan *Channa striata* di Danau Rawa Pening, Jawa Tengah, Indonesia. AACL Bioflux. 13(4):1917-1924.

Gagliano, Monica and M. I. McCormick. 2004. Feeding history influences otolith shape in tropical fish. Marine Ecology Progress Series, 278: 291-296.

Gumoloc, M. T and I. R. P. Gerasmio. 2024. Stock identification of brown-striped snapper, *Lutjanus vitta* (Qouy and Gaimard, 1824), inferred from otolith shape analysis off The Coast of Iligan Bay, Mindanao, Philippines. Asian Fisheries Society, 37 : 232-242.

UNIVERSITAS
GADJAH MADA
Karakteristik Morfologi dan Hubungan Morfometrik Otolith dengan Panjang dan Berat Ikan Kakap (Lutjanus vitta Quoy & Gaimard, 1824) yang Didaratkan di Pelabuhan Perikanan Pantai Bajomulyo Kabupaten Pati
Nur Diana Rizqi Azizah, Prof. Dr. Ir. Djumanto, M.Sc.
Universitas Gadjah Mada, 2026 | Diunduh dari http://etd.repository.ugm.ac.id/

He, T., J. Cheng, J. Qin., Y. Li., dan T. Gao. 2018. Comparative analysis of otolith morphology in three species of Scomber. Ichthyological Research, 65(2) : 192-201.

Holden, M. J and D. F. S. Raitt. 1974. Manual of fisheries science. Part 2: methods of resource investigation and their application FAO, Rome.

Lombarte., Antoni., and J. Lleonart. 1993. Otolit size changes related with body growth, habitat depth and temperature. Environmental biology of fishes, 37(3) : 297-306.

Mallik, A., S. Bhushan, G. B. Sreekanth, K. K. Ramteke, A. K. Jaiswar, and B. B. Nayak. 2024. Length-weight relationship, condition factor and otolith shape indices of four creek-associated fish of the Northeastern Arabian Sea. Acta Zoologica Bulgaria, 1-8.

Miloševič, D., A. Peśič., Z. Ikica., T. Mitrovič., and N. Paskaš. 2021. Biometry of the sagittal otolit for three demersal fish species from the Eastern Adriatic Sea (Montenegro). ACTA ADRIAT, 6(2) : 171-182.

Mogea, D., L. J. L. Lumigas., dan G. E. Mamuaya. 2019. Analisis biometrik otolit (sagitta) untuk pemisahan stok ikan cakalang, Katsuwonus pelamis (Linnaeus, 1758), yang didaratkan di Pelabuhan Perikanan Pantai Tumumpa (Manado), Sulawesi Utara. Jurnal Ilmiah Platax, 7(2): 401-412.

Morales-Nin, B and J. Panfili. 2002. Age estimation. Manual of fish sclerochronology (Eds J. Panfili, H. Troadec, H. de Pontual and PJ Wright) pp: 91-98.

Mourniaty, A. Z. A., M. A. Jabbar., I. N. Suyasa., dan A. Wujudi. 2020. Hubungan morfometrik otolit dengan ukuran ikan layang deles (*Decapterus Macrosoma* Bleeker, 1851) di perairan Bali Selatan. BAWAL. 12(2): 103-107.

Newman S. J and D. McB. Williams. 1996. Variation in reef-associated assemblages of the *Lutjanidae* and *Lethrinidae* at different distances offshore in the central Great Barrier Reef. Environmental Biology of Fishes, 46(2): 123-138.

Oktaviyani, S dan W. Kurniawan. 2017. Aspek reproduksi ikan kakap *(Lutjanus vitta) (Quoy & Gaimard*, 1824) di Teluk Jakarta dan sekitarnya. Jurnal Iktiologi Indonesia. 17(2) : 215-225.

Oktaviyani, S. 2018. Mengenal marga *Lutjanus*, salah satu komoditas unggulan dalam perikanan tangkap. Oseana, XLIII (3) : 29-39.

Palla, H. P., B. J. Gonzales., F. B. Sotto., A. S. Ilano., and K. Tachihara. 2016. Age, growth and mortality of brown stripe snapper *Lutjanus vitta* (Quoy and Gaimard 1824) from West Sulu Sea, Philippines. Asian Fisheries Science, 29 : 28-42.

Ponton, D. 2006. Is geometric morphometrics efficient for comparing otolit shape of different fish species?. Journal of Morphology, 267, 750–757. http://doi.org/ 10.1002/jmor.

UNIVERSITAS
GADJAH MADA
Karakteristik Morfologi dan Hubungan Morfometrik Otolith dengan Panjang dan Berat Ikan Kakap (Lutjanus vitta Quoy & Gaimard, 1824) yang Didaratkan di Pelabuhan Perikanan Pantai Bajomulyo Kabupaten Pati
Nur Diana Rizqi Azizah, Prof. Dr. Ir. Djumanto, M.Sc.
Universitas Gadjah Mada, 2026 | Diunduh dari http://etd.repository.ugm.ac.id/

Popper, A. N and R. R. Fay. 2011. Rethinking sound detection by fishes. Hear. Res. 273: 25-36.

Popper, A. N and Z. Lu. 2000. Structure-function relationships in fish otolith organs. Fisheries Research, 46(1-3): 15-25.

Ramachandran, S., D. M. Ali and C. V. Benjamin. 2013. Age, growth and maturity of brown stripe snapper *Lutjanus vitta* (Quoy & Gaimard, 1824) from southwest coast of India. Journal of the Marine Biological Association of India. 55(2) : 61-68.

Randall J. E. 1995. *Coastal Fishes of Oman.* University of Hawaii Press, Honolulu. 439 p.

Reader, J. M., A. Spares, M. J. Stokesbury., T. S. Avery, M. J. Dadswell. 2015. Elemental fingerprints of otoliths from smolt of Atlantic salmon, *Salmo salar* Linnaeus, 1758, from three maritime watersheds: Natural tag for stock discrimination. Proceedings of the Nova Scotian Institute of Science (NSIS), 48(1), 91-123.doi:10.15273/pnsis.v48i1.5908.

Sadighzadeh, Z., V. M. Tuset., T. Valinassab., M. R. Dadpour and A. Lombarte. 2012. Comparison of different otolith shape descriptors and morphometrics for the identification of closely related species of *Lutjanus spp.* from the Persian Gulf. Mar. Bio. Res., 8 : 802-814.

Sajina, A. M., S. M. Nair., V. V. R. Suresh, T. maity., J. Mukherjee, S. K. Banik, K. M. Sandhya., R. K. Manna, and R. Samanta. 2025. Application of otolith morphology to unvenil the stock structure of Hilsa shad (*Tenualosa ilisha*) in the Bhagirathi-Hooghly river system and the associated Northern Bay of Bengal. Elsevier, 90: 104467.

Sapa, A. S., W. Kantun., dan S. Wulandari. 2024. Analisis Indeks Morfometrik Otolit Ikan Swanggi (*Priacanthus tayenus* Richardson, 1846) dari Perairan Selat Makassar. Jurnal Riset Diwa Bahari, 2(2) : 86-90.

See, M., S. Marsham, C. W. Chang, V. C. Chong., A. Sasekumar, S. K. Dhillon, and K. H. Loh. 2016. The Use of Otolith Morphometrics in Determining the Size and Species Identification of Eight Mullets (Mugiliformes: *Mugilidae*) from Malaysia. Sains Malaysiana, 45(5): 735-743.

Vignon, M., and F. Morat. 2010. Environmental and genetic determinants of otolith shape revealed by a non-indigenous tropical fish. Marine Ecology Progress Series, 411 : 231-241.

WoRMS. 2026. World Database of Fishes. *Lutjanus vitta* (Qouy & Gaimard, 1824). Accessed through: World Register of Marine Species at: https://www.marinespecies.org.

UNIVERSITAS
GADJAH MADA
Karakteristik Morfologi dan Hubungan Morfometrik Otolith dengan Panjang dan Berat Ikan Kakap (Lutjanus vitta Quoy & Gaimard, 1824) yang Didaratkan di Pelabuhan Perikanan Pantai Bajomulyo Kabupaten Pati
Nur Diana Rizqi Azizah, Prof. Dr. Ir. Djumanto, M.Sc.
Universitas Gadjah Mada, 2026 | Diunduh dari http://etd.repository.ugm.ac.id/

Wujdi, A., Prihatiningsih, dan Suwarso. 2016. Karakteristik morfologi dan hubungan morfometrik *otolit* dengan ukuran ikan lemuru (*Sardinella lemuru* Bleeker, 1853) di Selat Bali. BAWAL. 8(3): 159-172.

Zengin, M., S. Saygin and N. Polat. 2015. Otolit shape analyses and dimensions of the anchovy, *Engraulis encrasicolus* L in the Black and Marmara Seas. Sains Malaysiana, 44(5) : 657–662.

Zischke, M. T., L. Litherland, B. R. Tilyard, N. J. Stratford, E. L. Jones, and Y. Wang. 2016. Otolith morphology of four mackerel species *(Scomberomorus spp.)* in Australia: Species differentiation and prediction for fisheries monitoring and assessment. Fisheries Research, 176: 39–47.