UNIVERSITAS
GADJAH MADA
ISOLATION AND IDENTIFICATION OF TOXIC COMPOUNDS AGAINTS ARTEMIA SALINA FROM THE SPONGE XESTOSPONGIA TESTUDINARIA COLLECTED IN GILI KETAPANG, PROBOLINGGO
Muhammad Arya Pradipta, Respati Tri Swasono S.Si, M.Phil. Ph.D, Associate Professor Keiichi Konoki
Universitas Gadjah Mada, 2026 | Diunduh dari http://etd.repository.ugm.ac.id/

REFERENCES

Abbott, W.S., 1925, A method of computing the effectiveness of an insecticide, *J. Econ. Entomol.*, 18(2), 265–267.

Ahmed, Y., Sohrab, M.H., Al-Reza, S.M., Tareq, F.S., Hasan, C.M. and Sattar, M.A., 2010, Antimicrobial and cytotoxic constituents from leaves of *Sapium baccatum*, *Food Chem. Toxicol.*, 48(2), 549–552.

Asadi, M.A., Iranawati, F. and Andini, A.W., 2018, Ecology of bivalves in the intertidal area of Gili Ketapang Island, East Java, Indonesia, *AACL Bioflux*, 11(1), 55–65.

Aue, W.P., Bartholdi, E. and Ernst, R.R., 1976, Two-dimensional spectroscopy. Application to nuclear magnetic resonance, *J. Chem. Phys.*, 64(5), 2229–2246.

Bayona, L.M., Van Leeuwen, G., Erol, Ö., Swierts, T., Van Der Ent, E., De Voogd, N.J. and Choi, Y.H., 2020, Influence of geographical location on the metabolic production of giant barrel sponges (*Xestospongia* spp.) revealed by metabolomics tools, *ACS Omega*, 5(21), 12398–12408.

Bintoro, R.S., Sholihah, Q., Marjono, Ciptadi, G. and Azhar, A., 2026, Suitability of physical oceanographic conditions to Ministerial Decree No. 64/KEPMEN-KP/2020 on the establishment of Gili Ketapang Marine Protected Area, *Int. J. Mar. Eng. Innov. Res.*, in press.

Chaturvedula, V.S.P. and Prakash, I., 2012, Isolation of stigmasterol and β-sitosterol from the dichloromethane extract of *Rubus suavissimus*, *Int. Curr. Pharm. J.*, 1(9), 239–242.

Claridge, T.D.W., 2016, *High-Resolution NMR Techniques in Organic Chemistry*, 3rd Ed., Elsevier, Amsterdam.

Dobretsov, S., Dahms, H., Tsoi, M. and Qian, P., 2005, Chemical control of epibiosis by Hong Kong sponges: the effect of sponge extracts on micro- and macrofouling communities, *Mar. Ecol. Prog. Ser.*, 297, 119–129.

Dodge, J.C., Yu, J., Sardi, S.P. and Shihabuddin, L.S., 2021, Sterol auto-oxidation adversely affects human motor neuron viability and is a neuropathological feature of amyotrophic lateral sclerosis, *Sci. Rep.*, 11, 803.

El-Gamal, A., Al-Massarani, S., Shaala, L., Alahdald, A., Al-Said, M., Ashour, A., Kumar, A., Abdel-Kader, M., Abdel-Mageed, W. and Youssef, D., 2016, Cytotoxic compounds from the Saudi Red Sea sponge *Xestospongia testudinaria*, *Mar. Drugs*, 14(5), 82.

UNIVERSITAS
GADJAH MADA
ISOLATION AND IDENTIFICATION OF TOXIC COMPOUNDS AGAINTS ARTEMIA SALINA FROM THE SPONGE XESTOSPONGIA
TESTUDINARIA COLLECTED IN GILI KETAPANG, PROBOLINGGO
Muhammad Arya Pradipta, Respati Tri Swasono S.Si, M.Phil. Ph.D, Associate Professor Keiichi Konoki
Universitas Gadjah Mada, 2026 | Diunduh dari http://etd.repository.ugm.ac.id/

El-Shitany, N.A., Shaala, L.A., Abbas, A.T., Abdel-Dayem, U.A., Azhar, E.I., Ali, S.S., Van Soest, R.W.M. and Youssef, D.T.A., 2015, Evaluation of the anti-inflammatory, antioxidant and immunomodulatory effects of the organic extract of the Red Sea marine sponge *Xestospongia testudinaria* against carrageenan induced rat paw inflammation, *PLoS ONE*, 10(9), e0138917.

Eprilurahman, R., 2026, *Surat Keterangan Identifikasi Spesimen Spons* No. 013/BI/SH/III/2026, Laboratorium Sistematika Hewan, Fakultas Biologi, Universitas Gadjah Mada, Yogyakarta.

Erngren, I., Smit, E., Pettersson, C., Cárdenas, P. and Hedeland, M., 2021, The effects of sampling and storage conditions on the metabolite profile of the marine sponge *Geodia barretti*, *Front. Chem.*, 9, 662659.

Finney, D.J., 1971, *Probit Analysis*, 3rd Ed., Cambridge University Press, Cambridge.

Gafni, J., Munsch, J.A., Lam, T.H., Catlin, M.C., Costa, L.G., Molinski, T.F. and Pessah, I.N., 1997, Xestospongins: potent membrane permeable blockers of the inositol 1,4,5-trisphosphate receptor, *Neuron*, 19(3), 723–733.

Gauvin, A., Smadja, J., Aknin, M. and Gaydou, E.M., 2004, Sterol composition and chemotaxonomic considerations in relation to sponges of the genus *Xestospongia*, *Biochem. Syst. Ecol.*, 32(5), 469–476.

Gottlieb, H.E., Kotlyar, V. and Nudelman, A., 1997, NMR chemical shifts of common laboratory solvents as trace impurities, *J. Org. Chem.*, 62(21), 7512–7515.

Gribble, G.W., 1999, The diversity of naturally occurring organobromine compounds, *Chem. Soc. Rev.*, 28(5), 335–346.

Hadi, T.A., Hafizt, M., Hadiyanto, Budiyanto, A. and Siringoringo, R.M., 2018, Shallow water sponges along the south coast of Java, Indonesia, *Biodiversitas*, 19(2), 535–543.

He, F., Mai, L., Longeon, A., Copp, B., Loaëc, N., Bescond, A., Meijer, L. and Bourguet-Kondracki, M., 2015a, Novel adociaquinone derivatives from the Indonesian sponge *Xestospongia* sp., *Mar. Drugs*, 13(5), 2617–2628.

He, W.F., Liang, L.F., Cai, Y.S., Gao, L.X., Li, Y.F., Li, J., Liu, H.L. and Guo, Y.W., 2015b, Brominated polyunsaturated lipids with protein tyrosine phosphatase-1B inhibitory activity from Chinese marine sponge *Xestospongia testudinaria*, *J. Asian Nat. Prod. Res.*, 17(8), 861–866.

Hooper, J.N.A. and Van Soest, R.W.M., 2002, *Systema Porifera, A Guide to the Classification of Sponges*, Kluwer Academic/Plenum Publishers, New York.

UNIVERSITAS
GADJAH MADA
ISOLATION AND IDENTIFICATION OF TOXIC COMPOUNDS AGAINTS ARTEMIA SALINA FROM THE SPONGE XESTOSPONGIA
TESTUDINARIA COLLECTED IN GILI KETAPANG, PROBOLINGGO
Muhammad Arya Pradipta, Respati Tri Swasono S.Si, M.Phil. Ph.D, Associate Professor Keiichi Konoki
Universitas Gadjah Mada, 2026 | Diunduh dari http://etd.repository.ugm.ac.id/

Huang, L., Zhu, X., Zhou, S., Cheng, Z., Shi, K., Zhang, C. and Shao, H., 2021, Phthalic acid esters: natural sources and biological activities, *Toxins*, 13(7), 495.

Ibraham, A., Rudiyansyah, R. and Alimuddin, A.H., 2021, Karakterisasi struktur lipid dari spons *Xestospongia testudinaria*, *Indo. J. Pure Appl. Chem.*, 4(1), 30–39.

Indrayani, E., Riniwati, H., Asmara, R., Solimun, S., Susilo, E. and Utami, T.N., 2025, The role of management institutions and community capacity and their suitability to the needs of Gili Ketapang Conservation Area, *WSEAS Trans. Environ. Dev.*, 21, 145–155.

Ishiyama, M., 1997, A highly water-soluble disulfonated tetrazolium salt as a chromogenic indicator for NADH as well as cell viability, *Talanta*, 44(7), 1299–1305.

Jiang, W., Liu, D., Deng, Z., De Voogd, N.J., Proksch, P. and Lin, W., 2011, Brominated polyunsaturated lipids and their stereochemistry from the Chinese marine sponge *Xestospongia testudinaria*, *Tetrahedron*, 67(1), 58–68.

Jomori, T., Higa, N., Tyas, T.A., Matsuura, N., Ueda, Y., Suetake, A., Miyazaki, S., Watanabe, S., Arizono, S., Hayashi, Y., Yasumoto, K., Ise, Y., Wakimoto, T., Yasumoto-Hirose, M., Tanaka, J. and Mori-Yasumoto, K., 2025, Onnamides and a novel analogue, onnamide G, as potent leishmanicidal agents, *Mar. Biotechnol.*, 27(5), 132.

Juniarta, H.P., Susilo, E. and Primyastanto, M., 2013, Kajian profil kearifan lokal masyarakat pesisir Pulau Gili Ketapang Kecamatan Sumberasih Kabupaten Probolinggo Jawa Timur, *J. ECSOFiM*, 1(1), 11–25.

Kalalinggi, S.Y., Soekamto, N.H., Firdaus, Appa, F.E. and Irmawan, M., 2023, Phytochemical screening of n-hexane extract of sponge *Xestospongia testudinaria* from Spermonde Archipelago, *J. Biotrop. Res. Nat. Technol.*, 1(2), 86–91.

Khodzori, F.A., Mazlan, N.B., Chong, W.S., Ong, K.H., Palaniveloo, K. and Shah, M.D., 2023, Metabolites and bioactivity of the marine *Xestospongia* sponges (Porifera, Demospongiae, Haplosclerida) of Southeast Asian waters, *Biomolecules*, 13(3), 484.

Kind, T. and Fiehn, O., 2006, Metabolomic database annotations via query of elemental compositions, mass accuracy is insufficient even at less than 1 ppm, *BMC Bioinformatics*, 7, 234.

Kind, T. and Fiehn, O., 2007, Seven golden rules for heuristic filtering of molecular formulas obtained by accurate mass spectrometry, *BMC Bioinformatics*, 8, 105.

UNIVERSITAS
GADJAH MADA
ISOLATION AND IDENTIFICATION OF TOXIC COMPOUNDS AGAINTS ARTEMIA SALINA FROM THE SPONGE XESTOSPONGIA
TESTUDINARIA COLLECTED IN GILI KETAPANG, PROBOLINGGO
Muhammad Arya Pradipta, Respati Tri Swasono S.Si, M.Phil. Ph.D, Associate Professor Keiichi Konoki
Universitas Gadjah Mada, 2026 | Diunduh dari http://etd.repository.ugm.ac.id/

Kovalerchik, D., Zovko, A., Hååg, P., Sierakowiak, A., Viktorsson, K., Lewensohn, R., Ilan, M. and Carmeli, S., 2022, Cytotoxic alkylynols of the sponge *Cribrochalina vasculum*: structure, synthetic analogs and SAR studies, *Mar. Drugs*, 20(4), 265.

Krisnawati, S. and Hidayah, Z., 2020, Pemetaan terumbu karang Pulau Gili Ketapang Probolinggo, *Juvenil*, 1(4), 437–450.

Li, D., Xu, Y., Shao, C., Yang, R., Zheng, C., Chen, Y., Fu, X., Qian, P., She, Z., De Voogd, N.J. and Wang, C., 2012, Antibacterial bisabolane-type sesquiterpenoids from the sponge-derived fungus *Aspergillus* sp., *Mar. Drugs*, 10(1), 234–241.

Lim, Y.J., Lee, C., Hong, J., Kim, D., Im, K.S. and Jung, J.H., 2001, Cytotoxic polyacetylenic alcohols from the marine sponge *Petrosia* species, *J. Nat. Prod.*, 64(12), 1565–1567.

Luu, P.V., Nguyen, H.M., Phan, P.M., Vo, A.D. and Ton-Nu, H.L., 2025, Testusterol, a new sterol of the sponge species *Xestospongia testudinaria* from Phu Quoc island, Vietnam, *Nat. Prod. Res.*, 39(15), 4382–4390.

Manger, R.L., Leja, L.S., Lee, S.Y., Hungerford, J.M. and Wekell, M.M., 1993, Tetrazolium-based cell bioassay for neurotoxins active on voltage-sensitive sodium channels: semiautomated assay for saxitoxins, brevetoxins, and ciguatoxins, *Anal. Biochem.*, 214(1), 190–194.

Meyer, B., Ferrigni, N., Putnam, J., Jacobsen, L., Nichols, D. and McLaughlin, J., 1982, Brine shrimp: a convenient general bioassay for active plant constituents, *Planta Med.*, 45(5), 31–34.

Moon, S., MacMillan, J.B., Olmstead, M.M., Ta, T.A., Pessah, I.N. and Molinski, T.F., 2002, (+)-7S-Hydroxyxestospongin A from the marine sponge *Xestospongia* sp. and absolute configuration of (+)-xestospongin D, *J. Nat. Prod.*, 65(3), 249–254.

Nakagawa, M., Endo, M., Tanaka, N. and Gen-Pei, L., 1984, Structures of xestospongin A, B, C and D, novel vasodilative compounds from marine sponge, *Tetrahedron Lett.*, 25(30), 3227–3230.

Nguta, J.M., Mbaria, J.M., Gakuya, D.W., Gathumbi, P.K., Kabasa, J.D. and Kiama, S.G., 2012, Evaluation of acute toxicity of crude plant extracts from Kenyan biodiversity using brine shrimp, *Artemia salina* L. (Artemiidae), *Open Conf. Proc. J.*, 3, 30–34.

Nguyen, H.M., Ito, T., Win, N.N., Vo, H.Q., Nguyen, H.T. and Morita, H., 2019, A new sterol from the Vietnamese marine sponge *Xestospongia testudinaria* and its biological activities, *Nat. Prod. Res.*, 33(8), 1175–1181.

Ntungwe, E.N., Domínguez-Martín, E.M., Roberto, A., Tavares, J., Isca, V.M.S., Pereira, P., Cebola, M.J. and Rijo, P., 2020, *Artemia* species: an important

UNIVERSITAS GADJAH MADA
ISOLATION AND IDENTIFICATION OF TOXIC COMPOUNDS AGAINTS ARTEMIA SALINA FROM THE SPONGE XESTOSPONGIA
TESTUDINARIA COLLECTED IN GILI KETAPANG, PROBOLINGGO
Muhammad Arya Pradipta, Respati Tri Swasono S.Si, M.Phil. Ph.D, Associate Professor Keiichi Konoki
Universitas Gadjah Mada, 2026 | Diunduh dari http://etd.repository.ugm.ac.id/

tool to screen general toxicity samples, *Curr. Pharm. Des.*, 26(24), 2892–2908.

Pham, N.B., Butler, M.S., Hooper, J.N.A., Moni, R.W. and Quinn, R.J., 1999, Isolation of xestosterol esters of brominated acetylenic fatty acids from the marine sponge *Xestospongia testudinaria*, *J. Nat. Prod.*, 62(10), 1439–1442.

Setiawan, E., De Voogd, N.J., Swierts, T., Hooper, J.N., Wörheide, G. and Erpenbeck, D., 2015, MtDNA diversity of the Indonesian giant barrel sponge *Xestospongia testudinaria* (Porifera: Haplosclerida), implications from partial cytochrome oxidase 1 sequences, *J. Mar. Biol. Assoc. U.K.*, 96(2), 323–332.

Sibero, M.T., Zhou, T., Igarashi, Y., Radjasa, O.K., Sabdono, A., Trianto, A., Bachtiarini, T.U. and Bahry, M.S., 2020, Chromanone-type compounds from marine sponge-derived *Daldinia eschscholtzii* KJMT FP 4.1, *J. Appl. Pharm. Sci.*, 10(1), 1–7.

Silverstein, R.M., Webster, F.X., Kiemle, D.J. and Bryce, D.L., 2014, *Spectrometric Identification of Organic Compounds*, 8th Ed., John Wiley and Sons, Hoboken.

Snyder, L.R., 1968, *Principles of Adsorption Chromatography*, Marcel Dekker, New York.

Still, W.C., Kahn, M. and Mitra, A., 1978, Rapid chromatographic technique for preparative separations with moderate resolution, *J. Org. Chem.*, 43(14), 2923–2925.

Swantara, I.M.D. and Rita, W.S., 2018, Toxicity of sponge extract *Xestospongia testudinaria*, *Int. J. Eng. Sci. Invent.*, 7(5), 55–58.

Swantara, M.D., Rita, W.S., Suartha, N. and Agustina, K.K., 2019, Anticancer activities of toxic isolate of *Xestospongia testudinaria* sponge, *Vet. World*, 12(9), 1434–1440.

Tai, B.H., Hang, D.T.T., Thung, D.C., Trang, D.T., Yen, P.H., Yen, D.T.H., Huong, P.T.T., Dung, D.T., Cuong, N.C., Thao, D.T., Nhiem, N.X. and Kiem, P.V., 2022, Xestospongiene VN1, a new brominated polyunsaturated lipid from marine sponge *Xestospongia testudinaria* (Lamarck, 1815), *Vietnam J. Chem.*, 60(4), 540–545.

Tominaga, H., Ishiyama, M., Ohseto, F., Sasamoto, K., Hamamoto, T., Suzuki, K. and Watanabe, M., 1999, A water-soluble tetrazolium salt useful for colorimetric cell viability assay, *Anal. Commun.*, 36(2), 47–50.

Viallon, J., Chinain, M. and Darius, H.T., 2020, Revisiting the neuroblastoma cell-based assay (CBA-N2a) for the improved detection of marine toxins active on voltage gated sodium channels (VGSCs), *Toxins*, 12(5), 281.