UNIVERSITAS
GADJAH MADA
Hubungan Perubahan NLR (Neutrofil Limfosit Rasio) sebagai Indikator Respon Antibiotik terhadap Pasien Sepsis di ICU RSUP DR. Sardjito Yogyakarta
TRIWIDODO BUDI MULYANTO, dr. Bowo Adiyanto, M.Sc, Sp.An-TI, Subsps.TI(K).; Dr.dr. Akhmad Yun Jufan, M.Sc
Universitas Gadjah Mada, 2026 | Diunduh dari http://etd.repository.ugm.ac.id/

DAFTAR PUSTAKA

Agustina, A.Y. (2024) *Perbandingan antara CRP/Albumin dengan Skor SOFA Tterhadap Angka Mortalitas Pasien Sepsis di ICU*. Universitas Gadjah Mada. Available at: https://etd.repository.ugm.ac.id/penelitian/detail/248024 (Accessed: November 5, 2025).

Alberts B, Johnson A, Lewis J, et al (2002) *Molecular Biology of the Cell*. 4th edition. New York: Garland Science.

Becker, K.L., Snider, R. and Nylen, E.S. (2010) "Procalcitonin in sepsis and systemic inflammation: a harmful biomarker and a therapeutic target," *British Journal of Pharmacology*, 159(2), pp. 253–264. Available at: https://doi.org/10.1111/j.1476-5381.2009.00433.x.

Buonacera, A. *et al.* (2022) "Neutrophil to Lymphocyte Ratio: An Emerging Marker of the Relationships between the Immune System and Diseases," *International Journal of Molecular Sciences*, 23(7), p. 3636. Available at: https://doi.org/10.3390/ijms23073636.

Caterino JM, Kahan S (2012) *Master Plan Kedaruratan Medik*. Indonesia: Binarupa Aksara Publisher.

Dahlan, M. S. (2013) *Besar Sampel dan Cara Pengambilan Sampel dalam Penelitian Kedokteran dan Kesehatan.* Ketiga. Jakarta: Salemba Medika.

De Jong, E. *et al.* (2016) "Efficacy and safety of procalcitonin guidance in reducing the duration of antibiotic treatment in critically ill patients: a randomised, controlled, open-label trial," *The Lancet Infectious Diseases*, 16(7), pp. 819–827. Available at: https://doi.org/10.1016/S1473-3099(16)00053-0.

Dobson, G.P., Letson, H.L. and Morris, J.L. (2024) "Revolution in sepsis: a symptoms-based to a systems-based approach?," *Journal of Biomedical Science*, 31(1), p. 57. Available at: https://doi.org/10.1186/s12929-024-01043-4.

Drăgoescu, A.N. *et al.* (2021) "Neutrophil to Lymphocyte Ratio (NLR)—A Useful Tool for the Prognosis of Sepsis in the ICU," *Biomedicines*, 10(1), p. 75. Available at: https://doi.org/10.3390/biomedicines10010075.

Evans, L. *et al.* (2021) "Surviving Sepsis Campaign: International Guidelines for Management of Sepsis and Septic Shock 2021," *Critical Care Medicine*, 49(11), pp. e1063–e1143. Available at: https://doi.org/10.1097/CCM.0000000000005337.

UNIVERSITAS
GADJAH MADA
Hubungan Perubahan NLR (Neutrofil Limfosit Rasio) sebagai Indikator Respon Antibiotik terhadap Pasien Sepsis di ICU RSUP DR. Sardjito Yogyakarta
TRIWIDODO BUDI MULYANTO, dr. Bowo Adiyanto, M.Sc, Sp.An-TI, Subsps.TI(K).; Dr.dr. Akhmad Yun Jufan, M.Sc
Universitas Gadjah Mada, 2026 | Diunduh dari http://etd.repository.ugm.ac.id/

Fauci AS, Kasper DL, Longo DL, Loscalzo J, et al (2009) "Harrison Manual Kedokteran." Indonesia: Karisma Publising Group.

Guyton, A. C. & Hall, E. J (2006) *Buku Ajar Fisiologi Kedokteran*. 11 ed. Jakarta: EGC.

Hotchkiss, R.S., Monneret, G. and Payen, D. (2013) "Sepsis-induced immunosuppression: from cellular dysfunctions to immunotherapy," *Nature Reviews Immunology*, 13(12), pp. 862–874. Available at: https://doi.org/10.1038/nri3552.

Iskandar, A. and Siska, F. (2025) "Analisis Hubungan Sequential Organ Failure Assessment (Sofa) Score Dengan Mortalitas Pasien Sepsis," *Jurnal Kesehatan Andalas*, 9(2), pp. 168–173. Available at: https://doi.org/10.25077/jka.v9i2.1221.

Khilnani, G.C. *et al.* (2022) "Guidelines for the Use of Procalcitonin for Rational Use of Antibiotics," *Indian Journal of Critical Care Medicine : Peer-reviewed, Official Publication of Indian Society of Critical Care Medicine*, 26(Suppl 2), pp. S77–S94. Available at: https://doi.org/10.5005/jp-journals-10071-24326.

Leliefeld, P.H.C. *et al.* (2016) "The role of neutrophils in immune dysfunction during severe inflammation," *Critical Care*, 20(1), p. 73. Available at: https://doi.org/10.1186/s13054-016-1250-4.

Li, X. *et al.* (2024) "Neutrophil-to-lymphocyte ratio, monocyte-to-lymphocyte ratio, platelet-to-lymphocyte ratio associated with 28-day all-cause mortality in septic patients with coronary artery disease: a retrospective analysis of MIMIC-IV database," *BMC Infectious Diseases*, 24(1), p. 749. Available at: https://doi.org/10.1186/s12879-024-09516-5.

Li, Y. *et al.* (2019) "The regulatory roles of neutrophils in adaptive immunity," *Cell Communication and Signaling*, 17(1), p. 147. Available at: https://doi.org/10.1186/s12964-019-0471-y.

Liang, P. and Yu, F. (2022) "Predictive Value of Procalcitonin and Neutrophil-to-Lymphocyte Ratio Variations for Bloodstream Infection with Septic Shock," *Medical Science Monitor*, 28. Available at: https://doi.org/10.12659/MSM.935966.

Liu, Q. *et al.* (2023) "Serum Cholinesterase, C-reactive Protein, Interleukin 6, and Procalcitonin Levels as Predictors of Mortality in Patients in the Intensive Care Unit," *Turkish Journal of Anaesthesiology and Reanimation*, 51(5), pp. 408–413. Available at: https://doi.org/10.4274/TJAR.2023.231349.

UNIVERSITAS
GADJAH MADA
Hubungan Perubahan NLR (Neutrofil Limfosit Rasio) sebagai Indikator Respon Antibiotik terhadap Pasien Sepsis di ICU RSUP DR. Sardjito Yogyakarta
TRIWIDODO BUDI MULYANTO, dr. Bowo Adiyanto, M.Sc, Sp.An-TI, Subsps.TI(K).; Dr.dr. Akhmad Yun Jufan, M.Sc
Universitas Gadjah Mada, 2026 | Diunduh dari http://etd.repository.ugm.ac.id/

Pasquau-Liaño, J. (2022) "Timing in antibiotic therapy: when and how to start, de-escalate and stop antibiotic therapy. Proposals from a stablished antimicrobial stewardship program," *Revista Española de Quimioterapia*, 35(Suppl3), pp. 102–107. Available at: https://doi.org/10.37201/req/s03.22.2022.

Paudel, R. *et al.* (2020) "Procalcitonin: A promising tool or just another overhyped test?," *Int. J. Med. Sci.*, 17.

Purba, A.K.R. *et al.* (2020) "The burden and costs of sepsis and reimbursement of its treatment in a developing country: An observational study on focal infections in Indonesia," *International Journal of Infectious Diseases*, 96, pp. 211–218. Available at: https://doi.org/10.1016/j.ijid.2020.04.075.

Radji, M., Fauziah, S. and Aribinuko, N. (2011) "Antibiotic sensitivity pattern of bacterial pathogens in the intensive care unit of Fatmawati Hospital, Indonesia," *Asian Pacific Journal of Tropical Biomedicine*, 1(1), pp. 39–42. Available at: https://doi.org/10.1016/S2221-1691(11)60065-8.

Reinhart, K. *et al.* (2012) "New Approaches to Sepsis: Molecular Diagnostics and Biomarkers," *Clinical Microbiology Reviews*, 25(4), pp. 609–634. Available at: https://doi.org/10.1128/CMR.00016-12.

Rhodes, A. *et al.* (2017) "Surviving Sepsis Campaign: International Guidelines for Management of Sepsis and Septic Shock: 2016," *Intensive Care Medicine*, 43(3), pp. 304–377. Available at: https://doi.org/10.1007/s00134-017-4683-6.

Rudd, K.E. *et al.* (2020) "Global, regional, and national sepsis incidence and mortality, 1990–2017: analysis for the Global Burden of Disease Study," *The Lancet*, 395(10219), pp. 200–211. Available at: https://doi.org/10.1016/S0140-6736(19)32989-7.

Runge MS, Greganti MA 1. (2009) *Netter's Internal Medicine*. 2nd ed. Philadelphia USA: Saunders Elsevier.

Sari, R. *et al.* (2019) "Neutrophil to lymphocyte ratio as a predictor of treatment response and mortality in septic shock patients in the intensive care unit," *TURKISH JOURNAL OF MEDICAL SCIENCES*, 49(5), pp. 1336–1349. Available at: https://doi.org/10.3906/sag-1901-105.

Schuetz, P. *et al.* (2017) "Procalcitonin to initiate or discontinue antibiotics in acute respiratory tract infections," *Cochrane Database of Systematic Reviews*. Edited by Cochrane Acute Respiratory Infections Group, 2019(5). Available at: https://doi.org/10.1002/14651858.CD007498.pub3.

UNIVERSITAS
GADJAH MADA
Hubungan Perubahan NLR (Neutrofil Limfosit Rasio) sebagai Indikator Respon Antibiotik terhadap Pasien Sepsis di ICU RSUP DR. Sardjito Yogyakarta
TRIWIDODO BUDI MULYANTO, dr. Bowo Adiyanto, M.Sc, Sp.An-TI, Subsps.TI(K).; Dr.dr. Akhmad Yun Jufan, M.Sc
Universitas Gadjah Mada, 2026 | Diunduh dari http://etd.repository.ugm.ac.id/

Shen, X. *et al.* (2017) "Neutrophil dysregulation during sepsis: an overview and update," *Journal of Cellular and Molecular Medicine*, 21(9), pp. 1687–1697. Available at: https://doi.org/10.1111/jcmm.13112.

Singer, M. *et al.* (2016) "The Third International Consensus Definitions for Sepsis and Septic Shock (Sepsis-3)," *JAMA*, 315(8), p. 801. Available at: https://doi.org/10.1001/jama.2016.0287.

Suba, M.-I. *et al.* (2026) "Early ΔNLR Outperforms Baseline Inflammatory Markers in Predicting Short-Term Outcomes in Sepsis," *Diagnostics*, 16(10), p. 1473. Available at: https://doi.org/10.3390/diagnostics16101473.

Taylor, S.P. *et al.* (2024) "Real-World Implications of Updated Surviving Sepsis Campaign Antibiotic Timing Recommendations*," *Critical Care Medicine*, 52(7), pp. 1002–1006. Available at: https://doi.org/10.1097/CCM.0000000000006240.

Uzan, M.M., Balcı, U.G. and Yılmazer, T.T. (2025) "Neutrophil/Lymphocyte Ratio as a Marker of Infections in Patients Hospitalized in the Oncology Palliative Care Service," *The Anatolian Journal of General Medical Research* [Preprint]. Available at: https://doi.org/10.4274/anatoljmed.2025.63825.

Van Der Poll, T., Shankar-Hari, M. and Wiersinga, W.J. (2021) "The immunology of sepsis," *Immunity*, 54(11), pp. 2450–2464. Available at: https://doi.org/10.1016/j.immuni.2021.10.012.

Van Engelen, T.S.R., Joost Wiersinga, W. and Van Der Poll, T. (2018) "Pathogenesis of Sepsis," in W.J. Wiersinga and C.W. Seymour (eds.) *Handbook of Sepsis*. Cham: Springer International Publishing, pp. 31–43. Available at: https://doi.org/10.1007/978-3-319-73506-1_3.

Vincent, J.-L. *et al.* (2006) "Sepsis in European intensive care units: Results of the SOAP study*:," *Critical Care Medicine*, 34(2), pp. 344–353. Available at: https://doi.org/10.1097/01.CCM.0000194725.48928.3A.

WHO (2024) *Sepsis*. Available at: https://www.who.int/news-room/fact-sheets/detail/sepsis (Accessed: May 30, 2025).

Wibowo, E.A.A., Kurniawaty, J. and Jufan, A.Y. (2020) "HUBUNGAN ANTARA NEUTROPHYL-LYMPHOCYTE RATIO DAN KADAR PROKALSITONIN PLASMA PADA PASIEN SEPSIS YANG DIRAWAT DI RUANG INTENSIF RSUP DR SARDJITO," *Jurnal Komplikasi Anestesi*, 7(2), pp. 9–20. Available at: https://doi.org/10.22146/jka.v7i2.7446.

UNIVERSITAS
GADJAH MADA
Hubungan Perubahan NLR (Neutrofil Limfosit Rasio) sebagai Indikator Respon Antibiotik terhadap Pasien Sepsis di ICU RSUP DR. Sardjito Yogyakarta
TRIWIDODO BUDI MULYANTO, dr. Bowo Adiyanto, M.Sc, Sp.An-TI, Subsps.TI(K).; Dr.dr. Akhmad Yun Jufan, M.Sc
Universitas Gadjah Mada, 2026 | Diunduh dari http://etd.repository.ugm.ac.id/

Wirz, Y. *et al.* (2018) "Effect of procalcitonin-guided antibiotic treatment on clinical outcomes in intensive care unit patients with infection and sepsis patients: a patient-level meta-analysis of randomized trials," *Critical Care*, 22(1), p. 191. Available at: https://doi.org/10.1186/s13054-018-2125-7.

Yadav, H. and Cartin-Ceba, R. (2016) "Balance between Hyperinflammation and Immunosuppression in Sepsis," *Seminars in Respiratory and Critical Care Medicine*, 37(01), pp. 042–050. Available at: https://doi.org/10.1055/s-0035-1570356.