Kadar Bilirubin Neonatus Aterm yang Diukur dengan Transcutaneous Bilirubinometri pada Hari Pertama
Sebagai Faktor Prognostik Fototerapi pada Minggu Pertama Kehidupan Neonatus
Ardiles Varian, Dr. dr. Tunjung Wibowo, MPH, M.Kes, Sp.A(K); dr. Endy Paryanto Prawirohartono, MPH, Sp.A(K)
Universitas Gadjah Mada, 2026 | Diunduh dari http://etd.repository.ugm.ac.id/

References:

1. Lee B, Piersante T, Calkins KL. Neonatal Hyperbilirubinemia. Pediatr Ann. 2022 Jun;51(6):e219-e227.

2. Lauer BJ, Spector ND. Hyperbilirubinemia in the Newborn. Pediatrics in Review. 2011; 32(8), 341–349.

3. National Institute for Health and Care Excellence. Neonatal jaundice: clinical guideline 98. May, 2010. https://www.nice.org.uk/guidance/ cg98 (accessed June 24, 2026).

4. Bhutani VK. Editorial: building evidence to manage newborn jaundice worldwide. Indian J Pediatr. 2012 Feb;79(2):253-5.

5. Greco C, Arnolda G, Boo NY, Iskander IF, Okolo AA, Rohsiswatmo R, Shapiro SM, Watchko J, Wennberg RP, Tiribelli C, Coda Zabetta CD. Neonatal Jaundice in Low- and Middle-Income Countries: Lessons and Future Directions from the 2015 Don Ostrow Trieste Yellow Retreat. Neonatology. 2016;110(3):172-80.

6. Campbell OM, Cegolon L, Macleod D, Benova L. Length of Stay After Childbirth in 92 Countries and Associated Factors in 30 Low- and Middle-Income Countries: Compilation of Reported Data and a Cross-sectional Analysis from Nationally Representative Surveys. PLoS Med. 2016 Mar 8;13(3):e1001972.

7. Bowers J, Cheyne H. Reducing the length of postnatal hospital stay: implications for cost and quality of care. BMC Health Serv Res. 2016 Jan 15;16:16.

8. Jain M, Bang A, Tiwari A, Jain S. Prediction of significant hyperbilirubinemia in term neonates by early non-invasive bilirubin measurement. World J Pediatr. 2017 Jun;13(3):222-227.

9. Jackson GL, Saumur M, Chandwani V, Engle WD. Evaluation of Early Transcutaneous Bilirubinometry to Predict Subsequent Hyperbilirubinemia in Neonates Admitted to a Well-Baby Nursery. Am J Perinatol. 2015 Aug;32(10):944-51.

Kadar Bilirubin Neonatus Aterm yang Diukur dengan Transcutaneous Bilirubinometri pada Hari Pertama
Sebagai Faktor Prognostik Fototerapi pada Minggu Pertama Kehidupan Neonatus
Ardiles Varian, Dr. dr. Tunjung Wibowo, MPH, M.Kes, Sp.A(K); dr. Endy Paryanto Prawirohartono, MPH, Sp.A(K)
Universitas Gadjah Mada, 2026 | Diunduh dari http://etd.repository.ugm.ac.id/

10. Kemper AR, Newman TB, Slaughter JL, Maisels MJ, Watchko JF, Downs SM et al. Clinical Practice Guideline Revision: Management of Hyperbilirubinemia in the Newborn Infant 35 or More Weeks of Gestation. Pediatrics. 2022; 150 (3): e2022058859.

11. Chowdhary S, Devi U, Giridhar S. Predicting Significant Hyperbilirubinemia in ABO Incompatibility: Is Cord Direct Antiglobulin Test Useful? Indian J Hematol Blood Transfus. 2022 Jul;38(3):591-595.

12. Khurana R, Batra P, Faridi M, Khan N. Revisiting ABO incompatibility as a risk factor for significant neonatal hyperbilirubinemia. Trop Doct. 2019 Jul;49(3):201-204.

13. Suda-Całus M, Dąbrowska K, Gulczyńska E. Infant of a diabetic mother: clinical presentation, diagnosis and treatment. Pediatr Endocrinol Diabetes Metab. 2024;30(1):36-41.

14. He J, Song J, Zou Z, Fan X, Tian R, Xu J et al. Association between neonatal hyperbilirubinemia and hypoglycemia in Chinese women with diabetes in pregnancy and influence factors. Sci Rep. 2022 Oct 10;12(1):16975.

15. Kallem VR, Pandita A, Pillai A. Infant of diabetic mother: what one needs to know? J Matern Fetal Neonatal Med. 2020 Feb;33(3):482-492.

16. Yang J, Cummings EA, O'connell C, Jangaard K. Fetal and neonatal outcomes of diabetic pregnancies. Obstet Gynecol. 2006 Sep;108(3 Pt 1):644-50.

17. Mimouni F, Miodovnik M, Siddiqi TA, Butler JB, Holroyde J, Tsang RC. Neonatal polycythemia in infants of insulin-dependent diabetic mothers. Obstet Gynecol. 1986 Sep;68(3):370-2.

18. Hanin EA, Rayan H, Hani T, Taleb J, Dany AH, Lama C. Breastfeeding and Readmission for Hyperbilirubinemia in Late Preterm and Term Infants in Beirut, Lebanon. Indian Pediatr. 2022 Mar 15;59(3):218-221.

UNIVERSITAS
GADJAH MADA
Kadar Bilirubin Neonatus Aterm yang Diukur dengan Transcutaneous Bilirubinometri pada Hari Pertama
Sebagai Faktor Prognostik Fototerapi pada Minggu Pertama Kehidupan Neonatus
Ardiles Varian, Dr. dr. Tunjung Wibowo, MPH, M.Kes, Sp.A(K); dr. Endy Paryanto Prawirohartono, MPH, Sp.A(K)
Universitas Gadjah Mada, 2026 | Diunduh dari http://etd.repository.ugm.ac.id/

19. Allam JB, Kumar S, Kurmi R, Ara R, Kumar AM. First Day Serum Bilirubin Level, as Predictor of Significant Hyperbilirubinemia in Neonates. Archives of Medicine and Health Sciences. 2018; 6(2): p 218-222.

20. Triasih R, Haksari EL, Surjono A. The first 24-hour bilirubin level as a predictor of hyperbilirubinemia in healthy term newborns. Paediatrica Indonesiana. 2003; 43: 85-90.

21. Ohishi A, Kondo M, Fujita T, Baba T, Iijima S. Accuracy of transcutaneous bilirubin level measured by a JM-105 bilirubinometer. Pediatr Neonatol. 2023 Jan;64(1):32-37.

22. Cordero N, Petrova A, Halari A, Hegyi T. Transcutaneous bilirubin measurements in preterm infants: the impact of race, age, and phototherapy. J Perinatol. 2026 Jan 5.