


DAFTAR PUSTAKA

- Abdelfattah, H.E., Bekhet, M.M.M., Tawfik, F.A., Elias, D.G., & Saleh, A.M.M.A.E.H. (2025). The association between insulin resistance and risk of developing depression and anxiety disorders in a sample of Egyptian population. *Egypt. J. Intern. Med.* 37 : 56.
- Aman, M., Resnawita, D., Rasyid, H., Kasim, H., Bakri, S., Umar, H., *et al.* (2021). The concordance of triglyceride glucose index (TyG index) and homeostatic model assessment for insulin resistance (Homa-IR) in non-diabetic subjects of adult Indonesian males. *Clin. Epidemiol. Glob. Health* 9 : 227–230.
- Ashwell, M., Gunn, P., & Gibson, S. (2012). Waist-to-height ratio is a better screening tool than waist circumference and BMI for adult cardiometabolic risk factors: systematic review and meta-analysis. *Obes. Rev.* 13 : 275–286.
- Baek, J.H., Kim, H., Kim, K.Y., & Jung, J. (2018). Insulin Resistance and the Risk of Diabetes and Dysglycemia in Korean General Adult Population. *Diabetes Metab. J.* 42 : 296.
- Ballena-Caicedo, J., Zuzunaga-Montoya, F.E., Loayza-Castro, J.A., Bustamante-Rodríguez, J.C., Vásquez Romero, L.E.M., Tapia-Limonchi, R., *et al.* (2025). Global prevalence of insulin resistance in the adult population: a systematic review and meta-analysis. *Front. Endocrinol.* 16 : 1646258.
- Bazyar, H., Zare Javid, A., Masoudi, M.R., Haidari, F., Heidari, Z., Hajializadeh, S., *et al.* (2024). Assessing the predictive value of insulin resistance indices for metabolic syndrome risk in type 2 diabetes mellitus patients. *Sci. Rep.* 14 : 8917.
- Bonora, E., Targher, G., Alberiche, M., Bonadonna, R.C., Saggiani, F., Zenere, M.B., *et al.* (2000). Homeostasis model assessment closely mirrors the glucose clamp technique in the assessment of insulin sensitivity: studies in subjects with various degrees of glucose tolerance and insulin sensitivity. *Diabetes Care* 23 : 57–63.
- Bowman, C.F., & Nichols, J.H. (2020). Comparison of Accuracy Guidelines for Hospital Glucose Meters. *J. Diabetes Sci. Technol.* 14 : 546–552.
- Cohen, J.F., Korevaar, D.A., Altman, D.G., Bruns, D.E., Gatsonis, C.A., Hooft, L., *et al.* (2016). STARD 2015 guidelines for reporting diagnostic accuracy studies: explanation and elaboration. *BMJ Open* 6 : e012799.
- Couto, A.N., Pohl, H.H., Bauer, M.E., & Schwanke, C.H.A. (2023). Accuracy of the triglyceride-glucose index as a surrogate marker for identifying metabolic syndrome in non-diabetic individuals. *Nutrition* 109 : 111978.
- DeFronzo, R.A. (2009). From the Triumvirate to the Ominous Octet: A New Paradigm for the Treatment of Type 2 Diabetes Mellitus. *Diabetes* 58 : 773–795.
- DeFronzo, R.A., Ferrannini, E., Groop, L., Henry, R.R., Herman, W.H., Holst, J.J., *et al.* (2015). Type 2 diabetes mellitus. *Nat. Rev. Dis. Primer* 1 : 15019.
- Dou, J., Guo, F., Guo, C., Wu, R.-Y., Hu, Y.-W., Sun, X.-L., *et al.* (2025). Triglyceride glucose-waist height ratio index is a promising and robust glucose and lipid metabolism indicator for predicting long-term type 2

- diabetes: insights from a Japanese prospective cohort data from 12 years. *Diabetes Res. Clin. Pract.* 227 : 112398.
- Du, M.-F., Zhang, X., Hu, G.-L., Mu, J.-J., Chu, C., Liao, Y.-Y., *et al.* (2023). Associations of lipid accumulation product, visceral adiposity index, and triglyceride-glucose index with subclinical organ damage in healthy Chinese adults. *Front. Endocrinol.* 14 : 1164592.
- Du, T., Yuan, G., Zhang, M., Zhou, X., Sun, X., & Yu, X. (2014). Clinical usefulness of lipid ratios, visceral adiposity indicators, and the triglycerides and glucose index as risk markers of insulin resistance. *Cardiovasc. Diabetol.* 13 : 146.
- Elkanawati, R., Sumiwi, S., & Levita, J. (2024). Impact of Lipids on Insulin Resistance: Insights from Human and Animal Studies. *Drug Des. Devel. Ther.* Volume 18 : 3337–3360.
- Er, L.-K., Wu, S., Chou, H.-H., Hsu, L.-A., Teng, M.-S., Sun, Y.-C., *et al.* (2016). Triglyceride Glucose-Body Mass Index Is a Simple and Clinically Useful Surrogate Marker for Insulin Resistance in Nondiabetic Individuals. *PLOS ONE* 11 : e0149731.
- Erqou, S., Adler, A.I., Challa, A.A., Fonarow, G.C., & Echouffo-Tcheugui, J.B. (2022). Insulin resistance and incident heart failure: a META-ANALYSIS. *Eur. J. Heart Fail.* 24 : 1139–1141.
- Gast, K.B., Tjeerdema, N., Stijnen, T., Smit, J.W.A., & Dekkers, O.M. (2012). Insulin Resistance and Risk of Incident Cardiovascular Events in Adults without Diabetes: Meta-Analysis. *PLoS ONE* 7 : e52036.
- Gastaldelli, A. (2022). Measuring and estimating insulin resistance in clinical and research settings. *Obesity* 30 : 1549–1563.
- Ghani, Z.A., College of Pharmacy, AL Bayan University, Baghdad, Iraq, Qaddori, H., Department of Physiology, College of Medicine, Al Nahrain University, Baghdad, Iraq, Al-Mayah, Q., & Department of Physiology, College of Medicine, Al Nahrain University, Baghdad, Iraq (2023). Triglyceride/high-density lipoprotein ratio as a predictor for insulin resistance in a sample of healthy Iraqi adults. *J. Med. Life* 16 : 668–674.
- Goh, L.P.W., Sani, S.A., Sabullah, M.K., & Gansau, J.A. (2022). The Prevalence of Insulin Resistance in Malaysia and Indonesia: An Updated Systematic Review and Meta-Analysis. *Medicina (Mex.)* 58 : 826.
- Gounden, V., Devaraj, S., & Jialal, I. (2024). The role of the triglyceride-glucose index as a biomarker of cardio-metabolic syndromes. *Lipids Health Dis.* 23 : 416.
- Gutch, M., Kumar, S., Razi, S., Gupta, K., & Gupta, A. (2015). Assessment of insulin sensitivity/resistance. *Indian J. Endocrinol. Metab.* 19 : 160.
- Jamar, G., Almeida, F.R.D., Gagliardi, A., Sobral, M.R., Ping, C.T., Sperandio, E., *et al.* (2017). Evaluation of waist-to-height ratio as a predictor of insulin resistance in non-diabetic obese individuals. A cross-sectional study. *Sao Paulo Med. J.* 135 : 462–468.
- Kahn, S.E., Hull, R.L., & Utzschneider, K.M. (2006). Mechanisms linking obesity to insulin resistance and type 2 diabetes. *Nature* 444 : 840–846.



Monsen, G., Jacobsen, C.E., & SKUP, (Scandinavian Evaluation of Laboratory Equipment for Primary Health Care) (2009). Contour® meter and test strips

- designed for glucose self-measurement and measurements by health care professionals manufactured by Bayer HealthCare: report from an evaluation organised by SKUP (No. SKUP/2009/75). Bergen, Norway : SKUP.
- Moriyama, K. (2025). Mini-review on insulin resistance assessment: Advances in surrogate indices and clinical applications. *World J. Clin. Cases* 13.
- Nakamura, M., Iso, H., Kitamura, A., Imano, H., Noda, H., Kiyama, M., *et al.* (2016). Comparison between the triglycerides standardization of routine methods used in Japan and the chromotropic acid reference measurement procedure used by the CDC Lipid Standardization Programme. *Ann. Clin. Biochem. Int. J. Lab. Med.* 53 : 632–639.
- Owei, I., Umekwe, N., Provo, C., Wan, J., & Dagogo-Jack, S. (2017). Insulin-sensitive and insulin-resistant obese and non-obese phenotypes: role in prediction of incident pre-diabetes in a longitudinal biracial cohort. *BMJ Open Diabetes Res. Care* 5 : e000415.
- Paramanathan, T., Sandrasegarampillai, B., Arasaratnam, V., & Thirunavukarasu, K. (2024). The discriminative ability of the triglyceride-glucose index to identify metabolic syndrome among adults of the northern Sri Lankan population. *BMC Endocr. Disord.* 24 : 101.
- Parcha, V., Heindl, B., Kalra, R., Li, P., Gower, B., Arora, G., *et al.* (2022). Insulin Resistance and Cardiometabolic Risk Profile Among Nondiabetic American Young Adults: Insights From NHANES. *J. Clin. Endocrinol. Metab.* 107 : e25–e37.
- Petersen, M.C., & Shulman, G.I. (2018). Mechanisms of Insulin Action and Insulin Resistance. *Physiol. Rev.* 98 : 2133–2223.
- Piqueras, P., Ballester, A., Durá-Gil, J.V., Martinez-Hervas, S., Redón, J., & Real, J.T. (2021). Anthropometric Indicators as a Tool for Diagnosis of Obesity and Other Health Risk Factors: A Literature Review. *Front. Psychol.* 12 : 631179.
- Pliego-Sandoval, J.E., Díaz-Barbosa, A., Reyes-Nava, L.A., Angeles Camacho-Ruiz, M., Iñiguez-Muñoz, L.E., & Pinto-Pérez, O. (2023). Development and Evaluation of a Low-Cost Triglyceride Quantification Enzymatic Biosensor Using an Arduino-Based Microfluidic System. *Biosensors* 13 : 826.
- Reaven, G.M. (2005). THE INSULIN RESISTANCE SYNDROME: Definition and Dietary Approaches to Treatment. *Annu. Rev. Nutr.* 25 : 391–406.
- Rohlfing, C., Petroski, G., Hatten-Beck, M., Hanson, S., Hoofnagle, A.N., Little, R.R., *et al.* (2025). The current status of serum insulin measurements and the need for standardization. *Clin. Chem. Lab. Med. CCLM* 63 : 2442–2446.
- Rosli, N., Kwon, H., Lim, J., Yoon, Y.A., & Jeong, J. (2022). Measurement comparability of insulin assays using conventional immunoassay kits. *J. Clin. Lab. Anal.* 36 : e24521.
- Samuel, V.T., & Shulman, G.I. (2016). The pathogenesis of insulin resistance: integrating signaling pathways and substrate flux. *J. Clin. Invest.* 126 : 12–22.
- Samuel, V.T., & Shulman, G.I. (2012). Mechanisms for Insulin Resistance: Common Threads and Missing Links. *Cell* 148 : 852–871.

- Sari, Y.R., Jesica, F., & Niken, N. (2023). Comparison between Pooled Sera and Commercial Serum on the Accuracy of Triglyceride Assessment. *Int. J. Multidiscip. Approach Res. Sci.* 1 : 375–380.
- Simmons, E., Johnson, D., Delhay, R., Colwell, J., Clor, E., Santa, P., *et al.* (2019). Development and validation of a noncompetitive electrochemiluminescence-based immunoassay (ECLIA) for specific determination of insulin lispro (Humalog®) in human serum to support pharmacokinetic assessments. *J. Immunoassay Immunochem.* 40 : 314–327.
- Sonagra AD, Zubair M, & Motiani A (2024). Hexokinase Method., in: *StatPearls*. Treasure Island (FL) : Treasure Island (FL): StatPearls Publishing.
- Song, S., Son, D.-H., Baik, S.-J., Cho, W.-J., & Lee, Y.-J. (2022). Triglyceride Glucose-Waist Circumference (TyG-WC) Is a Reliable Marker to Predict Non-Alcoholic Fatty Liver Disease. *Biomedicines* 10 : 2251.
- Sun, Y., Ji, H., Sun, W., An, X., & Lian, F. (2025). Triglyceride glucose (TyG) index: A promising biomarker for diagnosis and treatment of different diseases. *Eur. J. Intern. Med.* 131 : 3–14.
- Susilawati, M.D., Muljati, S., & Bantas, K. (2015). Perbandingan IMT dan Indikator Obesitas Sentral terhadap Kejadian Diabetes Melitus Tipe 2 (DMT2). *Bul. Penelit. Kesehat.* 43.
- Tárraga Marcos, P.J., López-González, Á.A., Martínez-Almoyna Rifá, E., Paublini Oliveira, H., Martorell Sánchez, C., Tárraga López, P.J., *et al.* (2025a). Risk of Insulin Resistance in 44,939 Spanish Healthcare Workers: Association with Sociodemographic Variables and Healthy Habits. *Diseases* 13 : 33.
- Tárraga Marcos, P.J., López-González, Á.A., Martínez-Almoyna Rifá, E., Paublini Oliveira, H., Martorell Sánchez, C., Tárraga López, P.J., *et al.* (2025b). Risk of Insulin Resistance in 44,939 Spanish Healthcare Workers: Association with Sociodemographic Variables and Healthy Habits. *Diseases* 13 : 33.
- Tosoratto, J., Tárraga López, P.J., López-González, Á.A., Busquets-Cortes, C., Obrador De Hevia, J., & Ramirez-Manent, J.I. (2025). Associations Between Shift Work and Insulin Resistance Markers in 53,053 Spanish Workers: A Sex-Stratified Cross-Sectional Analysis Using TyG, TyG-BMI, METS-IR, and SPISE-IR Indices. *J. Clin. Med.* 14 : 4604.
- Van Dijk, S.B., Takken, T., Prinsen, E.C., & Wittink, H. (2012). Different anthropometric adiposity measures and their association with cardiovascular disease risk factors: a meta-analysis. *Neth. Heart J.* 20 : 208–218.
- Vijayalakshmi, P., Manjula, H., & Usha, S. (2023). Association of Lipid Accumulation Product with Insulin Resistance in Type 2 Diabetes Mellitus: A Cross-sectional Study. *J. Clin. Diagn. Res.*
- Wakabayashi, I., & Daimon, T. (2014). A Strong Association between Lipid Accumulation Product and Diabetes Mellitus in Japanese Women and Men. *J. Atheroscler. Thromb.* 21 : 282–288.
- WHO (2004). Appropriate body-mass index for Asian populations and its implications for policy and intervention strategies. *The Lancet* 363 : 157–163.

- Wulandari, S.R., Permatasari, L., Dewi, A.S., Ruella, N., & Utami, W. (2024). Review: Metode-Metode Pemeriksaan Glukosa Darah. *Benzena Pharm. Sci. J.* 3 : 85–95.
- Yalcin, H., Toprak, B., & Colak, A. (2017). The independent relationship between hemoglobin A_{1c} and homeostasis model assessment of insulin resistance in non-diabetic subjects. *Turk. J. Biochem.* 42 : 31–36.
- Yamada, C., Mitsuhashi, T., Hiratsuka, N., Inabe, F., Araida, N., & Takahashi, E. (2011). Optimal reference interval for homeostasis model assessment of insulin resistance in a Japanese population: HOMA-IR reference interval in Japanese. *J. Diabetes Investig.* 2 : 373–376.
- Zatterale, F., Longo, M., Naderi, J., Raciti, G.A., Desiderio, A., Miele, C., *et al.* (2020). Chronic Adipose Tissue Inflammation Linking Obesity to Insulin Resistance and Type 2 Diabetes. *Front. Physiol.* 10 : 1607.
- Zheng, S., Shi, S., Ren, X., Han, T., Li, Y., Chen, Y., *et al.* (2016). Triglyceride glucose-waist circumference, a novel and effective predictor of diabetes in first-degree relatives of type 2 diabetes patients: cross-sectional and prospective cohort study. *J. Transl. Med.* 14 : 260.
- Zheng, Y., Yin, G., Chen, F., Lin, L., & Chen, Y. (2022). Evaluation of Triglyceride Glucose Index and Homeostasis Model of Insulin Resistance in Patients with Polycystic Ovary Syndrome. *Int. J. Womens Health* Volume 14 : 1821–1829.