Pemanfaatan SPUKTA Berbasis Citra RGB untuk Mengamati Keragaman Klon-Klon Teh (Camellia sinensis L.)
di Kebun Binorong PT Pagilaran
Miranda Cahayaningtyas, Dr. Panjisakti Basunanda, S.P., M.P.; Taufan Alam, S.P., M.Sc.
Universitas Gadjah Mada, 2026 | Diunduh dari http://etd.repository.ugm.ac.id/

DAFTAR PUSTAKA

Abdul, R., Nida, M., Xiongming, D. U., & Tehseen, A. M. (2020). Heritability and correlation analysis of morphological and yield traits in genetically modified cotton. *Journal of Cotton Research*, *3*(23), 1–9. https://doi.org/https://doi.org/10.1186/s42397-020-00067-z

Alves, A. K. S., Araújo, M. S., Chaves, S. F. S., Dias, L. A. S., Corrêdo, L. P., Pessoa, G. G. F. A., & Bezerra, A. R. G. (2024). High throughput phenotyping in soybean breeding using RGB image vegetation indices based on drone. *Scientific Reports*, *14*(32055), 1–11. https://doi.org/https://doi.org/10.1038/s41598-024-83807-4 1

An, Y., Mi, X., Zhao, S., Guo, R., Xia, X., & Liu, S. (2020). Revealing distinctions in genetic diversity and adaptive evolution between two varieties of camellia sinensis by whole-genome resequencing. *Frontiers in Plant Science*, *11*, 1–14. https://doi.org/10.3389/fpls.2020.603819

Anasiru, R. H., Hartono, H., Wicaksono, A., Saryoko, A., Fatmawati, A. A., & Bahrun, A. H. (2026). Pemantauan kehijauan daun padi sawah dengan drone RGB dan algoritma VARI : Validasi indeks SPAD. *Teknotan*, *20*(1), 137–146. https://doi.org/10.24198/jt.vol20n1.15

Anjarsari, I. R. D., Hamdani, J. S., Suherman, C., Nurmala, T., Khomaeni, H. S., Budidaya, D., Pertanian, F., & Padjadjaran, U. (2021). Studi Pemangkasan dan Aplikasi Sitokinin-Giberelin pada Tanaman Teh (Camellia sinensis (L .) O . Kuntze) Produktif Klon GMB 7 Study of Pruning and Cytokinin-Giberelin Applications on Productive Tea Plant. *Jurnal Agronomi Indonesia*, *49*(1), 89–96. https://doi.org/doi.org/10.24831/jai.v49i1.32046

Arkorful, E., Yu, Y., Chen, C., Lu, L., Hu, S., Yu, H., & Li, X. (2020). Untargeted metabolomic analysis using UPLC-MS / MS identi fi es metabolites involved in shoot growth and development in pruned tea plants (Camellia sinensis (L.) O. Kuntz). *Scientia Horticulturae*, *264*, 1–12. https://doi.org/10.1016/j.scienta.2019.109164

Badan Pusat Statistika. (2025). Badan Pusat Statistik, Jakarta. Indonesia.

Balasuriya, J. (1999). Shoot population density and shoot weight of clonal tea (Camellia sinensis) at di ff erent altitudes in Sri Lanka. *Europian Journal of Agronomy*, *11*, 123–130. https://doi.org/doi:10.1016/s1161-0301(99)00024-6

Bendig, J., Yu, K., Aasen, H., Bolten, A., Bennertz, S., Broscheit, J., Gnyp, M. L., & Bareth, G. (2015). Combining UAV-based plant height from crop surface models , visible , and near infrared vegetation indices for biomass monitoring in barley. *International Journal of Applied Earth Observation and Geoinformation*, *39*, 79–87. https://doi.org/10.1016/j.jag.2015.02.012

Benti, T., Debela, A., Bekele, Y., & Suleman, S. (2023). Effect of seasonal variation on yield and leaf quality of tea clone (Camellia sinensis (L.) O. Kuntze) in South West Ethiopia.

UNIVERSITAS GADJAH MADA
Pemanfaatan SPUKTA Berbasis Citra RGB untuk Mengamati Keragaman Klon-Klon Teh (Camellia sinensis L.) di Kebun Binorong PT Pagilaran
Miranda Cahayaningtyas, Dr. Panjisakti Basunanda, S.P., M.P.; Taufan Alam, S.P., M.Sc.
Universitas Gadjah Mada, 2026 | Diunduh dari http://etd.repository.ugm.ac.id/

Heliyon, *9*, 1–9. https://doi.org/10.1016/j.heliyon.2023.e14051

Budiherlando, A., & Taryono. (2019). Pendugaan parameter genetik beberapa komponen hasil calon galur teh keturunan kedua hasil perbanyakan vegetatif. *Vegetalika*, *8*(3), 153–165. https://doi.org/https://doi.org/10.22146/veg.42027

Buntaran, H., Maria, A., Vasquez, B., Gordillo, A., Sahr, M., Wimmer, V., & Piepho, H. P. (2022). Assessing the response to genomic selection by simulation. *Theoretical and Applied Genetics*, *135*, 2891–2905. https://doi.org/10.1007/s00122-022-04157-1

Carvalho, L. C., Gonçalves, E. F., Marques da Silva, J., & Costa, J. M. (2021). Potential phenotyping methodologies to assess inter- and intravarietal variability and to select grapevine genotypes tolerant to abiotic stress. *Frontiers in Plant Science*, *12*, 1–16. https://doi.org/10.3389/fpls.2021.718202

Chen, F., Liu, H., Zeng, Z., Zhou, X., & Tan, X. (2022). BES-Net : Boundary enhancing semantic context network for high-resolution image semantic segmentation. *Remote Sensing*, *14*, 1–21. https://doi.org/https://doi.org/10.3390/rs14071638

Eng, L. S., Ismail, R., Hashim, W., & Baharum, A. (2019). The use of VARI, GLI, and VIGREEN formulas in detecting vegetation in aerial images. *International Journal of Technology*, *10*(7), 1385–1394. https://doi.org/DOI:10.14716/ijtech.v10i7.3275

Esan, E. B., & Omolaja, S. S. (2002). Genotypic association, path analysis and pluck quality value in tea [Camellia sinensis (L.) O. Kuntze]. *Tropical Agriculture*, *79*(2), 1–13. https://journals.sta.uwi.edu/ojs/index.php/ta/article/view/1366

Fatawa, M. I., Santosa, E., Hapsari, D. P., & Krisantini. (2023). Changing rainfall and its adaptation strategies on tea plantation in West Java , Indonesia. *Indonesian Journal of Agronomy*, *51*(2), 257–268. https://doi.org/https://dx.doi.org/10.24831/ija.v51i2.47081 Edited

Hanafi, A., Syahrian, H., Rahadi, V. P., & Prayoga, M. K. (2025). Analisis karakter pertumbuhan 16 klon teh tipe Assamica (Camellia sinensis var. Assamica) pada fase pembibitan. *Jurnal Sains Teh Dan Kina*, *4*(1), 1–13. https://doi.org/10.22302/pptk.jur.jstk.v4i1.201

Irwin, H., Fadhilah, N., & Amalia, A. F. (2020). Teknik penggunaan drone dengan sensor kamera RGB dan algoritma VARI untuk mengidentifikasi tingkat stres tanaman jagung. *Buletin Teknik Pertanian*, *25*(2), 85–88. https://doi.org/https://doi.org/10.24198/jt.vol20n1.15

Khan, M. N., Tan, Y., He, L., & Dong, W. (2025). From air to space : A comprehensive approach to optimizing aboveground biomass estimation on UAV-based datasets. *Forest*, *16*(214), 1–25. https://doi.org/https://doi.org/10.3390/f16020214

Kirana, P. L., Mitrowihardjo, S., & Murti, R. H. (2015). Stabilitas hasil pucuk tujuh klon harapan teh (Camellia sinensis (L.) O. Kuntze) di Kebun Kayulandak. *Vegetalika*, *4*(3), 98–111. https://doi.org/https://doi.org/10.22146/veg.10481

UNIVERSITAS GADJAH MADA
Pemanfaatan SPUKTA Berbasis Citra RGB untuk Mengamati Keragaman Klon-Klon Teh (Camellia sinensis L.) di Kebun Binorong PT Pagilaran
Miranda Cahayaningtyas, Dr. Panjisakti Basunanda, S.P., M.P.; Taufan Alam, S.P., M.Sc.
Universitas Gadjah Mada, 2026 | Diunduh dari http://etd.repository.ugm.ac.id/

Krause, M. R., Mondal, S., Haghighattalab, A., Poland, J., Gore, M. A., & Sorrells, M. E. (2020). Aerial high-throughput phenotyping enables indirect selection for grain yield at the early generation , seed-limited stages in breeding programs. *Crop Science*, *60*, 3096–3114. https://doi.org/10.1002/csc2.20259

Kumar, P., Chaudhary, S., Sood, S., Mahant, S., Pundir, S., & Kalia, P. (2025). Tea beyond the cup: A comprehensive review of Camellia sinensis (L.) O. Kuntze. *Clinical Phytoscience*, *11*(23), 1–39. https://doi.org/https://doi.org/10.1186/s40816-025-00413-1

Li, H., Wang, Y., Fan, K., Mao, Y., Shen, Y., & Ding, Z. (2022). Evaluation of important phenotypic parameters of tea plantations using multi-source remote sensing data. *Frontiers in Plant Science*, *13*, 1–19. https://doi.org/10.3389/fpls.2022.898962

Li, J., Li, H., Liu, Z., Wang, Y., Chen, Y., Yang, N., Hu, Z., Li, T., & Zhuang, J. (2023). Molecular markers in tea plant (Camellia sinensis): Applications to evolution, genetic identification, and molecular breeding. *Plant Physiology and Biochemistry*, *198*(107704), 1–14. https://doi.org/10.1016/j.plaphy.2023.107704

Li, J., Wu, W., Zhao, C., Bai, X., Dong, L., & Tan, Y. (2025). Effects of solar elevation angle on the visible light vegetation index of a cotton field when extracted from the UAV. *Scientific Reports*, *15*(18497), 1–24. https://doi.org/https://doi.org/10.1038/s41598-025-00992-6 2

Li, Y., Yang, X., Tong, L., Wang, L., & Clarke, J. (2023). Phenomic selection in slash pine multi-temporally using UAV-multispectral imagery. *Frontiers in Plant Science*, *14*, 1–13. https://doi.org/10.3389/fpls.2023.1156430

Liu, X., Wu, X., Peng, Y., Mo, J., & Fang, S. (2023). Application of UAV-retrieved canopy spectra for remote evaluation of rice full heading date. *Science of Remote Sensing*, *7*, 1–11. https://doi.org/10.1016/j.srs.2023.100090

Lopez-cruz, M., Olson, E., Rovere, G., Crossa, J., Dreisigacker, S., Mondal, S., Singh, R., & Campos, G. D. L. (2020). Regularized selection indices for breeding value prediction using hyper-spectral image data. *Scientific Reports*, *10*, 1–12. https://doi.org/10.1038/s41598-020-65011-2

Lubanga, N., Bančič, J., Massawe, F., Mayes, S., & Gorjanc, G. (2023). Genomic selection strategies to increase genetic gain in tea breeding programs. *The Plant Genomme*, *16*, 1–18. https://doi.org/10.1002/tpg2.20282

Meng, X., Zhu, H., Yan, H., Wang, D., Yang, C., & Zhang, Y. (2018). Supplementary data contents page. *Journal of Agricultural and Food Chemistry*, *66*(27), 7150–7155. https://doi.org/https://doi.org/10.1021/acs.jafc.8b02066

Meyer, G. E., & Neto, J. C. (2008). Verification of color vegetation indices for automated crop imaging applications. *Elsevier*, *63*, 282–293. https://doi.org/10.1016/j.compag.2008.03.009

UNIVERSITAS GADJAH MADA
Pemanfaatan SPUKTA Berbasis Citra RGB untuk Mengamati Keragaman Klon-Klon Teh (Camellia sinensis L.) di Kebun Binorong PT Pagilaran
Miranda Cahayaningtyas, Dr. Panjisakti Basunanda, S.P., M.P.; Taufan Alam, S.P., M.Sc.
Universitas Gadjah Mada, 2026 | Diunduh dari http://etd.repository.ugm.ac.id/

Murti, R. H., Puspitasari, A., & Mitrowihardjo, S. (2014). Stability analysis of nine promising clones of tea (Camellia sinensis). *Agrivita*, *36*(1), 81–90. https://doi.org/http://dx.doi.org/10.17503/Agrivita-2014-36-1-p081-090

Mustamu, Y. A., Ustari, D., Wicaksono, A. A., Concibido, V., Suganda, T., & Karuniawan, A. (2024). Performance, stability analysis , and selection for new yellow-fleshed sweet potato (Ipomoea batatas [L.] Lam.) in West Java , Indonesia. *International Journal of Agronomy*, *2024*(1), 1–13. https://doi.org/10.1155/2024/6774420

Nardino, M., Pereira, J. M., Vinicius Torres Marques, Davila, F. C., Franco, F. D., & Barros, W. S. (2020). COEFFICIENT OF VARIATION : A NEW APPROACH FOR THE STUDY 1 Introduction. *Brazilian Journal of Biometrics*, *38*(2), 185–206. https://doi.org/10.28951/rbb.v38i2.440

Owuor, P. O., Kamau, D. M., & Jondiko, E. O. (2009). Responses of clonal tea to location of production and plucking intervals. *Food Chemistry*, *115*(1), 290–296. https://doi.org/10.1016/j.foodchem.2008.11.073

Rahadi, V. P., Khomaeni, H. S., & Sriyadi, B. (2016). Pengujian daya tumbuh setek klon teh (Camellia sinensis) hasil persilangan klon-klon generasi pertama. *Jurnal Pemuliaan Teh Dan Kina*, *19*(2), 124–130. https://doi.org/https://doi.org/10.22302/pptk.jur.jptk.v19i2.118

Rahimi, M., Mordrostami, M., & Mortezavi, M. (2019). Evaluation of tea (Camellia sinensis L .) biochemical traits in normal and drought stress conditions to identify drought tolerant clones. *Physiology and Molecular Biology of Plants*, *25*(1), 59–69. https://doi.org/10.1007/s12298-018-0564-x

Rahman, A., Tarwaca, E., Putra, S., & Waluyo, S. (2018). The effect of pyraclostrobin application on yield of tea (Camelia sinensis (L.) O. Kuntze) under wet season. *Agriculture Science*, *3*(3), 153–159. https://doi.org/https://doi.org/10.22146/ipas.26428

Rebetzke, G. J., Fischer, R. A., Deery, D. M., & Smith, D. J. (2019). Review : High-throughput phenotyping to enhance the use of crop genetic. *Plant Science*, *282*, 40–48. https://doi.org/10.1016/j.plantsci.2018.06.017

Sayuti, S., Maulidi, A., & Khaerudin, A. (2022). Pengembangan prototipe pesawat udara tanpa awak jenis fixed wing VTOL. *Jurnal Rekayasa Energi Dan Mekanika*, *2*(2), 128–138. https://doi.org/10.26760/jrem.v2i2.128

Sun, J., Rutkoski, J. E., Poland, J. A., Crossa, J., Jannink, J., & Sorrells, M. E. (2017). Multitrait, random regression , or simple repeatability model in high-throughput phenotyping data improve genomic prediction for wheat grain yield. *The Plant Genome*, *10*(2), 1–12. https://doi.org/10.3835/plantgenome2016.11.0111

Tanaka, T. S. T., Wang, S., Jørgensen, J. R., Gentili, M., Vidal, A. Z., Mortensen, A. K., Acharya, B. S., Beck, B. D., & Gislum, R. (2024). Review of crop phenotyping in field plot experiments using UAV-mounted sensors and algorithms. *Drones*, *8*(212), 1–21. https://doi.org/https://doi.org/10.3390/ drones8060212

UNIVERSITAS GADJAH MADA
Pemanfaatan SPUKTA Berbasis Citra RGB untuk Mengamati Keragaman Klon-Klon Teh (Camellia sinensis L.) di Kebun Binorong PT Pagilaran
Miranda Cahayaningtyas, Dr. Panjisakti Basunanda, S.P., M.P.; Taufan Alam, S.P., M.Sc.
Universitas Gadjah Mada, 2026 | Diunduh dari http://etd.repository.ugm.ac.id/

Tesfa, M., Tena, E., & Kebede, M. (2026). Analysis of variance components, heritability, and genetic advance for agronomic and biochemical traits in sugarcane genotypes. *Advances in Agriculture*, *2026*, 1–12. https://doi.org/10.1155/aia/8331924

Wang, S., Yu, C., Ma, J., Ouyang, J., & Zhao, Z. (2024). Tea yield estimation using UAV images and deep learning. *Industrial Crops & Products*, *212*(118358), 1–14. https://doi.org/10.1016/j.indcrop.2024.118358

Wang, X., Feng, H., Chang, Y., Ma, C., Wang, L., Hao, X., Li, A., Cheng, H., Wang, L., Cui, P., Jin, J., Wang, X., Wei, K., Ai, C., Zhao, S., Wu, Z., Li, Y., Liu, B., Wang, G., … Yang, Y. (2020). Population sequencing enhances understanding of tea plant evolution. *Nature Communications*, *11*, 1–10. https://doi.org/10.1038/s41467-020-18228-8

Wang, Y., Yang, Z., Kootstra, G., & Khan, H. A. (2023). The impact of variable illumination on vegetation indices and evaluation of illumination correction methods on chlorophyll content estimation using UAV imagery. *Plant Methods*, *19*(1), 1–16. https://doi.org/10.1186/s13007-023-01028-8

Wientjes, Y. C. J., Bijma, P., Calus, M. P. L., Zwaan, B. J., & Vitezica, Z. G. (2022). The long - term effects of genomic selection: 1 . Response to selection , additive genetic variance , and genetic architecture. *Genetics Selection Evolution*, *54*(19), 1–21. https://doi.org/10.1186/s12711-022-00709-7

Wijayanto, A., Indradewa, D., & Putra, E. T. S. (2015). Kuantitas dan kualitas hasil pucuk enam klon teh sinensis (Camellia sinensis (L.) O. Kuntze var Sinensis) di bagian Kebun Kayulandak, PT Pagilran. *Vegetalika*, *4*(3), 42–56. https://doi.org/https://doi.org/10.22146/veg.10476

Winursito, Suryadi, M., & Waluyo, S. (2012). Hasil dan keragaman genetik tujuh klon teh (Camellia sinensis (L.) Kuntze) di dua lokasi dengan ketinggian berbeda. *Vegetalika*, *1*(4), 1–13. https://doi.org/https://doi.org/10.22146/veg.1605

Xie, C., & Yang, C. (2020). A Review on plant high-throughput phenotyping traits using UAV-based sensors. *Computers and Electronics in Agriculture*, *178*, 1–14. https://doi.org/10.1016/j.compag.2020.105731

Xie, X., Sun, Z., Zhang, X., & Han, X. (2023). Novel aspects of regulation of nitrogen responses in the tea plant (Camellia sinensis (L.)). *Agronomy*, *13*(144), 1–14. https://doi.org/https:// doi.org/10.3390/agronomy13010144

Xu, R., Li, C., & Bernardes, S. (2021). Development and testing of a UAV-based multi-sensor system for plant phenotyping and precision agriculture. *Remote Sensing*, *13*(3517), 1–28. https://doi.org/https://doi.org/10.3390/rs13173517

Yang, X., Chen, J., Lu, X., Liu, H., Liu, Y., Bai, X., Qian, L., & Zhang, Z. (2025). Advances in UAV remote sensing for monitoring crop water and nutrient status: Modeling methods, influencing factors, and challenges. *Plants*, *14*(2544), 1–30. https://doi.org/https://doi.org/10.3390/plants14162544

UNIVERSITAS
GADJAH MADA
**Pemanfaatan SPUKTA Berbasis Citra RGB untuk Mengamati Keragaman Klon-Klon Teh (Camellia sinensis L.)
di Kebun Binorong PT Pagilaran**
Miranda Cahayaningtyas, Dr. Panjisakti Basunanda, S.P., M.P.; Taufan Alam, S.P., M.Sc.
Universitas Gadjah Mada, 2026 | Diunduh dari http://etd.repository.ugm.ac.id/

Yilmaz, G., Kandernir, N., & Kinalioglu, K. (2004). Effects of different pruning intervals on fresh shoot yield and some quality properties of tea (Camellia sinensis (L.) O. Kuntze) in Turkey. *Pakistan Journal of Biological Science*, *7*(7), 1208–1212. https://doi.org/10.3923/pjbs.2004.1208.1212

Yuan, Y., Bayer, P. E., Batley, J., & Edwards, D. (2021). Current status of structural variation studies in plants. *Plant Biotechnology Journal*, *19*, 2153–2163. https://doi.org/10.1111/pbi.13646

Zhang, C., Marzougui, A., & Sankaran, S. (2020). High-resolution satellite imagery applications in crop phenotyping: An overview. *Computers and Electronics in Agriculture*, *175*(105584), 1–10. https://doi.org/10.1016/j.compag.2020.105584

Zhang, Q., Wan, B., Cao, Z., Zhang, Q., & Wang, D. (2021). Exploring the potential of unmanned aerial vehicle (UAV) remote sensing for mapping plucking area of tea plantations. *Forest*, *12*(1214), 1–22. https://doi.org/https:// doi.org/10.3390/f12091214

Zhang, X., Chen, S., Shi, L., Gong, D., Zhang, S., Zhao, Q., Zhan, D., Vasseur, L., Wang, Y., Yu, J., Liao, Z., Xu, X., Qi, R., Wang, W., Ma, Y., Wang, P., Ye, N., Ma, D., Shi, Y., … Tang, H. (2021). Haplotype-resolved genome assembly provides insights into evolutionary history of the tea plant Camellia sinensis. *Nature Genetics*, *53*, 1250–1259. https://doi.org/10.1038/s41588-021-00895-y

Zhang, Y., Jiang, Y., Xu, B., Yang, G., Feng, H., Yang, X., Yang, H., Liu, C., Cheng, Z., & Feng, Z. (2024). Study on the estimation of leaf area index in rice based on. *Remote Sensing*, *16*(3049), 1–24. https://doi.org/https://doi.org/10.3390/rs16163049

Zhuang, Z., Tsai, H., & Chen, C.-I. (2025). Estimating Tea Plant Physiological Parameters Using Unmanned Aerial Vehicle Imagery and Machine Learning Algorithms. *Sensors*, *25*(1966), 1–28. https://doi.org/https:// doi.org/10.3390/s25071966

Ziyomo, C., & Bernardo, R. (2013). Drought tolerance in maize : Indirect selection through secondary traits versus genomewide selection. *Crop Science*, *53*, 1269–1275. https://doi.org/10.2135/cropsci2012.11.0651