UNIVERSITAS GADJAH MADA
Perbandingan Efek Takrolimus Dengan Triamsinolon Asetonida Terhadap Proliferasi Fibroblas Keloid Manusia
Lintang Unggul Rini, dr. Arief Budiyanto, Ph.D.Sp.KK, dr. Niken Trisnowati, M.Sc.Sp.KK
Universitas Gadjah Mada, 2026 | Diunduh dari http://etd.repository.ugm.ac.id/

DAFTAR PUSTAKA

Allen A, Siegfried E, Silverman R, Williams ML, Elias PM, Szabo SK, Korman NJ.Significant Absorption of Topical Tacrolimus in 3 Patients With Netherton Syndrome.*Arch Dermatol.2001;137:747-750*

Almatsier M, Azwar A, Djuanda A, Firmansyah R, Handaya, Ismail S, Muchtar A, Sani A, Setiabudy R.2007.MIMS Annual Full Prescribing Information(Eds.).Jakarta:CMPMedica

Bayat A, Arscott G, Ollier WE, et al.Description of site-specific morphology of keloid phenotypes in an Afrocaribbean population. *Br J Plast Surg 2004, 57:122-33*

Berman B, Flores F.The treatment of hypertrophic scars and keloids.Eur J Dermatol.1998 Dec;8(8):591-5

Bergman J, Rico MJ.Tacrolimus clinical studies for atopic dermatitis and other conditions.*Semin Cutan Med Surg. 2001 Dec;20(4):250-9*

Braun-Falco O, Plewig G, Wolff HH, Winkelman RK. 1984.Dermatology.Germany:Springer-Verlag

Brissett AE, Sherris DA.Scar contractures, hypertrophic scars, and keloids. *Facial Plast Surg 2001, 17:263-72*

Carroll LA, Hanasono MM, Mikulec AA, Kita M, Koch RJ. Triamcinolone stimulates bFGF production and inhibits TGF-β1 production by human dermal fibroblass. *Dermatol Surg 2002 Aug; 28(8):704-9*

Dennler S, André J, Alexaki I, Li A, Magnaldo T, ten Dijke P, Wang XJ, Verrecchia F, Mauviel A. Induction of sonic hedgehog mediators by transforming growth factor-beta: Smad3-dependent activation of Gli2 and Gli1 expression in vitro and in vivo.*Cancer Res. 2007 Jul 15;67(14):6981-6.*

Diegelmann RF, Cohen IK, McCoy BJ.Growth kinetics and collagen synthesis of normal skin, normal scar and keloid fibroblasts in vitro.*J Cell Physiol. 1979 Feb;98(2):341-6*

UNIVERSITAS GADJAH MADA
Perbandingan Efek Takrolimus Dengan Triamsinolon Asetonida Terhadap Proliferasi Fibroblas Keloid Manusia
Lintang Unggul Rini, dr. Arief Budiyanto, Ph.D.Sp.KK, dr. Niken Trisnowati, M.Sc.Sp.KK
Universitas Gadjah Mada, 2026 | Diunduh dari http://etd.repository.ugm.ac.id/

Duong HS, Zhang Q, Kobi A, Le A, Messadi DV.Assesment of Morphological and Immunohistological Alterations in Long-Term Keloid Skin Explants. *Cells Tissues Organs.2005;181:89-102*

Fine A, Goldstein RH.The effect of transforming growth factor-beta on cell proliferation and collagen formation by lung fibroblasts.*J Biol Chem. 1987 Mar 15;262(8):3897-902*

Freshney RI.2005.Culture of Animal Cells, a Manual of Basic Technique, 5th Ed. Hoboken NJ, John Wiley & Sons

Gabbiani G, Hirschel BJ, Ryan GB, Statkov PR, Majno G. Granulation tissue as a contractile organ.A study of structure and function.*J Exp Med.1972;135:719-34*

Gagliano N, Moscheni C, Tartaglia GM, Selleri S, Chiriva-Internati M, Cobos E, Torri C, Costa F, Pettinari L, Gioia M.A therapeutic dose of FK506 does not affect collagen turnover pathways in healthy human gingival fibroblasts.*Transplant Proc. 2008 Jun;40(5):1419-24.*

Gao Z, Wang Z, Shi Y, Lin Z, Jiang H, Hou T, Wang Q, Yuan X, Zhao Y, Wu H.Modulation of Collagen Synthesis in Keloid Fibroblasts by Silencing Smad2 With siRNA.*Youxin JinPlastic and Reconstructive Surgery 2006;118(6):1328-37*

Giordano A, Romano S, Mallardo M, D'Angelillo A, Calì G, Corcione N, Ferraro P, Romano MF. FK506 can activate transforming growth factor-β signalling in vascular smooth muscle cells and promote proliferation.*Oxford Journals Medicine Cardiovascular Research 2008 Volume79, Issue3 Pp. 519-526.*

Haimowitz JE, Margolis DJ.Moist wound healing, in Krasner D, Kane D (eds): Chronic Wound Care: A Clinical Source Book for Healthcare Professionals. Wayne, PA, Health *Management Publications, 1997, 49-55*

Hanifin JM, Ling MR, Langley R, Breneman D, Rafal E.Tacrolimus ointment for the treatment of atopic dermatitis in adult patients: part I, efficacy.*J Am Acad Dermatol.2001 Jan;44(1Suppl):S28-38*

Hamrick M, Boswell W, Carney D.Successful treatment of earlobe keloids in the pediatric population. *J. Pediatr. Surg. 2009 Jan; 44(1):286-8*

UNIVERSITAS GADJAH MADA
Perbandingan Efek Takrolimus Dengan Triamsinolon Asetonida Terhadap Proliferasi Fibroblas Keloid Manusia
Lintang Unggul Rini, dr. Arief Budiyanto, Ph.D.Sp.KK, dr. Niken Trisnowati, M.Sc.Sp.KK
Universitas Gadjah Mada, 2026 | Diunduh dari http://etd.repository.ugm.ac.id/

Howling GI, Dettmar PW, Goddard PA, Hampson FC, Dornish M, Wood EJ. The effect of chitin and chitosan on fibroblast populated collagen lattice contraction. *Biotechnology Application Biochemistry. 2002;36: 247-253*

Kane D.2001.Chronic wound healing and chronic wound management, in Krasner D, Rodeheaver GT, Sibbald RG. (eds): Chronic Wound Care: A Clinical Source Book for Healthcare Professionals, Third Edition. Wayne, PA, Health Management Publications, pp 7-17.

Katzung B.G,Trevor A.J,Masters,S.B.2008.Katsung and Trevor's Pharmacology Examination and Board Review Eighth Edition. USA:Mc Graw Hill

Keast DH, Orsted H.The basic principle of wound care.*Ostomy Wound Manage.1998 Aug;44(8):24-8,30-1*

Kelly AP.Medical and surgical therapies for keloids. *Dermatol Ther 2004, 17:212-8*

Kerstein MD.Introduction: moist wound healing.*American Journal of Surgery 1994;167(1A Suppl): 1S-6S*

Kim A, DiCarlo J, Cohen C, McCall C, Johnson D, McAlpine B, Quinn AG, McLaughlin ER, Arbiser JL. Are keloids really "gli-loids"?: High-level expression of *gli*-1 oncogene in keloids.
J Am Acad Dermatol. 2001 Nov;45(5):707-11

Kino T, Hatanaka H, Miyata S, Inamura N, Nishiyama M, Yajima T, Goto T, Okuhara M, Kohsaka M, Aoki H, et al.FK-506, a novel immunosuppressant isolated from a Streptomyces. II. Immunosuppressive effect of FK-506 in vitro.*J Antibiot(Tokyo). 1987 Sep;40(9):1256-65.*

Kischer CW, Hendrix MJ.Fibronectin (FN) in hypertrophic scars and keloids.*Cell Tissue Res. 1983;231(1):29-37*

Kuhns DB, Young HA, Gallin EK, Gallin JI. Ca2+-dependent production and release of IL-8 in human neutrophils. *J Immunol 1998; 161: 4332-9.*

MacLeod J (ed).1981.Davidson's Principles and Practice of Medicine, Thirteenth Edition. Edinburgh UK, pp 590-592

Marks JG, Miller JJ. 2006. Lookingbill and Marks' Principle of Dermatology Fourth Edition. USA : Elseiver

UNIVERSITAS GADJAH MADA
Perbandingan Efek Takrolimus Dengan Triamsinolon Asetonida Terhadap Proliferasi Fibroblas Keloid Manusia
Lintang Unggul Rini, dr. Arief Budiyanto, Ph.D.Sp.KK, dr. Niken Trisnowati, M.Sc.Sp.KK
Universitas Gadjah Mada, 2026 | Diunduh dari http://etd.repository.ugm.ac.id/

Marsland AM, Griffiths CE.The macrolide immunosuppressants in dermatology: mechanisms of action.*European Journal of Dermatology. 2002 12(6): 618-22.*

Messadi DV, Le A, Berg S, et al.Effect of TGF-beta 1 on PDGF receptors expression in human scar fibroblasts. *Front Biosci 1998, 3:a16-22*

McCance LK, Huether SE, Brashers VL, Rote NS.2010. Pathophysiology: The Biologic Basis for Disease in Adults and Children SixthEd. Philadelphia:Elseiver

Muroi N, Nishibori M, Fujii T, Yamagata M, Hosoi S, Nakaya N, Saeki K, Henmi K.Anaphylaxis From The Carboxymethylcellulose Component of Barium Sulfate Suspension. *The New England Journal of Medicine. 1997.Volume 337.*

Nath AK, D'Souza M.Localized Immediate Type Hypersensitivity Reaction Following Intralesional Triamcinolone Acetonide into a Keloid.*Egyptian Dermatology Online Journal 2010;6(1):16*

Newsome E, Bolling R, Langston K, Wang A, Jansen D.Wound Healing, Keloids. *Transplant Proc. 1990 February ; 22(1): 52-56.*

Nghiem P, Pearson G, Langley RG.Tacrolimus and pimecrolimus: from clever prokaryotes to inhibiting calcineurin and treating atopic dermatitis. *J Am Acad Dermatol 2002; 46:228-41.*

Postlethwaite AE, Keski-Oja J, Moses HL, Kang AH. Stimulation of the chemotactic migration of human fibroblasts by transforming growth factor beta.*J Exp Med. 1987 Jan 1;165(1):251-6*

Quele-Roussel, correspondence. Topical tacrolimus and pimecrolimus are not assosiated with skin atrophy.Reply from autho:*British Journal of Dermatology 2002;146:343*

Raghow R, Postlethwaite AE, Keski-Oja J, Moses HL, Kang AH.Transforming growth factor-beta increases steady state levels of type I procollagen and fibronectin messenger RNAs posttranscriptionally in cultured human dermal fibroblasts.*J Clin Invest. 1987 Apr;79(4):1285-8*

UNIVERSITAS GADJAH MADA
Perbandingan Efek Takrolimus Dengan Triamsinolon Asetonida Terhadap Proliferasi Fibroblas Keloid Manusia
Lintang Unggul Rini, dr. Arief Budiyanto, Ph.D.Sp.KK, dr. Niken Trisnowati, M.Sc.Sp.KK
Universitas Gadjah Mada, 2026 | Diunduh dari http://etd.repository.ugm.ac.id/

Reitamo S, Rissanen J, Remitz A, Granlund H, Erkko P, Elg P, Autio P, Lauerma AI.Tacrolimus ointment does not affect collagen synthesis: results of a single-center randomized trial.*J Invest Dermatol 1998; 111: 396-8*

Romano G, Cavarape A, Favret G, Bortolotti N, Bartoli E.Systemic and intratubular effect of cyclosporin-A and tacrolimus on the rat kidney.*European Journal of Pharmacology 2000;309(215-221)*

Santoso,S.2010.Statistik Nonparametrik.Elex Media.

Sastroasmoro S, Ismael S.1995.Dasar-Dasar Metodologi Penelitian Klinis.Jakarta : Binarupa Aksara

Satish L, Lyons-Weiler J, Hebda PA, Weils A.Gene expression patterns in isolated keloid fibroblast.*Wound Repair Regen 2006 Jul-Aug;14(4): 463-70*

Schäffer M, Fuchs N, Völker J, Schulz T, Kapischke M, Viebahn R.Differential effect of tacrolimus on dermal and intestinal wound healing.*J Invest Surg. 2005 Mar-Apr;18(2):71-9.*

Schreiber SL, Crabtree GR. The mechanism of action of cyclosporin A and FK506.*Immunol Today 1992; 13: 136-42.*

Sestini R, Pacini A, Di Gennaro P, Bacci S, Pinzani P, CesatiV, Carli P, Massi D, Genuardi M, Romagnoli P.Tacrolimus reduces the expression of GLI-1 by basal cell carcinoma.*Journal of Investigative Dermatology, 2004 Nov:123(5)A87-A97(11).*

Singer M, Salpeter M.M. 1961. Regeneration in vertebrates: the role of the wound epithelium. *In* Growth in Living Systems.Basic Books, Inc., New York. 277.

Slodownik D, Ingber A.The use of tacrolimus in dermatology. *Harefuah. 2003 Nov;142(11):770-4, 805*

Sperr WR, Agis H, Czerwenka K, Virgolini I, Bankl HC, Muller MR, Zsebo K, Lechner K, Valent P. Effects of cyclosporin A and FK-506 on stem cell factor-induced histamine secretion and growth of human mast cells. *J Allergy Clin Immunol 1996; 98:389-99.*

UNIVERSITAS GADJAH MADA
Perbandingan Efek Takrolimus Dengan Triamsinolon Asetonida Terhadap Proliferasi Fibroblas Keloid Manusia
Lintang Unggul Rini, dr. Arief Budiyanto, Ph.D.Sp.KK, dr. Niken Trisnowati, M.Sc.Sp.KK
Universitas Gadjah Mada, 2026 | Diunduh dari http://etd.repository.ugm.ac.id/

Sullivan STO, Shaughnessy MO, O'Connor TPF.1996. Aetiology and management of hypertrophic scars and keloids.*Ann R Coll Surg Engl 1996; 78: 168-175*

Sun T, Adra S,Smallwood R, Holcombe M,MacNeil S, Exploring Hypotheses of the Actions of TGF-b1 in Epidermal Wound Healing Using a 3D Computational Multiscale Model of the Human Epidermis. *PLoS ONE 2009;4(12): e8515*

Van der Windt AE, Jahr H, Farrell E, Verhaar JAN, Weinans H, Van Osch GJVM.Calcineurin inhibitors promote chondrogenic marker expression of dedifferentiated human adult chondrocytes via stimulation of TGFß1 production (doi: 10.1089/ten.TEA.2009.0082)

Vistica DT, Skehan P, Scudiero D, Monks A, Pittman A, Boyd MR.Tetrazolium-based assays for cellular viability: a critical examination of selected parameters affecting formazan production.*Cancer Res. 1991;51, 2515-2520*

Wahl LM, Wahl SM.1992.Inflammation, in Cohen IK, Diegelman RF, Lindblad WJ (eds): Wound Healing: Biochemical and Clinical Aspects. Philadelphia, PA, W.B. Saunders, pp 40-62

Wirohadidjojo YW, Etnawati K, Adiwinarni DR.The combination of suprakeloidal flap and pulsed light heat energy in keloid management: A case report.*Berkala Ilmu Kedokteran Vol. 40, No. 1, Maret 2008: 45-49*

Yao D, Dore Jr JJ, Leof EB.FKBP12 is a negative regulator of transforming groeth factor-beta receptor internalization,*J Biol Chem 275 (2000),pp.13149-13154*

Zhang JG, Walmsley MW, Moy JV, Cunningham AC, Talbot D, Dark JH, Kirby JA.Differential effects of cyclosporin A and tacrolimus on the production of TGF-beta: implications for the development of obliterative bronchiolitis after lung transplantation.*Transpl Int. 1998;11 Suppl 1:S325-7.*