

## DAFTAR PUSTAKA

- Angus, D., Van, D., 2013. *Severe Sepsis and Septic Shock*. *N Eng J Med* ; 369: 840- 851
- Bersten, A. D., Soni, N., 2014. *Oh's Intensive Care Manual*. 7<sup>th</sup> Edition. Adelaide : Sepsis ; 717-723
- Biro, G.P., 2013. *From the Atmosphere to the Mitochondria : The Oxygen Cascade; From : Hemoglobin-Based Oxygen Carriers as Red Cell Substitutes and Oxygen Therapeutics*. ISBN 978-3-642-40716-1: 27-53
- Cain, S.M., 1983. *Peripheral Oxygen Uptake And Delivery In Health And Disease*. Rockville. *Clinical Chest Medicine*: 4:139.
- Collis, C.S., Rice, E.C., 1996. *Free radicals in hypoxia and reoxygenation*. Tissue oxygen deprivation. New york : Cambridge University Press; p: 275-296.
- Correa, T.D., Vuda, M., Takala, J., Djafarzadeh, S., Silva, E., Jakob, SM., 2013. *Increasing mean arterial blood pressure in sepsis:effects on fluid balance, vasopressor load and renal function*. *Critical Care Medicine*; 17 : R21.
- Dellinger, R.P., Levy, M.M., 2013. *Surviving Campaign : International Guidelines For Management Of Severe Sepsis And Septic Shock 2012*. *Critical Care Medicine*; 41: 580-637.
- Dubin, A., Mario, O.P., Christian, A.C., Fernando, P., Gaston, M., Mirim, C., Vanina, S., Elisa, E., Can, I.2009.*Increasing arterial blood pressure with norepinephrine does not improve microcirculatory blood flow: a prospective study*.*Critical care*; 13:R92.
- Duranteau, J., Chandel, N.S., Kulish, A, Shao, Z, Schumacker, P.T., 1998. *Intracellular signaling by reactive oxygen species during hypoxia in cardiomyocytes*. *J Biol Chem*, 273..
- Elixhauser, A., Friedman, B., Stranges, E., 2009. *Septicemia in US Hospitals*. Agency Healthcare, Rockville. <http://www.hcupus.ahrq.gov/reports/statbriefs/sb12.pdf> (diakses pada 22 Mei 2016 20.00 wib)

- Gutierrez, G., Palizas, F., Doglio, G., 1992. *gastric Intramucosal Ph As A Therapeutic Index Of Tissue Oxygenation In Critically Ill Patients*. Lancet. 339:195.
- Hollenberg, S.M., Ahrens, T.S., Annane, D., 2004. *practice Parameters For Hemodynamic Support Of Sepsis In Adult Patients: update*. 32:1928.
- Hurtado, F.J., Buroni, M., Tenzi, J., 2009. Sepsis: *Clinical Approach, Evidence-Based at the Bedside*. In: gallo a, et al, editors. Intensive and Critical Care Medicine. Springer-Verlag Italia. 299-309.
- International Guidelines for Management of Severe Sepsis and Septic Shock. Surviving Sepsis Campaign. 2015. <https://www.sccm.org/SSC-Guidelines.pdf>. (diunduh pada tanggal 14 November 2016 pukul 06.00 WIB).
- Jing, Y., Si, Q., Chun, P., Hong, L., Shi, X., Shu, L., Ai, R., Ling, L., Ying, Z., Feng, M., Yi, Y., Hai, B. 2015. *A high mean arterial pressure target is associated with improved microcirculation in septic shock patients with previous hypertension : a prospective open label study*. Critical care; 19:130
- John, M.A., Michael, M.G., 2013. Resuscitation fluids. *New Engl J of Med*; 369:13
- Jones, D.P., 1996. *Cellular Energetics And Biochemistry Of Hypoxia*. In: Haddad GG, Lister G (eds). *Tissue oxygen deprivation*. Marcel Dekker: New york,: 25-50.
- Josef, F., Sugiman, T., Tampubolon, O., Rupi'i., 2014. *Penatalaksanaan sepsis dan syok septik*. Surviving sepsis campagne bundle. Perdici.
- Kula, R., Chyle, V., Szturz, P., Sklienka, P., Sukenik, Tichy, Maca, Neiser., 2009. *A Response To Infection In Patients With Severe Sepsis – Do We Need a “Stage-Directed Therapy Concept?”*. Bratisl Lek Listy : 459-464
- Leach, R., 2014. *Critical Care Medicine At A Glance*. 3<sup>rd</sup> edition. John wiley & Son. Chapter 25
- LeDoux, Astiz, M.E., Carpati, Rackrow., 2002. *Effects Of Perfusion Pressure On Tissue Perfusion In Septic Shock*. Critical Care Medicine : 2729-32

- Leone, M., Pierre, A., Peter, R., Jean, L., Claude, M.2015.*Optimizing mean arterial pressure in septic shock: a critical reappraisal of the literature*. Critical care:19:101
- Marino, P.L., 1990. *The ICU Book 2nd Edition. Oxygen and Carbondioxide Transport*. William & Wilkins; 31-51.
- Mervyn, S., Clifford, S., Christopher, W., Manu, S.,2016.*The third internasional consensus definitions for sepsis and syok septic*. JAMA vol 315.801-810.
- Muller, L., Toumi, M., Bousquet, P.J., 2011. *An Increasing In Aortic Blood Flow After An Infusion Of 100ml Colloid Over 1 Minute Can Predict Fluid Responsiveness : the mini-fluid challenge study*. Anesthesiology; 115(3): 541-7
- McLellan, S.A., and Walsh, T.S., 2004. Oxygen Delivery and Haemoglobin. *British Journal of Anesthesia Volume 4 Number 4*. 23-126
- Nebout, S., Pirracchio, R., 2012. Review Article : *Should we Monitor ScVO2 in Critically Ill Patients?* *Cardiologi Research and Practice*.1-7 Retrieved from: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC3177360/>. (diakses pada tanggal 19 Desember 2016 pukul 17.00 wib).
- Nguyen, B.,Rivers, E., Abrahamian, F., Moran, G., Abraham, E.2006. *Severe sepsis and septic shock: Review of the literature and emergency. Department management guidelines*. *Annals of Emergency Medicine*. 48; 28-54
- Peake S., Delaney, A., Bailey .,Bellomo. R., Cameron, P.A., 2014. *goal-Directed Resuscitation For Patients With Early Septic Shock*. *NEJM*; 371:1496.
- Pierre, A., Ferhat, Bruno, Conrad., 2014. *High Versus Low Blood-Pressure Target in Patients with septic shock*. *New England Journal Medicine* : 370:1583-1593
- Poeze, M., Solberg, B.C., Greve, J.W., Ramsay, G., 2005. *Monitoring Global Volume-Related Hemodynamic Or Regional Variables After Initial Resuscitation: What Is A Better Predictor Of Outcome In Critically Ill Septic Patients?*. *Critical Care Medicine* ; 33:2494.

- Rivers, E., Nguyen, B., Havstad, S., 2001. *Early Goal-Directed Therapy In The Treatment Of Severe Sepsis And Septic Shock*. *N Engl J Med* ; 345:1368.
- Sastroasmoro, A., Ismael, S., 1995. *Dasar-Dasar Metodologi Penelitian Klinis*. Binarupa Aksara.; 52-77
- Sedarmayanti, Hidayat, S., 2002. *Metodologi Penelitian. Mandar Maju Bandung*.; 144-146
- Sessler, C.N., Perry, J.C., Varney, K.L. 2004. *management Of Severe Sepsis And Septic Shock*. *Current Critical Care*; 10:354.
- Textoris, J., Fouche, L., Wiramus, S., Antonini, F., Tho, S., Martin, C., and Leone, M., 2011. *High Central Venous Oxygen Saturation In The Latter Stages ofSeptic Shock is Associated With Increase Mortality*.*Critical Care Medicine Journal*. 1-6. Retrieved from: <http://ccforum.com/content/15/4/R 176>
- Vallet, B., 1998. *Vascular Reactivity And Tissue Oxygenation*. *Intensive Care Med*; 24: 3.
- Van der Poll, T., Opal, S. 2008. *Host-Pathogen Interaction In Sepsis*. *Lancet Infect*: 8:32-43
- Vincent, J.L. 2008. *Clinical Sepsis And Septic Shock-Definition, Diagnosis And Management Principles*. *Langenbecks Arch Surg*; 393:817-24
- Yealy, D.M., Kellum, J.A. 2014. *a Randomized Trial Of Protocol-Based Care For Early Septic Shock*. *N Engl J Med*; 370:1683.