

DAFTAR PUSTAKA

- [1] C. A. W. Glas, “Missing Data A2 - Peterson, Penelope,” E. Baker and B. B. T.-I. E. of E. (Third E. McGaw, Eds. Oxford: Elsevier, 2010, pp. 283–288.
- [2] L. G. Dmitrović, V. Dušak, and J. Dobša, “Missing data problems in non-Gaussian probability distributions,” *Informatologia*, vol. 49, no. 3–4, pp. 138–152, 2016.
- [3] J. L. Schafer, “Multiple imputation: a primer,” *Stat Methods Med Res*, vol. 8, no. 1, pp. 3–15, 1999.
- [4] K. I. Vaden, M. Gebregziabher, S. E. Kuchinsky, and M. A. Eckert, “Multiple imputation of missing fMRI data in whole brain analysis,” *Neuroimage*, vol. 60, no. 3, pp. 1843–1855, 2012.
- [5] I. Eekhout, H. C. W. De Vet, J. W. R. Twisk, J. P. L. Brand, M. R. De Boer, and M. W. Heymans, “Missing data in a multi-item instrument were best handled by multiple imputation at the item score level,” *J. Clin. Epidemiol.*, vol. 67, no. 3, pp. 335–342, 2014.
- [6] E.-L. Silva-Ramírez, R. Pino-Mejías, and M. López-Coello, “Single imputation with multilayer perceptron and multiple imputation combining multilayer perceptron and k-nearest neighbours for monotone patterns,” *Appl. Soft Comput. J.*, vol. 29, pp. 65–74, 2015.
- [7] P. and S. H. Berglund, *Multiple Imputation of Missing Data Using SAS*. SAS Institute Inc., Cary, North California, USA, 2014.
- [8] P. Royston, “Multiple imputation of missing values,” *Stata J.*, vol. 4, no. 3, pp. 227–241, 2004.
- [9] X. Yang, G. Da Su, J. Chen, and Y. S. Moon, “Restoration of low resolution car plate images using PCA based image super-resolution,” in *Proceedings - International Conference on Image Processing, ICIP*, 2010, pp. 2789–2792.
- [10] C. Pous, D. Caballero, and B. Lopez, “Diagnosing Patients Combining

- Principal Components Analysis and Case Based Reasoning,” in *2008 Eighth International Conference on Hybrid Intelligent Systems*, 2008, pp. 819–824.
- [11] C.-F. Tsai and F.-Y. Chang, “Combining instance selection for better missing value imputation,” *J. Syst. Softw.*, vol. 122, pp. 63–71, Dec. 2016.
- [12] J. Geronimi and G. Saporta, “Variable selection for multiply-imputed data with penalized generalized estimating equations,” *Comput. Stat. Data Anal.*, vol. 110, pp. 103–114, 2017.
- [13] X. Yang, T. R. Belin, and W. J. Boscardin, “Imputation and variable selection in linear regression models with missing covariates,” *Biometrics*, vol. 61, no. 2, 2005.
- [14] A. R. T. Donders, G. J. M. G. van der Heijden, T. Stijnen, and K. G. M. Moons, “Review: A gentle introduction to imputation of missing values,” *J. Clin. Epidemiol.*, vol. 59, no. 10, pp. 1087–1091, 2006.
- [15] Z. Zhang, “Missing data imputation: focusing on single imputation.,” *Ann. Transl. Med.*, vol. 4, no. 1, p. 9, 2016.
- [16] M. P. Gómez-Carracedo, J. M. Andrade, P. López-Mahía, S. Muniategui, and D. Prada, “A practical comparison of single and multiple imputation methods to handle complex missing data in air quality datasets,” *Chemom. Intell. Lab. Syst.*, vol. 134, pp. 23–33, 2014.
- [17] M. N. Karim, C. M. Reid, L. Tran, A. Cochrane, and B. Billah, “Missing Value Imputation Improves Mortality Risk Prediction Following Cardiac Surgery: An Investigation of an Australian Patient Cohort,” *Hear. Lung Circ.*, vol. 26, no. 3, pp. 301–308, 2017.
- [18] J. Kaiser, “Dealing with Missing Values in Data,” *J. Syst. Integr.*, pp. 42–51, 2014.
- [19] A. C. Acock, “Working with missing values,” *J. Marriage Fam.*, vol. 67, no. 4, pp. 1012–1028, 2005.

- [20] D. Little, RJA and Rublin, “Statistical Analysis with Missing Data,” *Wiley, New York.*, p. 381, 1987.
- [21] S. G. Karadogan, L. Marchegiani, L. Kai Hansen, and J. Larsen, *How efficient is estimation with missing data?* 2011.
- [22] B. A. Bell, J. D. Kromrey, and J. M. Ferron, “Missing Data and Complex Samples: The Impact of Listwise Deletion vs. Subpopulation Analysis on Statistical Bias and Hypothesis Test Results When Data Are MCAR and MAR,” in *Proceedings of the Joint Statistical Meetings, Survey Research Methods Section*, 2009, pp. 4759–4770.
- [23] H. W. Marsh, “Pairwise deletion for missing data in structural equation models: Nonpositive definite matrices, parameter estimates, goodness of fit, and adjusted sample sizes,” *Struct. Equ. Model.*, vol. 5, no. 1, pp. 22–36, 1998.
- [24] S. F. Nielsen, “Nonparametric conditional mean imputation,” *J. Stat. Plan. Inference*, vol. 99, no. 2, pp. 129–150, 2001.
- [25] A. Gelman and J. Hill, “Missing-data imputation,” in *Data Analysis Using Regression and Multilevel/Hierarchical Models*, 2007, pp. 529–543.
- [26] S. B. Woolley, A. A. Cardoni, and J. W. Goethe, “Last-Observation-Carried-Forward Imputation Method in Clinical Efficacy Trials: Review of 352 Antidepressant Studies,” *Pharmacother. J. Hum. Pharmacol. Drug Ther.*, vol. 29, no. 12, pp. 1408–1416, 2009.
- [27] and I. S. M. Jatra, R. R. Isnanto, “Identifikasi Iris Mata Menggunakan Metode Analisis Komponen Utama Dan Perhitungan Jarak Euclidean,” Universitas Diponegoro, 2011.
- [28] R. J. and D. Wichern, *Applied Multivariate Statistical Analysis*, Keenam. Prentice Hall, 2007.