



DAFTAR PUSTAKA

- Adamis, A. P. (2002) 'Is diabetic retinopathy an inflammatory disease?', *The British journal of ophthalmology*. BMJ Publishing Group, 86(4), pp. 363–5. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11914197> (Accessed: 13 January 2018).
- Afari, M. E. and Bhat, T. (2016) 'Neutrophil to lymphocyte ratio (NLR) and cardiovascular diseases: An update', *Expert Review of Cardiovascular Therapy*, 14(5), pp. 573–577. doi: 10.1586/14779072.2016.1154788.
- Ahmed, A. M. (2002) 'History of diabetes mellitus.', *Saudi medical journal*, 23(4), pp. 373-8. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11953758> (Accessed: 3 September 2017).
- Al-Rubeaan, K., Abu El-Asrar, A. M., Youssef, A. M., Subhani, S. N., Ahmad, N. A., Al-Sharqawi, A. H., Alguwaihes, A., Alotaibi, M. S., et al. (2015) 'Diabetic retinopathy and its risk factors in a society with a type 2 diabetes epidemic: A Saudi National Diabetes Registry-based study', *Acta Ophthalmologica*, 93(2), pp. e140–e147. doi: 10.1111/aos.12532.
- Al-Sarraf, A. A., Al-Bannai, S. K., Al-Furaih, A. M. and El-Shaziy, M. K. (2010) 'Prevalence and Factors Associated With Diabetic Retinopathy, a Multi-Centric Study in Kuwait', *Alexandria Bulletin*, (2).
- Alberti, K. G. M. M. and Zimmet, P. Z. (1998) 'Definition, diagnosis and classification of diabetes mellitus and its complications. Part 1: diagnosis and classification of diabetes mellitus. Provisional report of a WHO Consultation', *Diabetic Medicine*, 15(7), pp. 539–553. doi: 10.1002/(SICI)1096-9136(199807)15:7<539::AID-DIA668>3.0.CO;2-S.
- Alrawahi, A. H., Rizvi, S. G. A., Al-Riyami, D. and Al-Anqoodi, Z. (2012) 'Prevalence and Risk Factors of Diabetic Nephropathy in Omani Type 2 Diabetics in Al-Dakhiliyah Region', *Oman Medical Journal*, 27(3), pp. 212–216. doi: 10.5001/omj.2012.48.
- AlSaraj, F. (2015) 'Pathogenesis of Type 2 Diabetes Mellitus', in *Treatment of Type 2 Diabetes*. InTech. doi: 10.5772/59183.
- American Diabetes Association (2002) 'Diabetic Retinopathy', *Diabetes Care*. American Diabetes Association, 25(Supplement 1), pp. S90–S93. doi: 10.2337/diacare.25.2007.S90.
- American Diabetes Association (2002) 'Implications of the United Kingdom Prospective Diabetes Study', *Diabetes Care*, 25(Supplement 1), pp.S28-S32.
- American Diabetes Association (2004) 'Diagnosis and Classification of Diabetes Mellitus', *Diabetes Care*, 27(Supplement 1), pp. S5–S10. doi: 10.2337/diacare.27.2007.S5



- American Diabetes Association (2014) ‘Diagnosis and classification of diabetes mellitus.’, *Diabetes care*. American Diabetes Association, 37 Suppl 1(Supplement 1), pp. S81-90. doi: 10.2337/dc14-S081.
- American Diabetes Association (2017) ‘6. Glycemic Targets: Standards of Medical Care in Diabetes—2018’, *Diabetes Care*, 41(Supplement 1), pp. S55-S64.
- An, X., Mao, H.-P., Wei, X., Chen, J.-H., Yang, X., Li, Z.-B., Yu, X.-Q. and Li, Z.-J. (2012) ‘Elevated neutrophil to lymphocyte ratio predicts overall and cardiovascular mortality in maintenance peritoneal dialysis patients’, *International Urology and Nephrology*, 44(5), pp. 1521–1528. doi: 10.1007/s11255-012-0130-3.
- Andersohn, F., Konzen, C. and Garbe, E. (2007) ‘Systematic review: agranulocytosis induced by nonchemotherapy drugs.’, *Annals of internal medicine*, 146(9), pp. 657–65. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17470834> (Accessed: 2 April 2018).
- Araszkiewicz, A. and Zozulinska-Ziolkiewicz, D. (2016) ‘Retinal Neurodegeneration in the Course of Diabetes-Pathogenesis and Clinical Perspective.’, *Current neuropharmacology*. Bentham Science Publishers, 14(8), pp. 805–809. doi: 10.2174/1570159X14666160225154536.
- Azab, B., Camacho-Rivera, M. and Taioli, E. (2014) ‘Average values and racial differences of neutrophil lymphocyte ratio among a nationally representative sample of United States subjects.’, *PloS one*. Public Library of Science, 9(11), p. e112361. doi: 10.1371/journal.pone.0112361.
- Badan Penelitian dan Pengembangan Kesehatan (2013) ‘Riset Kesehatan Dasar (RISKESDAS) 2013’, *Laporan Nasional 2013*, pp. 1–384.
- Barber, A. J., Gardner, T. W. and Abcouwer, S. F. (2011) ‘The significance of vascular and neural apoptosis to the pathology of diabetic retinopathy.’, *Investigative ophthalmology & visual science*. Association for Research in Vision and Ophthalmology, 52(2), pp. 1156–63. doi: 10.1167/iovs.10-6293.
- Barg, S., Eliasson, L., Renström, E. and Rorsman, P. (2002) ‘A subset of 50 secretory granules in close contact with L-type Ca²⁺ channels accounts for first-phase insulin secretion in mouse beta-cells.’, *Diabetes*. American Diabetes Association, 51 Suppl 1(suppl 1), pp. S74-82. doi: 10.2337/DIABETES.51.2007.S74.
- Bhutta, H., Agha, R., Wong, J., Tang, T. Y., Wilson, Y. G. and Walsh, S. R. (2011) ‘Neutrophil-lymphocyte ratio predicts medium-term survival following elective major vascular surgery: A cross-sectional study’, *Vascular and Endovascular Surgery*, 45(3), pp. 227–231. doi: 10.1177/1538574410396590.
- Bierhaus, A. and Nawroth, P. P. (2009) ‘Multiple levels of regulation determine the role of the receptor for AGE (RAGE) as common soil in inflammation,



immune responses and diabetes mellitus and its complications', *Diabetologia*, 52(11), pp. 2251–2263. doi: 10.1007/s00125-009-1458-9.

Bilous, R. and Donnelly, R. (2010) *Handbook of Diabetes*. Oxford, UK: Wiley-Blackwell. doi: 10.1002/9781444391374.

Böni-Schnetzler, M., Thorne, J., Parnaud, G., Marselli, L., Ehses, J. A., Kerr-Conte, J., Pattou, F., Halban, P. A., et al. (2008) 'Increased interleukin (IL)-1 β messenger ribonucleic acid expression in β -cells of individuals with type 2 diabetes and regulation of IL-1 β in human islets by glucose and autostimulation', *Journal of Clinical Endocrinology and Metabolism*. The Endocrine Society, 93(10), pp. 4065–4074. doi: 10.1210/jc.2008-0396.

Bourne, R. R. A., Stevens, G. A., White, R. A., Smith, J. L., Flaxman, S. R., Price, H., Jonas, J. B., Keeffe, J., et al. (2013) 'Causes of vision loss worldwide, 1990–2010: A systematic analysis', *The Lancet Global Health*, 1(6), pp. 339–349. doi: 10.1016/S2214-109X(13)70113-X.

Boutens, L. and Stienstra, R. (2016) 'Adipose tissue macrophages: going off track during obesity.', *Diabetologia*. Springer, 59(5), pp. 879–94. doi: 10.1007/s00125-016-3904-9.

Bouzakri, K., Roques, M., Gual, P., Espinosa, S., Guebre-Egziabher, F., Riou, J.-P., Laville, M., Le Marchand-Brustel, Y., et al. (2003) 'Reduced activation of phosphatidylinositol-3 kinase and increased serine 636 phosphorylation of insulin receptor substrate-1 in primary culture of skeletal muscle cells from patients with type 2 diabetes.', *Diabetes*. American Diabetes Association, 52(6), pp. 1319–25. doi: 10.2337/DIABETES.52.6.1319.

Brass, D., McKay, P. and Scott, F. (2014) 'Investigating an incidental finding of lymphopenia.', *BMJ (Clinical research ed.)*. British Medical Journal Publishing Group, 348, p. g1721. doi: 10.1136/BMJ.G1721.

Burtis, C., Ashwood, E. and Bruns, D. (2012) 'Tietz Textbook of Clinical Chemistry and Molecular Diagnostics.' 5th ed. London: Elsevier Health Sciences.

Buyukkaya, E., Karakaş, M. F., Karakaş, E., Akçay, A. B., Tanboga, I. H., Kurt, M. and Sen, N. (2014) 'Correlation of Neutrophil to Lymphocyte Ratio With the Presence and Severity of Metabolic Syndrome', *Clinical and Applied Thrombosis/Hemostasis*, 20(2), pp. 159–163. doi: 10.1177/1076029612459675.

Caprná, M., Kubatka, P., Saxena, S., Valasková, J., Stefanicková, J., Kobyliak, N., Zulli, A. and Kružliak, P. (2017) 'The Impact of Hyperglycemia on VEGF Secretion in Retinal Endothelial Cells', *Folia Medica*, 59(2), pp. 183–189. doi: 10.1515/folmed-2017-0029.

Carvalho Vidigal, F. de, Guedes Cocate, P., Gonçalves Pereira, L. and Cássia Gonçalves Alfenas, R. de (2012) 'The role of hyperglycemia in the induction of oxidative stress and inflammatory process', *Nutrición Hospitalaria*.



- Sociedad Española de Nutrición Parenteral y Enteral (SENPE), 27(5), pp. 1391–1398. doi: 10.3305/nh.2012.27.5.5917.
- Center for Disease Control and Prevention (2016) *At A Glance 2016 Diabetes*.
- Cerf, M. E. (2013) ‘Beta cell dysfunction and insulin resistance.’, *Frontiers in endocrinology*. Frontiers Media SA, 4, p. 37. doi: 10.3389/fendo.2013.00037.
- Chatham, W. (2017) *Glucocorticoid effects on the immune system - UpToDate, UpToDate*. Available at: <https://www.uptodate.com/contents/glucocorticoid-effects-on-the-immune-system#H3739728875> (Accessed: 4 April 2018).
- Chen, L., Tuo, B. and Dong, H. (2016) ‘Regulation of Intestinal Glucose Absorption by Ion Channels and Transporters.’, *Nutrients*. Multidisciplinary Digital Publishing Institute (MDPI), 8(1). doi: 10.3390/nu8010043.
- Cheung, N., Mitchell, P. and Wong, T. Y. (2010) ‘Diabetic retinopathy’, *The Lancet*, 376(9735), pp. 124–136. doi: 10.1016/S0140-6736(09)62124-3.
- Cheung, N. and Wong, T. Y. (2008) ‘Diabetic retinopathy and systemic vascular complications’, *Progress in Retinal and Eye Research*, 27(2), pp. 161–176. doi: 10.1016/j.preteyeres.2007.12.001.
- Chibber, R., Ben-Mahmud, B. M., Chibber, S. and Kohner, E. M. (2007) ‘Leukocytes in diabetic retinopathy.’, *Current diabetes reviews*, 3(1), pp. 3–14. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18220651> (Accessed: 19 August 2017).
- Cho, H. and Sobrin, L. (2014) ‘Genetics of diabetic retinopathy.’, *Current diabetes reports*. NIH Public Access, 14(8), p. 515. doi: 10.1007/s11892-014-0515-z.
- Chung, F.-M., Tsai, J. C.-R., Chang, D.-M., Shin, S.-J. and Lee, Y.-J. (2005) ‘Peripheral total and differential leukocyte count in diabetic nephropathy: the relationship of plasma leptin to leukocytosis.’, *Diabetes care*. American Diabetes Association, 28(7), pp. 1710–7. doi: 10.2337/DIACARE.28.7.1710.
- Ciray, H., Aksoy, A., Ulu, N., Cizmecioglu, A., Gaipov, A., and Solak, Y. (2015) ‘Nephropathy, but not Angiographically Proven Retinopathy, is Associated with Neutrophil to Lymphocyte Ratio in Patients with Type 2 Diabetes.’, *Experimental and Clinical Endocrinology & Diabetes*, 123(05), 267–271. <http://doi.org/10.1055/s-0035-1547257>
- Cnop, M., Foufelle, F. and Velloso, L. A. (2012) ‘Endoplasmic reticulum stress, obesity and diabetes’, *Trends in Molecular Medicine*, 18(1), pp. 59–68. doi: 10.1016/j.molmed.2011.07.010.
- Crawley, J. B., Rawlinson, L., Lali, F. V, Page, T. H., Saklatvala, J. and Foxwell, B. M. (1997) ‘T cell proliferation in response to interleukins 2 and 7 requires p38MAP kinase activation.’, *The Journal of biological chemistry*. American Society for Biochemistry and Molecular Biology, 272(23), pp. 15023–7. doi:



10.1074/JBC.272.23.15023.

- Cruz, N. G., Sousa, L. P., Sousa, M. O., Pietrani, N. T., Fernandes, A. P. and Gomes, K. B. (2013) 'The linkage between inflammation and Type 2 diabetes mellitus', *Diabetes Research and Clinical Practice*, 99(2), pp. 85–92. doi: 10.1016/j.diabres.2012.09.003.
- Curtis, B. R. (2014) 'Drug-induced immune neutropenia/agranulocytosis.', *Immunochemistry*, 30(2), pp. 95–101. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/25247619> (Accessed: 2 April 2018).
- Cybulska-Heinrich, A. K., Baertschi, M., Loesche, C. C., Schoetzau, A., Konieczka, K., Josifova, T. and Flammer, J. (2015) 'Patients with diabetic retinopathy have high retinal venous pressure.', *The EPMA journal*. Springer, 6(1), p. 5. doi: 10.1186/s13167-015-0027-1.
- Davids, M. S. (2018) *Approach to the adult with lymphocytosis or lymphocytopenia - UpToDate*, UpToDate. Available at: <https://www.uptodate.com/contents/approach-to-the-adult-with-lymphocytosis-or-lymphocytopenia> (Accessed: 4 April 2018).
- Davis, K. E., Prasad, C., Vijayagopal, P., Juma, S. and Imrhan, V. (2016) 'Advanced Glycation End Products, Inflammation, and Chronic Metabolic Diseases: Links in a Chain?', *Critical Reviews in Food Science and Nutrition*, 56(6), pp. 989–998. doi: 10.1080/10408398.2012.744738.
- DeFronzo, R. A. and Tripathy, D. (2009) 'Skeletal muscle insulin resistance is the primary defect in type 2 diabetes.', *Diabetes care*. American Diabetes Association, 32 Suppl 2(Suppl 2), pp. S157-63. doi: 10.2337/dc09-S302.
- Dilektasli, E., Inaba, K., Haltmeier, T., Wong, M. D., Clark, D., Benjamin, E. R., Lam, L. and Demetriades, D. (2016) 'The prognostic value of neutrophil-to-lymphocyte ratio on mortality in critically ill trauma patients', *Journal of Trauma and Acute Care Surgery*, 81(5), pp. 882–888. doi: 10.1097/TA.0000000000000980.
- Donath, M. Y., Dalmas, É., Sauter, N. S. and Böni-Schnetzler, M. (2013) 'Inflammation in obesity and diabetes: Islet dysfunction and therapeutic opportunity', *Cell Metabolism*, 17(6), pp. 860–872. doi: 10.1016/j.cmet.2013.05.001.
- Donath, M. Y. and Shoelson, S. E. (2011) 'Type 2 diabetes as an inflammatory disease', *Nature Reviews Immunology*, 11(2), pp. 98–107. doi: 10.1038/nri2925.
- Eguchi, K., Manabe, I., Oishi-Tanaka, Y., Ohsugi, M., Kono, N., Ogata, F., Yagi, N., Ohto, U., et al. (2012) 'Saturated Fatty Acid and TLR Signaling Link β Cell Dysfunction and Islet Inflammation', *Cell Metabolism*, 15(4), pp. 518–533. doi: 10.1016/j.cmet.2012.01.023.
- El-Asrar, A. M. A. (2012) 'Role of inflammation in the pathogenesis of diabetic



- retinopathy.', *Middle East African journal of ophthalmology*. Wolters Kluwer -- Medknow Publications, 19(1), pp. 70–4. doi: 10.4103/0974-9233.92118.
- Espósito, K., Nappo, F., Marfella, R., Giugliano, G., Giugliano, F., Ciotola, M., Quagliaro, L., Ceriello, A., et al. (2002) 'Inflammatory cytokine concentrations are acutely increased by hyperglycemia in humans: role of oxidative stress.', *Circulation*, 106(16), pp. 2067–72. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/12379575> (Accessed: 21 January 2018).
- Fan, M., Su, M., Tan, Y., Liu, Q., Ren, Y., Li, L. and Lv, J. (2015) 'Gender, Age, and Education Level Modify the Association between Body Mass Index and Physical Activity: A Cross-Sectional Study in Hangzhou, China', *PLOS ONE*. Edited by X. Miao. Public Library of Science, 10(5), p. e0125534. doi: 10.1371/journal.pone.0125534.
- Feng, J.-R., Qiu, X., Wang, F., Chen, P.-F., Gao, Q., Peng, Y.-N., Lin, X., Liu, Q., et al. (2017) 'Diagnostic Value of Neutrophil-to-Lymphocyte Ratio and Platelet-to-Lymphocyte Ratio in Crohn's Disease', *Gastroenterology Research and Practice*. Hindawi, 2017, pp. 1–5. doi: 10.1155/2017/3526460.
- Fernández-Real, J. M. and Pickup, J. C. (2008) 'Innate immunity, insulin resistance and type 2 diabetes.', *Trends in endocrinology and metabolism: TEM*, 19(1), pp. 10–16. doi: 10.1016/j.tem.2007.10.004.
- Fernández-Real, J. M. and Pickup, J. C. (2012) 'Innate immunity, insulin resistance and type 2 diabetes', *Diabetologia*, 55(2), pp. 273–278. doi: 10.1007/s00125-011-2387-y.
- Fong, D., Aiello, L., Gardner, T., King, G., Blankenship, G., Cavallerano, J., Ferris, F. and Klein, R. (2003) 'Retinopathy in Diabetes', *Diabetes Care*, 27(Supplement 1), pp. S84–S87.
- Forget, P., Khalifa, C., Defour, J.-P., Latinne, D., Van Pel, M.-C. and De Kock, M. (2017) 'What is the normal value of the neutrophil-to-lymphocyte ratio?', *BMC research notes*. BioMed Central, 10(1), p. 12. doi: 10.1186/s13104-016-2335-5.
- Fribley, A., Zhang, K. and Kaufman, R. J. (2009) 'Regulation of apoptosis by the unfolded protein response.', *Methods in molecular biology (Clifton, N.J.). NIH Public Access*, 559, pp. 191–204. doi: 10.1007/978-1-60327-017-5_14.
- Garnita, D. (2012) 'Faktor Risiko Diabetes Melitus di Indonesia (Analisis Data Sakerti 2007)', *Fakultas Kesehatan Masyarakat Universitas Indonesia*.
- Gärtner, V. and Eigenthaler, T. K. (2008) 'Pathogenesis of diabetic macro- and microangiopathy.', *Clinical nephrology*, 70(1), pp. 1–9.
- Geerlings, S. E. and Hoepelman, A. I. (1999) 'Immune dysfunction in patients with diabetes mellitus (DM).', *FEMS immunology and medical microbiology*, 26(3–4), pp. 259–65. Available at:



<http://www.ncbi.nlm.nih.gov/pubmed/10575137> (Accessed: 9 December 2017).

Gerich, J. E. (2002) 'Is reduced first-phase insulin release the earliest detectable abnormality in individuals destined to develop type 2 diabetes?', *Diabetes*. American Diabetes Association, 51 Suppl 1(suppl 1), pp. S117-21. doi: 10.2337/DIABETES.51.2007.S117.

Goldberg, M. F. and Jampol, L. M. (1987) 'Knowledge of diabetic retinopathy before and 18 years after the Airlie House Symposium on Treatment of Diabetic Retinopathy.', *Ophthalmology*, 94(7), pp. 741-6. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/2889177> (Accessed: 14 January 2018).

Gologorsky, D., Thanos, A. and Vavvas, D. (2012) 'Therapeutic interventions against inflammatory and angiogenic mediators in proliferative diabetic retinopathy.', *Mediators of inflammation*. Hindawi Limited, 2012, p. 629452. doi: 10.1155/2012/629452.

Goldstein, D., Little, R., Lorenz, R., Malone, J., Nathan, D., Peterson, C. and Sacks, D. (2004) 'Tests of Glycemia in Diabetes.', *Diabetes Care*, [online] 27(7), pp.1761-1773. Available at: <http://care.diabetesjournals.org/content/27/7/1761.long> [Accessed 25 Apr. 2018].

Goodman, D. A., Goodman, C. B. and Monk, J. S. (1995) 'Use of the neutrophil:lymphocyte ratio in the diagnosis of appendicitis.', *The American surgeon*, 61(3), pp. 257-9. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/7887542> (Accessed: 31 January 2018).

Grossmann, V., Schmitt, V. H., Zeller, T., Panova-Noeva, M., Schulz, A., Laubert-Reh, D., Juenger, C., Schnabel, R. B., et al. (2015) 'Profile of the Immune and Inflammatory Response in Individuals With Prediabetes and Type 2 Diabetes.', *Diabetes care*. American Diabetes Association, 38(7), pp. 1356-64. doi: 10.2337/dc14-3008.

Gunnlaugsdottir, E., Halldorsdottir, S., Klein, R., Eiriksdottir, G., Klein, B. E., Benediktsson, R., Harris, T. B., Launer, L. J., et al. (2012) 'Retinopathy in old persons with and without diabetes mellitus: the Age, Gene/Environment Susceptibility--Reykjavik Study (AGES-R).', *Diabetologia*. NIH Public Access, 55(3), pp. 671-80. doi: 10.1007/s00125-011-2395-y.

Guo, X., Zhang, S., Zhang, Q., Liu, L., Wu, H., Du, H., Shi, H., Wang, C., et al. (2015) 'Neutrophil:lymphocyte ratio is positively related to type 2 diabetes in a large-scale adult population: a Tianjin Chronic Low-Grade Systemic Inflammation and Health cohort study.', *European journal of endocrinology*. European Society of Endocrinology, 173(2), pp. 217-25. doi: 10.1530/EJE-15-0176.

Gupta, S. (2002) 'Tumor necrosis factor-alpha-induced apoptosis in T cells from aged humans: a role of TNFR-I and downstream signaling molecules.', *Experimental gerontology*, 37(2-3), pp. 293-9. Available at:



<http://www.ncbi.nlm.nih.gov/pubmed/11772515> (Accessed: 24 January 2018).

Hammes, H.-P., Welp, R., Kempe, H.-P., Wagner, C., Siegel, E., Holl, R. W. and Mellitus, D. I. B. C. N. D. (2015) 'Risk Factors for Retinopathy and DME in Type 2 Diabetes—Results from the German/Austrian DPV Database', *PLOS ONE*. Edited by O. Strauß. Public Library of Science, 10(7), p. e0132492. doi: 10.1371/journal.pone.0132492.

Hernández, C. and Simó, R. (2012) 'Erythropoietin produced by the retina: Its role in physiology and diabetic retinopathy', *Endocrine*, 41(2), pp. 220–226. doi: 10.1007/s12020-011-9579-6.

Hietala, K., Forsblom, C., Summanen, P. and Groop, P. (2008) 'Heritability of Proliferative Diabetic Retinopathy', *Diabetes*, 57(8), pp. 2176-2180.

Hinman, R. M. and Cambier, J. C. (2014) 'Role of B Lymphocytes in the Pathogenesis of Type 1 Diabetes', *Current Diabetes Reports*, 14(11), p. 543. doi: 10.1007/s11892-014-0543-8.

Hotamisligil, G. S., Arner, P., Caro, J. F., Atkinson, R. L. and Spiegelman, B. M. (1995) 'Increased adipose tissue expression of tumor necrosis factor-alpha in human obesity and insulin resistance.', *Journal of Clinical Investigation*, 95(5), pp. 2409–2415. doi: 10.1172/JCI117936.

Hotamisligil, G. S., Shargill, N. S. and Spiegelman, B. M. (1993) 'Adipose expression of tumor necrosis factor-alpha: direct role in obesity-linked insulin resistance.', *Science (New York, N.Y.)*, 259(5091), pp. 87–91. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/7678183> (Accessed: 11 January 2018).

Huang, L., Xie, Y., Dai, S. and Zheng, H. (2017) 'Neutrophil-to-lymphocyte ratio in diabetic microangiopathy', 10(2), pp. 1223–1232.

Imtiaz, F., Shafique, K., Mirza, S. S., Ayoob, Z., Vart, P. and Rao, S. (2012) 'Neutrophil lymphocyte ratio as a measure of systemic inflammation in prevalent chronic diseases in Asian population.', *International archives of medicine*. BioMed Central, 5(1), p. 2. doi: 10.1186/1755-7682-5-2.

International Diabetes Federation (2017) *IDF Diabetes Atlas*. 8th edn. Edited by S. Karurangan, J. da R. Fernandes, Y. Huang, and B. Malanda.

Ip, B. C., Hogan, A. E. and Nikolajczyk, B. S. (2015) 'Lymphocyte roles in metabolic dysfunction: Of men and mice', *Trends in Endocrinology and Metabolism*. Elsevier Ltd, 26(2), pp. 91–100. doi: 10.1016/j.tem.2014.12.001.

Joussen, A. M., Murata, T., Tsujikawa, A., Kirchhof, B., Bursell, S. E. and Adamis, A. P. (2001) 'Leukocyte-mediated endothelial cell injury and death in the diabetic retina.', *The American journal of pathology*. American Society for Investigative Pathology, 158(1), pp. 147–52. doi: 10.1016/S0002-9440(10)63952-1.



- Joussen, A. M., Poulaki, V., Mitsiades, N., Cai, W.-Y., Suzuma, I., Pak, J., Ju, S.-T., Rook, S. L., *et al.* (2003) ‘Suppression of Fas-FasL-induced endothelial cell apoptosis prevents diabetic blood–retinal barrier breakdown in a model of streptozotocin-induced diabetes’, *The FASEB Journal*, 17(1), pp. 76–78. doi: 10.1096/fj.02-0157fje.
- Kahn, S. E., Cooper, M. E. and Del Prato, S. (2014) ‘Pathophysiology and treatment of type 2 diabetes: perspectives on the past, present, and future.’, *Lancet (London, England)*. NIH Public Access, 383(9922), pp. 1068–83. doi: 10.1016/S0140-6736(13)62154-6.
- Kalani, M. (2008) ‘The importance of endothelin-1 for microvascular dysfunction in diabetes.’, *Vascular health and risk management*. Dove Press, 4(5), pp. 1061–8. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19183753> (Accessed: 21 January 2018).
- Von Känel, R., Mills, P. J. and Dimsdale, J. E. (2001) ‘Short-term hyperglycemia induces lymphopenia and lymphocyte subset redistribution’, *Life Sciences*, 69(3), pp. 255–262. doi: 10.1016/S0024-3205(01)01127-4.
- Kashyap, S. R., Belfort, R., Berria, R., Suraamornkul, S., Pratipranawatr, T., Finlayson, J., Barrentine, A., Bajaj, M., *et al.* (2004) ‘Discordant effects of a chronic physiological increase in plasma FFA on insulin signaling in healthy subjects with or without a family history of type 2 diabetes’, *American Journal of Physiology-Endocrinology and Metabolism*, 287(3), pp. E537–E546. doi: 10.1152/ajpendo.00541.2003.
- Katusić, D., Tomić, M., Jukić, T., Kordić, R., Sikić, J., Vukojević, N. and Sarić, B. (2005) ‘Obesity--a risk factor for diabetic retinopathy in type 2 diabetes?’, *Collegium antropologicum*, 29 Suppl 1, pp. 47–50. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16193676> (Accessed: 9 December 2017).
- Kern, T. S. (2007) ‘Contributions of inflammatory processes to the development of the early stages of diabetic retinopathy.’, *Experimental diabetes research*. Hindawi, 2007, p. 95103. doi: 10.1155/2007/95103.
- Khan, S., Raghuram, G. V., Bhargava, A., Pathak, N., Chandra, D. H., Jain, S. K. and Mishra, P. K. (2011) ‘Role and clinical significance of lymphocyte mitochondrial dysfunction in type 2 diabetes mellitus’, *Translational Research*, 158(6), pp. 344–359. doi: 10.1016/j.trsl.2011.08.007.
- Khandare, S. A., Chittawar, S., Nahar, N., Dubey, T. N. and Qureshi, Z. (2017) ‘Study of Neutrophil-lymphocyte Ratio as Novel Marker for Diabetic Nephropathy in Type 2 Diabetes.’, *Indian journal of endocrinology and metabolism*. Wolters Kluwer -- Medknow Publications, 21(3), pp. 387–392. doi: 10.4103/ijem.IJEM_476_16.
- Laugesen, K., Jørgensen, J. O. L., Sørensen, H. T. and Petersen, I. (2017) ‘Systemic glucocorticoid use in Denmark: A population-based prevalence study’, *BMJ Open*, 7(5). doi: 10.1136/bmjopen-2016-015237.



- Leasher, J. L., Bourne, R. R. A., Flaxman, S. R., Jonas, J. B., Keeffe, J., Naidoo, K., Pesudovs, K., Price, H., *et al.* (2016) ‘Global estimates on the number of people blind or visually impaired by diabetic retinopathy: A meta-analysis from 1990 to 2010’, *Diabetes Care*, 39(9), pp. 1643–1649. doi: 10.2337/dc15-2171.
- Lee, R., Wong, T. Y. and Sabanayagam, C. (2015) ‘Epidemiology of diabetic retinopathy, diabetic macular edema and related vision loss.’, *Eye and vision (London, England)*. BioMed Central, 2, p. 17. doi: 10.1186/s40662-015-0026-2.
- Lemasters, J. J., Qian, T., He, L., Kim, J.-S., Elmore, S. P., Cascio, W. E. and Brenner, D. A. (2002) ‘Role of Mitochondrial Inner Membrane Permeabilization in Necrotic Cell Death, Apoptosis, and Autophagy’, *Antioxidants & Redox Signaling*, 4(5), pp. 769–781. doi: 10.1089/152308602760598918.
- Ley, S. H., Hamdy, O., Mohan, V. and Hu, F. B. (2014) ‘Prevention and management of type 2 diabetes: dietary components and nutritional strategies.’, *Lancet (London, England)*. NIH Public Access, 383(9933), pp. 1999–2007. doi: 10.1016/S0140-6736(14)60613-9.
- Li, X., Shen, J., Lu, Z., Chen, M., Fang, X. and Wang, G. (2017) ‘High neutrophil-to-lymphocyte ratio is associated with increased carotid artery intima-media thickness in type 2 diabetes.’, *Journal of diabetes investigation*. Wiley-Blackwell, 8(1), pp. 101–107. doi: 10.1111/jdi.12541.
- Li, Y., Xu, W., Liao, Z., Yao, B., Chen, X., Huang, Z., Hu, G. and Weng, J. (2004) ‘Induction of long-term glycemic control in newly diagnosed type 2 diabetic patients is associated with improvement of beta-cell function.’, *Diabetes care*, 27(11), pp. 2597–602. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15504992> (Accessed: 14 December 2017).
- Lima, V. C., Cavalieri, G. C., Lima, M. C., Nazario, N. O. and Lima, G. C. (2016) ‘Risk factors for diabetic retinopathy: a case-control study.’, *International journal of retina and vitreous*. BioMed Central, 2, p. 21. doi: 10.1186/s40942-016-0047-6.
- Lou, M., Luo, P., Tang, R., Peng, Y., Yu, S., Huang, W. and He, L. (2015) ‘Relationship between neutrophil-lymphocyte ratio and insulin resistance in newly diagnosed type 2 diabetes mellitus patients.’, *BMC endocrine disorders*. BioMed Central, 15, p. 9. doi: 10.1186/s12902-015-0002-9.
- Maghbooli, Z., Pasalar, P., Keshtkar, A., Farzadfar, F. and Larijani, B. (2014) ‘Predictive factors of diabetic complications: a possible link between family history of diabetes and diabetic retinopathy’, *Journal of Diabetes & Metabolic Disorders*, 13(1), p. 55. doi: 10.1186/2251-6581-13-55.
- Marseglia, L., Manti, S., D’Angelo, G., Nicotera, A., Parisi, E., Di Rosa, G., Gitto, E. and Arrigo, T. (2014) ‘Oxidative stress in obesity: a critical



component in human diseases.', *International journal of molecular sciences*. Multidisciplinary Digital Publishing Institute (MDPI), 16(1), pp. 378–400. doi: 10.3390/ijms16010378.

Meigs, J. B. (2017) *The metabolic syndrome (insulin resistance syndrome or syndrome X)* - UpToDate. Available at: https://www.uptodate.com/contents/the-metabolic-syndrome-insulin-resistance-syndrome-or-syndrome-x?source=see_link (Accessed: 14 December 2017).

Mertoglu, C. and Gunay, M. (2017) 'Neutrophil-Lymphocyte ratio and Platelet-Lymphocyte ratio as useful predictive markers of prediabetes and diabetes mellitus', *Diabetes & Metabolic Syndrome: Clinical Research & Reviews*, 11, pp. S127–S131. doi: 10.1016/j.dsx.2016.12.021.

Moursy, E. Y., Megallaa, M. H., Mouftah, R. F. and Ahmed, S. M. (2015) 'Relationship Between Neutrophil-Lymphocyte Ratio and Microvascular Complications in Egyptian Patients with Type 2 ... Relationship Between Neutrophil-Lymphocyte Ratio and Microvascular Complications in Egyptian Patients with Type 2 Diabetes', 3(December), pp. 250–255. doi: 10.11648/j.ajim.20150306.16.

Naeim, F., Nagesh Rao, P., Song, S. X., Grody, W. W., Naeim, F., Nagesh Rao, P., Song, S. X. and Grody, W. W. (2013) 'Granulocytic Disorders', in *Atlas of Hematopathology*. Elsevier, pp. 663–673. doi: 10.1016/B978-0-12-385183-3.00060-7.

Nakahara, K., Monden, Y., Ohno, K., Fujii, Y., Hashimoto, J., Kitagawa, Y. and Kawashima, Y. (1987) 'Importance of biologic status to the postoperative prognosis of patients with stage III nonsmall cell lung cancer.', *Journal of surgical oncology*, 36(3), pp. 155–60.

Nentwich, M. M. and Ulbig, M. W. (2015) 'Diabetic retinopathy - ocular complications of diabetes mellitus.', *World journal of diabetes*. Baishideng Publishing Group Inc, 6(3), pp. 489–99. doi: 10.4239/wjd.v6.i3.489.

Ni, A. (2016) 'Reference Values of Neutrophil-Lymphocyte Ratio, Platelet-Lymphocyte Ratio and Mean Platelet Volume in Healthy Adults in North Central Nigeria', *Journal of Blood & Lymph*. OMICS International, 6(1). doi: 10.4172/2165-7831.1000143.

Nieto-Vazquez, I., Fernández-Veledo, S., Krämer, D. K., Vila-Bedmar, R., García-Guerra, L. and Lorenzo, M. (2008) 'Insulin resistance associated to obesity: the link TNF-alpha', *Archives of Physiology and Biochemistry*, 114(3), pp. 183–194. doi: 10.1080/13813450802181047.

Ntzani, E., Liberopoulos, G., Manolio, T. and Ioannidis, J. (2011) 'Consistency of genome-wide associations across major ancestral groups', *Human Genetics*, 131(7), pp. 1057-1071. doi: 10.1007/s00439-011-1124-4.

Otton, R., Soriano, F. G., Verlengia, R. and Curi, R. (2004) 'Diabetes induces



- apoptosis in lymphocytes', *Journal of Endocrinology*, 182(1), pp. 145–156. doi: 10.1677/joe.0.1820145.
- Perseghin, G., Ghosh, S., Gerow, K. and Shulman, G. I. (1997) 'Metabolic defects in lean nondiabetic offspring of NIDDM parents: a cross-sectional study.', *Diabetes*. American Diabetes Association, 46(6), pp. 1001–9. doi: 10.2337/DIAB.46.6.1001.
- Perseghin, G., Petersen, K. and Shulman, G. I. (2003) 'Cellular mechanism of insulin resistance: potential links with inflammation', *International Journal of Obesity*. Nature Publishing Group, 27(S3), pp. S6–S11. doi: 10.1038/sj.ijo.0802491.
- Petersen, K. F. and Shulman, G. I. (2006) 'Etiology of insulin resistance.', *The American journal of medicine*. NIH Public Access, 119(5 Suppl 1), pp. S10-6. doi: 10.1016/j.amjmed.2006.01.009.
- Porte, D. and Kahn, S. E. (1995) 'The key role of islet dysfunction in type II diabetes mellitus.', *Clinical and investigative medicine. Medecine clinique et experimentale*, 18(4), pp. 247–54. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/8549009> (Accessed: 10 January 2018).
- Ramachandran, A., Ma, R. C. W. and Snehalatha, C. (2010) 'Diabetes in Asia', *Lancet (London, England)*. Elsevier, 375(9712), pp. 408–18. doi: 10.1016/S0140-6736(09)60937-5.
- Reaven, G. M. (2005) 'Insulin resistance, the insulin resistance syndrome, and cardiovascular disease.', *Panminerva medica*, 47(4), pp. 201–10. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16489319> (Accessed: 14 December 2017).
- Rifkin, R. M. (2010) 'Neutrophilia', in *Decision Making in Medicine*. Elsevier, pp. 240–241. doi: 10.1016/B978-0-323-04107-2.50089-2.
- Riva, C. E., Logean, E. and Falsini, B. (2005) 'Visually evoked hemodynamical response and assessment of neurovascular coupling in the optic nerve and retina', *Progress in Retinal and Eye Research*, 24(2), pp. 183–215. doi: 10.1016/j.preteyes.2004.07.002.
- Robinson, R., Barathi, V. A., Chaurasia, S. S., Wong, T. Y. and Kern, T. S. (2012) 'Update on animal models of diabetic retinopathy: from molecular approaches to mice and higher mammals.', *Disease models & mechanisms*. The Company of Biologists Ltd, 5(4), pp. 444–56. doi: 10.1242/dmm.009597.
- Roy, S., Ha, J., Trudeau, K. and Beglova, E. (2010) 'Vascular basement membrane thickening in diabetic retinopathy', *Current Eye Research*, 35(12), pp. 1045–1056. doi: 10.3109/02713683.2010.514659.
- Russo, G. T., Giorda, C. B., Cercone, S., Nicolucci, A., Cucinotta, D., Leotta, S., Arcangeli, A., Calabrese, M., et al. (2014) 'Factors associated with beta-cell dysfunction in type 2 diabetes: The BETADECLINE study', *PLoS ONE*,



- 9(10). doi: 10.1371/journal.pone.0109702.
- Saghizadeh, M., Ong, J. M., Garvey, W. T., Henry, R. R. and Kern, P. A. (1996) ‘The expression of TNF alpha by human muscle. Relationship to insulin resistance.’, *Journal of Clinical Investigation*, 97(4), pp. 1111–1116. doi: 10.1172/JCI118504.
- Sakowicz-Burkiewicz, M., Kocbuch, K., Grden, M., Szutowicz, A. and Pawelczyk, T. (2006) ‘Diabetes-induced decrease of adenosine kinase expression impairs the proliferation potential of diabetic rat T lymphocytes.’, *Immunology*. Wiley-Blackwell, 118(3), pp. 402–12. doi: 10.1111/j.1365-2567.2006.02380.x.
- Sanders, L. (2002) ‘From Thebes to Toronto and the 21st Century: An Incredible Journey’, *Diabetes Spectrum*, 15(1), pp. 56–60. doi: 10.2337/diaspect.15.1.56.
- Sasongko, M. B., Widyaputri, F., Agni, A. N., Wardhana, F. S., Kotha, S., Gupta, P., Widayanti, T. W., Haryanto, S., et al. (2017) ‘Prevalence of Diabetic Retinopathy and Blindness in Indonesian Adults With Type 2 Diabetes’, *American Journal of Ophthalmology*, 181, pp. 79–87. doi: 10.1016/j.ajo.2017.06.019.
- Sayiner, Z. A., Eraydin, A., Atakur, S. and Özkaya, M. (2017) ‘The Relationship Between Microvascular Complications in Type 2 Diabetes with Mean Platelet Volume, Red Blood Cell Distribution Width, Neutrophile Lymphocyte Ratio’, *Turkiye Klinikleri Journal of Internal Medicine*, 2(3), pp. 113–117. doi: 10.5336/intermed.2017-57111.
- Sefil, F., Ulutas, K. T., Dokuyucu, R., Sumbul, A. T., Yengil, E., Yagiz, A. E., Yula, E., Ustun, I., et al. (2014) ‘Investigation of neutrophil lymphocyte ratio and blood glucose regulation in patients with type 2 diabetes mellitus’, *Journal of International Medical Research*, 42(2), pp. 581–588. doi: 10.1177/0300060513516944.
- Selders, G. S., Fetz, A. E., Radic, M. Z. and Bowlin, G. L. (2017) ‘An overview of the role of neutrophils in innate immunity, inflammation and host-biomaterial integration.’, *Regenerative biomaterials*. Oxford University Press, 4(1), pp. 55–68. doi: 10.1093/rb/rbw041.
- Semeraro, F., Cancarini, A., Dell’Omo, R., Rezzola, S., Romano, M. R. and Costagliola, C. (2015) ‘Diabetic retinopathy: Vascular and inflammatory disease’, *Journal of Diabetes Research*. Hindawi Publishing Corporation, 2015(July). doi: 10.1155/2015/582060.
- Sharma, Y., Saxena, S., Mishra, A., Saxena, A. and Natu, S. M. (2012) ‘Advanced glycation end products and diabetic retinopathy.’, *Journal of ocular biology, diseases, and informatics*. Springer, 5(3–4), pp. 63–9. doi: 10.1007/s12177-013-9104-7.
- Shiny, A., Bibin, Y. S., Shanthirani, C. S., Regin, B. S., Anjana, R. M.,



- Balasubramanyam, M., Jebarani, S. and Mohan, V. (2014) ‘Association of Neutrophil-Lymphocyte Ratio with Glucose Intolerance: An Indicator of Systemic Inflammation in Patients with Type 2 Diabetes’, *Diabetes Technology & Therapeutics*, 16(8), pp. 524–530. doi: 10.1089/dia.2013.0264.
- Simó-Servat, O., Hernández, C. and Simó, R. (2013) ‘Genetics in diabetic retinopathy: current concepts and new insights.’, *Current genomics*, 14(5), pp. 289–99. doi: 10.2174/13892029113149990008.
- Simó, R., Carrasco, E., García-Ramírez, M. and Hernández, C. (2006) ‘Angiogenic and antiangiogenic factors in proliferative diabetic retinopathy.’, *Current diabetes reviews*, 2(1), pp. 71–98. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18220619> (Accessed: 21 January 2018).
- Soehnlein, O., Steffens, S., Hidalgo, A. and Weber, C. (2017) ‘Neutrophils as protagonists and targets in chronic inflammation’, *Nature Reviews Immunology*. Nature Publishing Group, 17(4), pp. 248–261. doi: 10.1038/nri.2017.10.
- Song, H., Wang, L. and Hui, Y. (2006) ‘Expression of CD18 on the neutrophils of patients with diabetic retinopathy’, *Graefe’s Archive for Clinical and Experimental Ophthalmology*, 245(1), pp. 24–31. doi: 10.1007/s00417-006-0379-2.
- Soro-Paavonen, A., Zhang, W.-Z., Venardos, K., Coughlan, M. T., Harris, E., Tong, D. C., Brasacchio, D., Paavonen, K., et al. (2010) ‘Advanced glycation end-products induce vascular dysfunction via resistance to nitric oxide and suppression of endothelial nitric oxide synthase’, *Journal of Hypertension*, 28(4), pp. 780–788. doi: 10.1097/HJH.0b013e328335043e.
- Srivastava, B. K. and Rema, M. (2005) ‘Does hypertension play a role in diabetic retinopathy?’, *Journal of Association of Physicians of India*, pp. 803–808.
- Stemmelin, G. R., Doti, C. A., Shanley, C. M., Ceresetto, J. M., Rabinovich, O. M., Vicente Reparaz, M. A., Vukovic, M. G. and Bullorsky, E. O. (2004) ‘Smoking as a Cause for Mild Chronic Neutrophilia.’, *Blood*, 104(11). Available at: <http://www.bloodjournal.org/content/104/11/3796?sso-checked=true> (Accessed: 29 January 2018).
- Stumvoll, M., Goldstein, B. J. and van Haeften, T. W. (2005) ‘Type 2 diabetes: principles of pathogenesis and therapy.’, *Lancet (London, England)*, 365(9467), pp. 1333–46. doi: 10.1016/S0140-6736(05)61032-X.
- Talchai, C., Lin, H. V., Kitamura, T. and Accili, D. (2009) ‘Genetic and biochemical pathways of β-cell failure in type 2 diabetes’, *Diabetes, Obesity and Metabolism*, 11(SUPPL. 4), pp. 38–45. doi: 10.1111/j.1463-1326.2009.01115.x.
- Tanaka, S., Isoda, F., Ishihara, Y., Kimura, M. and Yamakawa, T. (2001) ‘T lymphopenia in relation to body mass index and TNF-alpha in human



obesity: adequate weight reduction can be corrective.', *Clinical endocrinology*, 54(3), pp. 347–54. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11298087> (Accessed: 9 December 2017).

Tang, C., Naassan, A. E., Chamson-Reig, A., Koulajian, K., Goh, T. T., Yoon, F., Oprescu, A. I., Ghanim, H., et al. (2013) 'Susceptibility to Fatty Acid-Induced β -Cell Dysfunction Is Enhanced in Prediabetic Diabetes-Prone BioBreeding Rats: A Potential Link Between β -Cell Lipotoxicity and Islet Inflammation', *Endocrinology*. Oxford University Press, 154(1), pp. 89–101. doi: 10.1210/en.2012-1720.

Tang, J. and Kern, T. S. (2011) 'Inflammation in diabetic retinopathy.', *Progress in retinal and eye research*. NIH Public Access, 30(5), pp. 343–58. doi: 10.1016/j.preteyes.2011.05.002.

Taouk Y, F. E., EK, F., Y, T., TE, N., JJ, W., TY, W. and EL, L. (2014) 'Relative Importance and Contribution of Risk Factors for Diabetic Retinopathy and Macular Edema', *Journal of Diabetes & Metabolism*. OMICS International, 5(2), pp. 1–7. doi: 10.4172/2155-6156.1000337.

Tapp, R. J., Shaw, J. E., Harper, C. A., de Courten, M. P., Balkau, B., McCarty, D. J., Taylor, H. R., Welborn, T. A., et al. (2003) 'The prevalence of and factors associated with diabetic retinopathy in the Australian population.', *Diabetes care*. American Diabetes Association, 26(6), pp. 1731–7. doi: 10.2337/DIACARE.26.6.1731.

Tarr, J. M., Kaul, K., Chopra, M., Kohner, E. M. and Chibber, R. (2013) 'Pathophysiology of Diabetic Retinopathy', *ISRN Ophthalmol*, 2013(January), pp. 1–13. doi: 10.1155/2013/343560.

Templeton, A. J., McNamara, M. G., Šeruga, B., Vera-Badillo, F. E., Aneja, P., Ocaña, A., Leibowitz-Amit, R., Sonpavde, G., et al. (2014) 'Prognostic Role of Neutrophil-to-Lymphocyte Ratio in Solid Tumors: A Systematic Review and Meta-Analysis', *JNCI: Journal of the National Cancer Institute*, 106(6), p. dju124. doi: 10.1093/jnci/dju124.

Treffers, L. W., Hiemstra, I. H., Kuijpers, T. W., van den Berg, T. K. and Matlung, H. L. (2016) 'Neutrophils in cancer', *Immunological Reviews*, 273(1), pp. 312–328. doi: 10.1111/imr.12444.

Ulu, S., Bucak, A., Ulu, M. S., Ahsen, A., Duran, A., Yucedag, F. and Aycicek, A. (2014) 'Neutrophil-lymphocyte ratio as a new predictive and prognostic factor at the hearing loss of diabetic patients', *European Archives of Oto-Rhino-Laryngology*. Springer Berlin Heidelberg, 271(10), pp. 2681–2686. doi: 10.1007/s00405-013-2734-3.

Ulu, S. M., Dogan, M., Ahsen, A., Altug, A., Demir, K., Acartürk, G. and Inan, S. (2013) 'Neutrophil-to-Lymphocyte Ratio as a Quick and Reliable Predictive Marker to Diagnose the Severity of Diabetic Retinopathy', *Diabetes Technology & Therapeutics*, 15(11), pp. 942–947. doi:



10.1089/dia.2013.0097.

- Vidro, E. K., Gee, S., Unda, R., Ma, J. and Tsin, A. (2008) ‘Glucose and TGFbeta2 modulate the viability of cultured human retinal pericytes and their VEGF release.’, *Current eye research*. NIH Public Access, 33(11), pp. 984–93. doi: 10.1080/02713680802450976.
- Vigili de Kreutzenberg, S., Coracina, A., Volpi, A., Fadini, G. P., Frigo, A. C., Guarneri, G., Tiengo, A. and Avogaro, A. (2011) ‘Microangiopathy is independently associated with presence, severity and composition of carotid atherosclerosis in type 2 diabetes’, *Nutrition, Metabolism and Cardiovascular Diseases*. Elsevier Ltd, 21(4), pp. 286–293. doi: 10.1016/j.numecd.2009.10.003.
- Wang, J., Xu, X., Elliott, M. H., Zhu, M. and Le, Y.-Z. (2010) ‘Müller cell-derived VEGF is essential for diabetes-induced retinal inflammation and vascular leakage.’, *Diabetes*. American Diabetes Association, 59(9), pp. 2297–305. doi: 10.2337/db09-1420.
- Wang, Q., Shen, J. J. and Cochran, C. (2016) ‘Unemployment Rate, Smoking in China: Are They Related?’, *International journal of environmental research and public health*. Multidisciplinary Digital Publishing Institute (MDPI), 13(1). doi: 10.3390/ijerph13010113.
- Wang, R., Zhang, J., Li, Y., Liu, T. and Yu, K. (2015) ‘Neutrophil–Lymphocyte ratio is associated with arterial stiffness in diabetic retinopathy in type 2 diabetes’, *Journal of Diabetes and its Complications*, 29(2), pp. 245–249. doi: 10.1016/j.jdiacomp.2014.11.006.
- Weisberg, S. P., McCann, D., Desai, M., Rosenbaum, M., Leibel, R. L. and Ferrante, A. W. (2003) ‘Obesity is associated with macrophage accumulation in adipose tissue’, *Journal of Clinical Investigation*, 112(12), pp. 1796–1808. doi: 10.1172/JCI19246.
- Wellen, K. E. and Hotamisligil, G. S. (2005) ‘Inflammation, stress, and diabetes.’, *The Journal of clinical investigation*. American Society for Clinical Investigation, 115(5), pp. 1111–9. doi: 10.1172/JCI25102.
- WHO (2006) ‘Definition and Diagnosis of Diabetes Mellitus and Intermediate Hyperglycemia’, *Who2*, p. 50. doi: ISBN 92 4 159493 4.
- ‘WHO | Diabetes mellitus’ (2010) *WHO*. World Health Organization. Available at: <http://www.who.int/mediacentre/factsheets/fs138/en/> (Accessed: 3 September 2017).
- Wilkinson, C. ., Ferris, F. L., Klein, R. E., Lee, P. P., Agardh, C. D., Davis, M., Dills, D., Kampik, A., et al. (2003) ‘Proposed international clinical diabetic retinopathy and diabetic macular edema disease severity scales’, *Ophthalmology*, 110(9), pp. 1677–1682. doi: 10.1016/S0161-6420(03)00475-5.



- Wong, T. Y., Cheung, C. M. G., Larsen, M., Sharma, S. and Simó, R. (2016) ‘Diabetic retinopathy’, *Nature Reviews Disease Primers*, 2(March). doi: 10.1038/nrdp.2016.12.
- Woo, S. J., Ahn, S. J., Ahn, J., Park, K. H. and Lee, K. (2011) ‘Elevated Systemic Neutrophil Count in Diabetic Retinopathy and Diabetes: A Hospital-Based Cross-sectional Study of 30,793 Korean Subjects’, *Investigative Ophthalmology & Visual Science*. The Association for Research in Vision and Ophthalmology, 52(10), p. 7697. doi: 10.1167/iovs.11-7784.
- World Health Organization (2016) ‘Global Report on Diabetes’, *Isbn*, 978, p. 88. doi: ISBN 978 92 4 156525 7.
- Wu, L., Fernandez-Loaiza, P., Sauma, J., Hernandez-Bogantes, E. and Masis, M. (1991) ‘Grading diabetic retinopathy from stereoscopic color fundus photographs--an extension of the modified Airlie House classification. ETDRS report number 10. Early Treatment Diabetic Retinopathy Study Research Group.’, *Ophthalmology*, 98(5 Suppl), pp. 786–806. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/2062513> (Accessed: 14 January 2018).
- Wu, L., Fernandez-Loaiza, P., Sauma, J., Hernandez-Bogantes, E. and Masis, M. (2013) ‘Classification of diabetic retinopathy and diabetic macular edema’, *World journal of diabetes*. Baishideng Publishing Group Inc, 4(6), pp. 290–4. doi: 10.4239/wjd.v4.i6.290.
- Xia, C., Rao, X. and Zhong, J. (2017) ‘Role of T Lymphocytes in Type 2 Diabetes and Diabetes-Associated Inflammation’, *Journal of Diabetes Research*. Hindawi, 2017, pp. 1–6. doi: 10.1155/2017/6494795.
- Xu, H., Barnes, G. T., Yang, Q., Tan, G., Yang, D., Chou, C. J., Sole, J., Nichols, A., et al. (2003) ‘Chronic inflammation in fat plays a crucial role in the development of obesity-related insulin resistance.’, *The Journal of clinical investigation*. American Society for Clinical Investigation, 112(12), pp. 1821–30. doi: 10.1172/JCI19451.
- Xu, H. and Chen, M. (2017) ‘Diabetic retinopathy and dysregulated innate immunity’, *Vision Research*. Elsevier Ltd, 139, pp. 39–46. doi: 10.1016/j.visres.2017.04.013.
- Xu, H., Chen, Y., Li, Y., Xia, F., Han, B., Zhang, H., Zhai, H., Wu, H., et al. (2014) ‘Mitochondrial apoptosis of lymphocyte is induced in type 2 diabetes’, *Chin Med J (Engl)*, 127(2), pp. 213–217. doi: 10.3760/cma.j.issn.0366-6999.20131906.
- Yau, J. W. Y., Rogers, S. L., Kawasaki, R., Lamoureux, E. L., Kowalski, J. W., Bek, T., Chen, S.-J., Dekker, J. M., et al. (2012) ‘Global Prevalence and Major Risk Factors of Diabetic Retinopathy’, *Diabetes Care*, 35(3), pp. 556–564. doi: 10.2337/dc11-1909.
- Yue, S., Zhang, J., Wu, J., Teng, W., Liu, L., and Chen, L. (2015) ‘Use of the Monocyte-to-Lymphocyte Ratio to Predict Diabetic Retinopathy’,



International Journal of Environmental Research and Public Health, 12(8),
10009–10019. doi: 10.3390/ijerph120810009.

- Zeng, C., Shi, X., Zhang, B., Liu, H., Zhang, L., Ding, W. and Zhao, Y. (2012) ‘The imbalance of Th17/Th1/Tregs in patients with type 2 diabetes: Relationship with metabolic factors and complications’, *Journal of Molecular Medicine*. Springer-Verlag, 90(2), pp. 175–186. doi: 10.1007/s00109-011-0816-5.
- Zhang, X., Saaddine, J. B., Chou, C.-F., Cotch, M. F., Cheng, Y. J., Geiss, L. S., Gregg, E. W., Albright, A. L., *et al.* (2010) ‘Prevalence of Diabetic Retinopathy in the United States, 2005–2008’, *JAMA*, 304(6), p. 649. doi: 10.1001/jama.2010.1111.
- Zhou, B., Lu, Y., Hajifathalian, K., Bentham, J., Di Cesare, M., Danaei, G., Bixby, H., Cowan, M. J., *et al.* (2016) ‘Worldwide trends in diabetes since 1980: A pooled analysis of 751 population-based studies with 4.4 million participants’, *The Lancet. NCD Risk Factor Collaboration*. Open Access article distributed under the terms of CC BY, 387(10027), pp. 1513–1530. doi: 10.1016/S0140-6736(16)00618-8.
- Zhou, R., Tardivel, A., Thorens, B., Choi, I. and Tschopp, J. (2010) ‘Thioredoxin-interacting protein links oxidative stress to inflammasome activation’, *Nature Immunology*, 11(2), pp. 136–140. doi: 10.1038/ni.1831.
- Zhou, W., Chu, Y. and An, G. (2017) ‘Significant difference of neutrophil-lymphocyte ratio between colorectal cancer, adenomatous polyp and healthy people’, *European Review for Medical and Pharmacological Sciences*, 21, pp. 5386–5391.
- Zhu, Q. and Song, H.-P. (2015) ‘CD18 expression in granulocytes infiltrating the vitreous fluid in patients with diabetic retinopathy.’, *International journal of ophthalmology*. Press of International Journal of Ophthalmology, 8(3), pp. 508–12. doi: 10.3980/j.issn.2222-3959.2015.03.13.
- Zorrilla, E. P., Luborsky, L., McKay, J. R., Rosenthal, R., Houldin, A., Tax, A., McCorkle, R., Seligman, D. A., *et al.* (2001) ‘The Relationship of Depression and Stressors to Immunological Assays: A Meta-Analytic Review’, *Brain, Behavior, and Immunity*, 15(3), pp. 199–226. doi: 10.1006/bbri.2000.0597.