UNIVERSITAS
GADJAH MADA
**TINGKAT AKURASI METODE MAGNIFIKASI CANINE INCISOR INDEX DALAM PENENTUAN LETAK IMPAKSI KANINUS
MAKSILA PADA RADIOGRAF PANORAMIK RAS DEUTRO MELAYU**
ANTON, Drg. Rahardjo, S.U., Sp BM (K); drg. Bambang Dwiraharjo, Sp.BM (K)
Universitas Gadjah Mada, 2018 | Diunduh dari http://etd.repository.ugm.ac.id/

DAFTAR PUSTAKA

Al-Khateeb, S.N., Abu Alhaija, E.S.J., 2006. Tooth size discrepancies and arch parameters among different malocclusions in a Jordanian sample. *Angle Orthod*.76,459–465.doi:10.1043/00033219

Amal, N., 2011. *Hubungan analisis Bolton anterior dengan jarak gigit gigi insisivus sentralis.* http://referensi/hubungan-analisis-bolton-anterior.html (17 april 2011).

An, S., Wang, J., Li, J., Cheng, Q., Jiang, C.M.M., Wang, Y.T.T., Huang, Y.-F.F., Yu, W.J.J., Gou, Y.C.C., Xiao, L., 2013. Comparison of methods for localization of impacted maxillary canines by panoramic radiographs. *Dentomaxillofacial Radiol.* 42, 1–6. doi:10.1259/dmfr.20130129.

Armstrong, C., Johnston, C., Burden, D., Stevenson, M., 2003. Localizing ectopic maxillary canines - Horizontal or vertical parallax? *Eur. J. Orthod.* 25, 585–589. doi:10.1093/ejo/25.6.585.

Bennett, J., McLaughlin, R.P., 2002. *Orthodontic Management of the Dentition with the Preadjusted Appliance*. Isis Medical, Oxford (ISBN 899066 91 8). Republished in 2002 by Mosby, Edinburgh.

Boel, T, 2009. *Dental Radiologi : Prinsip dan Teknik,* USU Press, Medan.

Bayram, M., Özer, M., Sener, I., 2007. Maxillary canine impactions related to impacted central incisors: Two case reports. *J. Contemp. Dent. Pract.* 8, 72–81. doi:1526-3711-434 [pii].

Bedoya, M.M., Park, J.H., 2009. A Review of the Diagnosis and Management of Impacted Maxillary Canines. *J Am Dent Assoc* 140, 1485–1493. doi:140/12/1485 [pii].

Berkhout, W.E.R., 2015. The ALARA-principle. Backgrounds and enforcement in dental practices. *Ned. Tijdschr. Tandheelkd*. 122, 263–270.

Bishara, S.E., 1992. Impacted maxillary canines:a review.*Am J Orthod Dentofacial Orthop*.,101(2):159-171.

Bishara, S.E., 2001. *Textbook of Orthodontics*. 1st ed., Saunders. doi:10.1017/CBO9781107415324.004.

Bishara, S.E., 1998. Clinical management of impactedmaxillary canines. *Semin. Orthod.* 4, 87–98. doi:10.1016/S1073-8746(98)80006-6.

Budiman, J.A., Yashadana, E.D.D., Sadosa, S.D., Masbirin, P.I., 1997. Hubungan

UNIVERSITAS
GADJAH MADA
**TINGKAT AKURASI METODE MAGNIFIKASI CANINE INCISOR INDEX DALAM PENENTUAN LETAK IMPAKSI KANINUS
MAKSILA PADA RADIOGRAF PANORAMIK RAS DEUTRO MELAYU**
ANTON, Drg. Rahardjo, S.U., Sp BM (K); drg. Bambang Dwiraharjo, Sp.BM (K)
Universitas Gadjah Mada, 2018 | Diunduh dari http://etd.repository.ugm.ac.id/

rasio anterior dengan overjet dan overbite pada perawatan ortodontik. *Jurnal Kedokteran Gigi Universitas Indonesia.* 4(3): 19-25.

Budiraharjo, R., Pradopo, S., 2002. Ukuran mesiodistal gigi anak usia 12 tahun populasi Jawa dan Madura di Kabupaten Jember. *Jurnal PDGI*. 490-3.

Bujang, MA dan Baharum, N, 2017, Guidelines of the minimum sample size requirements for cohen's kappa, *EBPH*, 14(2):1-10

Chaushu, S., Chaushu, G., Becker, a, 1999. The use of panoramic radiographs to localize displaced maxillary canines. *Oral Surg. Oral Med. Oral Pathol. Oral Radiol. Endod.* 88, 511–516. doi:10.1016/S1079-2104(99)70072-7.

Christopher, P., Watanabe, A., Farman, A., Gon, M., Watanabe, D.C., Paulo, J., Issa, M., 2008. Radiographic Signals Detection of Systemic Disease . *Int. J. Morphol.* 26, 915–926. doi:10.4067/S0717-95022008000400021.

Cohen J.A., 1960. Coefficient of agreement for nominal scales. *Educ Psychol Meas*. 20:37–46.

Cohen, M.D., 2015. ALARA, Image Gently and CT-induced cancer. *Pediatr. Radiol*. 45, 465–470. doi:10.1007/s00247-014-3198-3.

Counihan, K., Al-Awadhi, E.A, Butler, J., 2013. Guidelines for the assessment of the impacted maxillary canine. *Dent. Update* 40, 770–772, 775–777.

Ericson, S., Kurol, J., 1988. Early treatment of palatally erupting maxillary canines by extraction of the primary canines. *Eur. J. Orthod.* 10, 283–295. doi:10.1093/ejo/10.4.283.

Faber, J., Berto, P.M., Quaresma, M., 2006. Rapid prototyping as a tool for diagnosis and treatment planning for maxillary canine impaction. *Am. J.* Orthod. Dentofac. Orthop. 129, 583–589. doi:10.1016/j.ajodo.2005.12.015.

Farman, A.G., Clark, S.J., Friedlander, A.H., Jacobs, W.R., Khan, Z., Kushner, G.M., Norman, K.M., Nortjé, C.J., Silveira, A., Wood, R.E., Yaggy, S.T., 2007. *Panoramic radiology: Seminars on maxillofacial imaging and interpretation*, doi:10.1007/978-3-540-46230-9.

Fox, N.A., Fletcher, G.A., Horner, K., 1995. Localising maxillary canines using dental panoramic tomography. *Br. Dent. J.* 179, 416–20. doi:10.1038/sj.bdj.4808945.

Fragiskos, F.D., Stefanou, E., Angelopoulou, E., Alexandridis, C., Giamarellou, H., Pefanis, A., Perdikaris, G., 2007. Oral surgery, *The New England journal of medicine*. doi:2006939050.

UNIVERSITAS
GADJAH MADA
**TINGKAT AKURASI METODE MAGNIFIKASI CANINE INCISOR INDEX DALAM PENENTUAN LETAK IMPAKSI KANINUS
MAKSILA PADA RADIOGRAF PANORAMIK RAS DEUTRO MELAYU**
ANTON, Drg. Rahardjo, S.U., Sp BM (K); drg. Bambang Dwiraharjo, Sp.BM (K)
Universitas Gadjah Mada, 2018 | Diunduh dari http://etd.repository.ugm.ac.id/

Frank, C. a, 2000. Treatment options for impacted teeth. *J. Am. Dent. Assoc.* 131, 623–632. doi:10.14219/jada.archive.2000.0236.

Gavel, V., Dermaut, L., 1999. The effect of tooth position on the image of unerupted canines on panoramic radiographs. *Eur. J. Orthod.* 21, 551–560. doi:10.1093/ejo/21.5.551.

Green-Thompson, N.F., 2007. Measuring and predicting Leeway space in themixed dentition on panoramic xrays using computer imaging analysis. *Thesis*. Johanesburg : University of the Witwatersrand. 1-35.

Haspeni., 2011. *Ukuran dan bentuk lengkung gigi pada mahasiswa suku batak FKG dan FT USU*. Skripsi FKG USU. Medan.

Harty, F.J., 1995. *Kamuskedokterangigi*. EGC, Jakarta . 13, 229.

Hartono, E., 2014. Karakteristik analisis sefalometri pada kasus impaksi molar ketiga mandibula. *Skripsi*. Universitas Hasanuddin. Makasar.

Ho, C.T., Freer, T.J., 1994. Clinical application of the graphical analysis of tooth width discrepancy. *Aust Orthod J*. 13(3):137-43.

Hong, Q., Tan, J., Koirala, R., Lina, Y., Shimizu, T., Nakano, K., 2008. A study of Bolton's and Pont's analysis on Permanent Dentition of Nepalese. *Journal of Hard Tissue Biology*. 17 (3) : 91-98.

Hendra, C. 2005. Analisis Hubungan Bentuk dan Ukuran Gigi Geligi, Bentuk dan Ukuran Lengkung Gigi Geligi Terhadap Terjadinya Impaksi Gigi Molar Tiga Rahang Bawah (Studi Epidemiologi pada Suku Bugis dan Suku Toraja). *Disertasi*. Universitas Hasanuddin: Makassar. 35-37.

Jacobs, S.G., Woods, M.G., 2000. Radiographic localization of unerupted teeth: Further findings about the vertical tube shift method and other localization techniques. *Am. J. Orthod. Dentofac. Orthop*. 118, 439–447. doi:10.1067/mod.2000.108782.

Jacoby, H., 1983. The etiology of maxillary canine impactions. *Am. J. Orthod.* 84, 125–132. doi:10.1016/0002-9416(83)90176-8.

Katsnelson, A., Flick, W.G., Susarla, S., Tartakovsky, J. V., Miloro, M., 2010. Use of Panoramic X-Ray to Determine Position of Impacted Maxillary Canines. *J. Oral Maxillofac. Surg*. 68, 996–1000. doi:10.1016/j.joms.2009.09.022.

Khalid, W.H., 2008. *Variations in tooth size, dental arch dimensions and shape among Malay school children.* Masters thesis, Universiti Sains Malaysia. 1(18): 1-24.

UNIVERSITAS
GADJAH MADA
**TINGKAT AKURASI METODE MAGNIFIKASI CANINE INCISOR INDEX DALAM PENENTUAN LETAK IMPAKSI KANINUS
MAKSILA PADA RADIOGRAF PANORAMIK RAS DEUTRO MELAYU**
ANTON, Drg. Rahardjo, S.U., Sp BM (K); drg. Bambang Dwiraharjo, Sp.BM (K)
Universitas Gadjah Mada, 2018 | Diunduh dari http://etd.repository.ugm.ac.id/

Kiliaridis, S., Georgiakaki, I., Katsaros, C., 2003. Masseter muscle thickness and maxillary dental arch width. *Eur J Orthod*. 25: 259-263.

Kumar, S., Mehrotra, P., Bhagchandani, J., Singh, A., Garg, A., Kumar, S., Sharma, A., Yadav, H., 2015. Localization of impacted canines. *J. Clin. Diagnostic Res.* 9, 11–14. doi:10.7860/JCDR/2015/10529.5480.

Kurol, J., 2002. Early treatment of tooth-eruption disturbances. *Am. J. Orthod. Dentofac. Orthop.* 121, 588–591. doi:10.1067/mod.2002.124173.

Langlais, R.P., Broadus, R., Glass, B.J., 1985. Bifid mandibular canals in panoramic radiographs. *J Am Dent Assoc* . 110(6): 925.

Leifert S, Jonas I E 2003 Dental anomalies as a microsymptom of palatal canine displacement. *Journal of Orofacial Orthopedics*. 64: 108–120.

Lindauer, S.J., Rubenstein, L.K., Hang, W.M., Anderson, W.C., Isaacson, R.J., 1992. Canine impaction early with panoramic radiographs.*Am J Dent Assoc.*123:917.

Ludicke, G., Harzer, W., Tausche, E., 2008. Incisor inclination--risk factor for palatally-impacted canines. *J. Orofac. Orthop*. 69, 357–364.

Mah, J., Enciso, R., Jorgensen, M., 2003. Management of impacted cuspids using 3-D volumetric imaging. *J. Calif. Dent. Assoc.* 31, 835–41.

Mason, C., Papadakou, P., Roberts, G.J., 2001. The radiographic localization of impacted maxillary canines: A comparison of methods. *Eur. J. Orthod.* 23, 25–34. doi:10.1093/ejo/23.1.25.

Margono, G., 1998. *Radiologi Intraoral : Teknik, Prosesing, Interpretasi Radiogram*, EGC, Jakarta.

Mehmet, B., Özer, M., Sener, I.,2007. Bilaterally Impacted Maxillary Central Incisors: Surgical Exposure and Orthodontic Treatment: A Case Report. *J Contemp Dent Pract* . (7)4:098-105.

Mudjosemedi, M., Widyaningrum, R., Gracea, S., 2015. Perbedaan Hasil Pengukuran Horizontal pada Tulang Mandibula dengan Radiograf Panoramik . *Majalah Kedokteran Gigi Indonesia.* 1 (1), 78–85.

Nagpal, A., Pai, K.M., Setty, S., Sharma, G., 2009. Localization of impacted maxillary canines using panoramic radiography. *J. Oral Sci.* 51, 37–45. doi:10.2334/josnusd.51.37.

Nishikawa, K., Suehiro, A., Sekine, H., Kousuge, Y., Wakoh, M., Sano, T., 2010.

UNIVERSITAS GADJAH MADA
TINGKAT AKURASI METODE MAGNIFIKASI CANINE INCISOR INDEX DALAM PENENTUAN LETAK IMPAKSI KANINUS MAKSILA PADA RADIOGRAF PANORAMIK RAS DEUTRO MELAYU
ANTON, Drg. Rahardjo, S.U., Sp BM (K); drg. Bambang Dwiraharjo, Sp.BM (K)
Universitas Gadjah Mada, 2018 | Diunduh dari http://etd.repository.ugm.ac.id/

Is linear distance measured by panoramic radiography reliable? *Oral Radiol.* 26, 16–19. doi:10.1007/s11282-010-0038-5.

Ostrofsky, M.K., 1976. Localization of impacted canines with Status-X radiography. *Oral Surgery, Oral Med. Oral Pathol.* 42, 529–533. doi:10.1016/0030-4220(76)90301-7.

Paramesthi, G.A.M.D.H., Farmasyanti, C.A., dan Karunia, D., 2011. Besar Indeks Pont dan Korkhaus Serta Hubungan Antara Lebar dan Panjang Lengkung Gigi terhadap Tinggi Palatum. *Maj Ked Gi.* 18(1) :6-10

Peck, S., Peck, L., Kataja, M., 1994. The palatally displaced canine as a dental anomaly of genetic origin. *Angle Orthod.* doi:10.1043/0003-3219(1994)064<0249:WNID>2.0.CO;2.

Peckmann, T.R., Logar, C., Garrido-Varas, C.E., Meek, S., Pinto, X.T., 2016. Sex determination using the mesio-distal dimension of permanent maxillary incisors and canines in a modern Chilean population. *Sci. Justice* 56, 84–89. doi:10.1016/j.scijus.2015.10.002.

Pirinen, S., Arte, S., Apajalahti, S., 1996. Palatal displacement of canine is genetic and related to congenital absence of teeth.*J Dent Res.*75(10):1742-6.

Pittayapat, P., Willems, G., Alqerban, A., Coucke, W., Ribeiro-Rotta, R.F., Souza, P.C., Westphalen, F.H., Jacobs, R., 2014. Agreement between cone beam computed tomography images and panoramic radiographs for initial orthodontic evaluation. *Oral Surg. Oral Med. Oral Pathol. Oral Radiol.* 117, 111–119. doi:10.1016/j.oooo.2013.10.016.

Pramanthi, S., 2007. Perbedaan Ukuran Maxillae dan Mandibulae antara Laki-laki dengan Wanita di Bagian Anatomi FakultasKedokteran Universitas Airlangga. *Jurnal PDGI*. 57 (1) : 26-29

Rachmini, S.L., 1990. Validitas ratio ukuran, mesiodistal ukuran gigi rahang bawah dan rahang atas menurut Bolton. *Tesis* Universitas Indonesia. Jakarta.

Rai, B., Anand, S.C., Madan, M., Dhattarwal, S., 2007. Criteria for determination of sex from mandible. *J Dent Sci*. 4(2): 1-9.

Rai, B., Anand, S.C., 2007. Gender determination by diagonal distances of teeth. *J Biological Anthropology*. 1(1): 1-8.

Richardson, G., Russell, K.A., 2000. A review of impacted permanent maxillary cuspids - Diagnosis and prevention. *J. Can. Dent. Assoc.* 66(9):497-501.

Rieuwpassa, I. E., Hamrun, N., Riksavianti, F., 2013.Ukuran mesiodistal dan

UNIVERSITAS
GADJAH MADA
**TINGKAT AKURASI METODE MAGNIFIKASI CANINE INCISOR INDEX DALAM PENENTUAN LETAK IMPAKSI KANINUS
MAKSILA PADA RADIOGRAF PANORAMIK RAS DEUTRO MELAYU**
ANTON, Drg. Rahardjo, S.U., Sp BM (K); drg. Bambang Dwiraharjo, Sp.BM (K)
Universitas Gadjah Mada, 2018 | Diunduh dari http://etd.repository.ugm.ac.id/

servikoinsisal gigi insisivus sentralis suku Bugis, Makassar, dan Toraja tidak menunjukkan perbedaan yang bermakna. *Dentofasial*, 12 (1) : 1-4.

Santoro, M., Ayoub, M.E., 2000. Mesiodistal crown dimensions and tooth size discrepancy of the permanent dentition of Dominican Americans. *Am J Orthod.* 70 : 303-07.

Shapira, Y., Kuftinec, M.M., 1998. Early diagnosis and interception of potential maxillary canine impaction. *J. Am. Dent. Assoc*. 129, 1450–4. doi:10.14219/jada.archive.1998.0080.

Sylvia, M., 2013. Hubungan lebar mesiodistal gigi dengan kecembungan profil jaringan lunak wajah pada mahasiswa FKG USU ras Deutromelayu. *Skripsi* FKG USU. Medan.

Sodhi, K.S., Krishna, S., Saxena, A.K., Sinha, A., Khandelwal, N., Lee, E.Y., 2015. Clinical application of "Justification" and "Optimization" principle of ALARA in pediatric CT imaging: "How many children can be protected from unnecessary radiation?" *Eur. J. Radiol.* 84, 1752–1757. doi:10.1016/j.ejrad.2015.05.030.

Stivaros, N., Mandall, N.A., 2000. Radiographic factors affecting the management of impacted upper permanent canines. *J. Orthod.* 27, 169–173. doi:10.1093/ortho/27.2.169.

Sudjana, 1996. *Teknik Analisis Regresi Dan Korelasi*. Tarsito: Bandung.

Suri, S., Utreja, A., Rattan, V., 2004. Orthodontic treatment of bilaterally impacted maxillary canines in an adult. *Am. J. Orthod. Dentofac. Orthop.* 122, 429–437. doi:10.1067/mod.2002.126406.

Swasono, S., Mieke, S.M., Susilowati., 2004. Variasi normal lebar mesiodistal gigi pada orang Bugis dan Toraja. *Dent. Journal.* 37 (1) : 1-3.

Thilander, B dan Myrberg, N., 1973. The prevalence of malocclusion in Swedish schoolchildren, *Scand J Dent Res.*, 81:12.

Van der Linden, F.P.G.M. *Treatment of impacted canines in orthodonthic consepts and strategies*, Quintessence publishing Co, Ltd., London 2000; 161-176.

Veli, I., Yuksel, B., Uysal, T., 2014. Prevalence of Maxillary Permanent Canine Impaction in Relation to Anomalous Lateral Incisors. *Turkish J.* Orthod. 27, 90–99. doi:10.13076/TJO-D-14-00025.

Warford, J.H., Grandhi, R.K., Tira, D.E., 2003. Prediction of maxillary canine impaction using sectors and angular measurement. *Am. J. Orthod. Dentofac.*

UNIVERSITAS
GADJAH MADA
**TINGKAT AKURASI METODE MAGNIFIKASI CANINE INCISOR INDEX DALAM PENENTUAN LETAK IMPAKSI KANINUS
MAKSILA PADA RADIOGRAF PANORAMIK RAS DEUTRO MELAYU**
ANTON, Drg. Rahardjo, S.U., Sp BM (K); drg. Bambang Dwiraharjo, Sp.BM (K)
Universitas Gadjah Mada, 2018 | Diunduh dari http://etd.repository.ugm.ac.id/

Orthop. 124, 651–655. doi:10.1016/S0889-5406(03)00621-8.

Whaites, E., 2013. *Essential of Dental Radiography and Radiology*, 5th ed. Churchill Livingstone,Elsevier . 137-141, 147-9, 151-165.

White, S.C., Pharoah, M.J., 2009. *Oral radiology: Principles and Interpretation*, 6th ed. St Louis: Elsevier. 161-186.

Wolf, J.E., Mattila, K., 1979. Localization of impacted maxillary canines by panoramic tomography. *Dentomaxillofacial Radiol*. 8, 85–91. doi:10.1259/dmfr.1979.0011.

Yacoob, H., Narnbar, P., Naidu, M.D.K., 1996. Racial Characteristic of Human Teeth with Special Emphasis on the Mongoloid Dentition. *Malaysian J. Pathol*. 18 : 1-7

Yavuz, M.S., Aras, M.H., Büyükkurt, M.C., Tozoglu, S., 2007. Impacted mandibular canines. *J. Contemp. Dent. Pract.* 8, 078–085. doi:10.1016/j.joms.2006.02.033.