

## References

- Adcock, R. A., Dale, C., Fisher, M., Aldebot, S., Genevsky, A., Simpson, G. V., Nagarajan, A., Vinogradov, S. (2009). When top-down meets bottom-up: auditory training enhances verbal memory in Schizophrenia. *Schizophrenia Bulletin*, 35(6), 1132–1141. <https://doi.org/10.1093/schbul/sbp068>
- Almeida, C. P. B. de, Skupien, E. C., & Silva, D. R. (2015). Health care seeking behavior and patient delay in tuberculosis diagnosis. *Cadernos de Saúde Pública*, 31(2), 321–330. <https://doi.org/10.1590/0102-311X00195413>
- American Psychiatric Association (Ed.). (2013). *Diagnostic and statistical manual of mental disorders: DSM-5* (5th ed). Washington, D.C: American Psychiatric Association.
- Azwar, S. (2004). *Penyusunan skala psikologi* (6th ed.). Yogyakarta: Pustaka Pelajar.
- Badan Penelitian Dan Pengembangan Kesehatan. (2013). *Riset kesehatan dasar* (Riskesdas). Jakarta: Kementerian Kesehatan.
- Belge, J.-B., Maurage, P., Manginckx, C., Leleux, D., Delatte, B., & Constant, E. (2017). Facial decoding in schizophrenia is underpinned by basic visual processing impairments. *Psychiatry Research*. <https://doi.org/10.1016/j.psychres.2017.04.007>
- Butler, P. D., & Javitt, D. C. (2005). Early-stage visual processing deficits in schizophrenia. *Current Opinion in Psychiatry*, 18(2), 151–157.
- Carbon, C.-C. (2014). Understanding human perception by human-made illusions. *Frontiers in Human Neuroscience*, 8. <https://doi.org/10.3389/fnhum.2014.00566>
- Chaumon, M., Kveraga, K., Barrett, L. F., & Bar, M. (2014). Visual predictions in the orbitofrontal cortex rely on associative content. *Cerebral Cortex (New York, NY)*, 24(11), 2899–2907. <https://doi.org/10.1093/cercor/bht146>
- Cook, J., Barbalat, G., & Blakemore, S.-J. (2012). Top-down modulation of the perception of other people in schizophrenia and autism. *Frontiers in Human Neuroscience*, 6. <https://doi.org/10.3389/fnhum.2012.00175>
- Corbett, B. A., Newsom, C., Key, A. P., Qualls, L. R., & Edmiston, E. K. (2014). Examining the relationship between face processing and social interaction behavior in children with and without autism spectrum disorder. *Journal of Neurodevelopmental Disorders*, 6(1), 35.
- Dima, D., Dietrich, D. E., Dillo, W., & Emrich, H. M. (2010). Impaired top-down processes in schizophrenia: A DCM study of ERPs. *NeuroImage*, 52(3), 824–832. <https://doi.org/10.1016/j.neuroimage.2009.12.086>
- Dima, D., Dillo, W., Bonnemann, C., Emrich, H. M., & Dietrich, D. E. (2011). Reduced P300 and P600 amplitude in the hollow-mask illusion in patients with schizophrenia. *Psychiatry Research: Neuroimaging*, 191(2), 145–151. <https://doi.org/10.1016/j.psychresns.2010.09.015>
- Dima, D., Roiser, J. P., Dietrich, D. E., Bonnemann, C., Lanfermann, H., Emrich, H. M., & Dillo, W. (2009). Understanding why patients with schizophrenia

- do not perceive the hollow-mask illusion using dynamic causal modelling. *NeuroImage*, 46(4), 1180–1186. <https://doi.org/10.1016/j.neuroimage.2009.03.033>
- Elgar, K., & Campbell, R. (2001). Annotation: the cognitive neuroscience of face recognition: implications for developmental disorders. *Journal of Child Psychology and Psychiatry*, 42(6), 705–717. <https://doi.org/10.1017/S0021963001007508>
- Farah, M. J., Wilson, K. D., Drain, H. M., & Tanaka, J. R. (1995). The inverted face inversion effect in prosopagnosia: evidence for mandatory, face-specific perceptual mechanisms. *Vision Research*, 35(14), 2089–2093.
- Friston, K. J., Harrison, L., & Penny, W. (2003). Dynamic causal modelling. *NeuroImage*, 19(4), 1273–1302. [https://doi.org/10.1016/S1053-8119\(03\)00202-7](https://doi.org/10.1016/S1053-8119(03)00202-7)
- Gilbert, C. D., & Sigman, M. (2007). Brain states: top-down influences in sensory processing. *Neuron*, 54(5), 677–696. <https://doi.org/10.1016/j.neuron.2007.05.019>
- Girma, E., & Tesfaye, M. (2011). Patterns of treatment seeking behavior for mental illnesses in Southwest Ethiopia: a hospital based study. *BMC Psychiatry*, 11, 138. <https://doi.org/10.1186/1471-244X-11-138>
- Glabus, M. F. (2005). *Neuroimaging*. Elsevier Science. Retrieved from <https://books.google.co.id/books?id=d2LmMLTsIJMC>
- Gold, J. M., Fuller, R. L., Robinson, B. M., Braun, E. L., & Luck, S. J. (2007). Impaired top-down control of visual search in schizophrenia. *Schizophrenia Research*, 94(1–3), 148–155.
- Gross, C. (1992). Representation of visual stimuli in inferior temporal cortex. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 335(1273), 3–10. <https://doi.org/10.1098/rstb.1992.0001>
- Hauser, L. (n.d.). Spatial frequency 1. Retrieved April 26, 2018, from <http://www.ucalgary.ca/pip369/mod4/spatial/frequency1>
- Healy, D. (2008). *Psychiatric drugs explained E-book*. Elsevier Health Sciences. Retrieved from <https://books.google.co.id/books?id=xurNluX8g7IC>
- Hebart, M. N., & Hesselmann, G. (2012). What visual information is processed in the human dorsal stream? *Journal of Neuroscience*, 32(24), 8107–8109. <https://doi.org/10.1523/JNEUROSCI.1462-12.2012>
- Jenkins, N. (2016, March 21). Thousands of mentally ill Indonesians live in chains: report. Retrieved September 15, 2017, from <http://time.com/4265623/indonesia-mental-illness-chains-pasung-hrw/>
- Katsuki, F., & Constantinidis, C. (2014). Bottom-up and top-down attention: different processes and overlapping neural systems. *The Neuroscientist*, 20(5), 509–521. <https://doi.org/10.1177/1073858413514136>
- Keane, B. P., Silverstein, S. M., Wang, Y., & Papathomas, T. V. (2013). Reduced depth inversion illusions in schizophrenia are state-specific and occur for multiple object types and viewing conditions. *Journal of Abnormal Psychology*, 122(2), 506–512. <https://doi.org/10.1037/a0032110>

- Khakbaban, Z., Taghipour, A., Latifnejad Roudsari, R., & Mohammadi, E. (2014). Help seeking behavior of women with self-discovered breast cancer symptoms: a meta-ethnographic synthesis of patient delay. *PLoS ONE*, 9(12). <https://doi.org/10.1371/journal.pone.0110262>
- Kveraga, K., Boshyan, J., & Bar, M. (2007). Magnocellular projections as the trigger of top-down facilitation in recognition. *Journal of Neuroscience*, 27(48), 13232–13240. <https://doi.org/10.1523/JNEUROSCI.3481-07.2007>
- Lee, C. U., Shenton, M. E., Salisbury, D. F., Kasai, K., Onitsuka, T., Dickey, C. C., ... McCarley, R. W. (2002). Fusiform gyrus volume reduction in first-episode schizophrenia: a magnetic resonance imaging study. *Archives of General Psychiatry*, 59(9), 775–781.
- Leopold, D. A., & Rhodes, G. (2010). A comparative view of face perception. *Journal of Comparative Psychology (Washington, D.C. : 1983)*, 124(3), 233–251. <https://doi.org/10.1037/a0019460>
- Lestari, W., & Wardhani, Y. F. (2014). Stigma and management on people with severe mental disorders with “Pasung” (physical restraint). *Buletin Penelitian Sistem Kesehatan*, 17(2), 10.
- Levitt, J. M., Saka, N., Hunter Romanelli, L., & Hoagwood, K. (2007). Early identification of mental health problems in schools: The status of instrumentation. *Journal of School Psychology*, 45(2), 163–191. <https://doi.org/10.1016/j.jsp.2006.11.005>
- Li, J., Liu, J., Liang, J., Zhang, H., Zhao, J., Huber, D. E., ... Shi, G. (2009). A distributed neural system for top-down face processing. *Neuroscience Letters*, 451(1), 6–10. <https://doi.org/10.1016/j.neulet.2008.12.039>
- Liu, C.-S. J., Bryan, R. N., Miki, A., Woo, J. H., Liu, G. T., & Elliott, M. A. (2006). Magnocellular and parvocellular visual pathways have different blood oxygen level-dependent signal time courses in human primary visual cortex. *American Journal of Neuroradiology*, 27(8), 1628–1634.
- Lokuge, K., Caleo, G., Greig, J., Duncombe, J., McWilliam, N., Squire, J., ... others. (2016). Successful control of Ebola Virus disease: analysis of service based data from rural Sierra Leone. *PLoS Neglected Tropical Diseases*, 10(3), e0004498.
- Lusia Kus Ana. (2012, February 11). Gangguan jiwa masih diabaikan. Retrieved May 27, 2018, from <https://lifestyle.kompas.com/read/2012/02/11/07363466/Gangguan.Jiwa.Masih.Diabaikan>
- Martínez, A., Hillyard, S. A., Dias, E. C., Hagler, D. J., Butler, P. D., Guilfoyle, D. N., Jalbrzikowski, M., Silipo, G., Javitt, D. C. (2008). Magnocellular pathway impairment in schizophrenia: evidence from Functional Magnetic Resonance Imaging. *Journal of Neuroscience*, 28(30), 7492–7500. <https://doi.org/10.1523/JNEUROSCI.1852-08.2008>
- Maslim, R. (2013). *Diagnosis gangguan jiwa, rujukan ringkas dari PPDGJ-III dan DSM-5*. Jakarta: Bagian Ilmu Kedokteran Jiwa FK-Unika Atmajaya.
- McDonald, B., Highley, J. R., Walker, M. A., Herron, B. M., Cooper, S. J., Esiri, M. M., & Crow, T. J. (2000). Anomalous Asymmetry of Fusiform and

- Parahippocampal Gyrus Gray Matter in Schizophrenia: A Postmortem Study. *American Journal of Psychiatry*, 157(1), 40–47.  
<https://doi.org/10.1176/ajp.157.1.40>
- Mishra, N., Nagpal, S. S., Chadda, R. K., & Sood, M. (2011). Help-seeking behavior of patients with mental health problems visiting a tertiary care center in north India. *Indian Journal of Psychiatry*, 53(3), 234–238.  
<https://doi.org/10.4103/0019-5545.86814>
- Muhana, S. U., Partosuwido, S. R., Ul Hasanat, N., & Ramdhani, N. (1999). *Skala kepribadian Universitas Gadjah Mada: alat diagnosis gangguan psikologis masyarakat Indonesia*. Lembaga Penelitian Universitas Gadjah Mada.
- Nestor, A., Vettel, J. M., & Tarr, M. J. (2008). Task-Specific Codes for Face Recognition: How they Shape the Neural Representation of Features for Detection and Individuation. *PLoS ONE*, 3(12), e3978.  
<https://doi.org/10.1371/journal.pone.0003978>
- Onitsuka, T., Shenton, M. E., Kasai, K., Nestor, P. G., Toner, S. K., Kikinis, R., ... McCarley, R. W. (2003). Fusiform gyrus volume reduction and facial recognition in chronic schizophrenia. *Archives of General Psychiatry*, 60(4), 349–355.
- Prasetyo, N. H., & Subandi, M. A. (2014). Program Intervensi Narimo Ing Pandum untuk Meningkatkan Kesejahteraan Psikologis Keluarga Pasien Skizofrenia. *JIP: Jurnal Intervensi Psikologi*, 6(2).
- Saygin, Z. M., Osher, D. E., Koldewyn, K., Reynolds, G., Gabrieli, J. D. E., & Saxe, R. R. (2011). Anatomical connectivity patterns predict face selectivity in the fusiform gyrus. *Nature Neuroscience*, 15(2), 321–327.  
<https://doi.org/10.1038/nn.3001>
- Schwarzlose, R. F. (2005). Separate Face and Body Selectivity on the Fusiform Gyrus. *Journal of Neuroscience*, 25(47), 11055–11059.  
<https://doi.org/10.1523/JNEUROSCI.2621-05.2005>
- Seiferth, N. Y., Pauly, K., Kellermann, T., Shah, N. J., Ott, G., Herpertz-Dahlmann, B., ... Habel, U. (2009). Neuronal correlates of facial emotion discrimination in early onset schizophrenia. *Neuropsychopharmacology*, 34(2), 477–487.
- Sembiring, S., & Sismudjito. (2015). Pengetahuan dan pemanfaatan metode pengobatan tradisional pada masyarakat Desa Suka Nalu Kecamatan Barus Jahe. *Perspektif Sosiologi*, 3(1), 14.
- Silverstein, S. M., All, S. D., Kasi, R., Berten, S., Essex, B., Lathrop, K. L., & Little, D. M. (2010). Increased fusiform area activation in schizophrenia during processing of spatial frequency-degraded faces, as revealed by fMRI. *Psychological Medicine*, 40(07), 1159–1169.  
<https://doi.org/10.1017/S0033291709991735>
- Subandi, M. A. (2008). Ngemong: dimensi keluarga pasien psikotik di Jawa. *Jurnal Psikologi*, 35(1), 62–79.
- Suharto, B. (2014). Budaya pasung dan dampak yuridis sosiologis (studi tentang upaya pelepasan pasung dan pencegahan tindakan pemasungan di Kabupaten Wonogiri). *Indonesian Journal on Medical Science*, 1(2), 1–10.

- Sussman, T. J., Jin, J., & Mohanty, A. (2016). Top-down and bottom-up factors in threat-related perception and attention in anxiety. *Biological Psychology*, 121, 160–172. <https://doi.org/10.1016/j.biopsycho.2016.08.006>
- Suyasa, I. G. P., Krisnandari, A. I. W., Onajiati, N. W., & Diyu, I. A. N. P. (2014). Keluhan-keluhan lanjut usia yang datang ke pengobatan gratis di salah satu wilayah pedesaan di Bali. In *Keluhan-Keluhan Lanjut Usia yang Datang Ke Pengobatan Gratis di Salah Satu Wilayah Pedesaan di Bali* (p. 7). Denpasar.
- Takahashi, T., Zhou, S.-Y., Nakamura, K., Tanino, R., Furuichi, A., Kido, M., Kurachi, M., Suzuki, M. (2011). A follow-up MRI study of the fusiform gyrus and middle and inferior temporal gyri in schizophrenia spectrum. *Progress in Neuro-Psychopharmacology and Biological Psychiatry*, 35(8), 1957–1964. <https://doi.org/10.1016/j.pnpbp.2011.07.009>
- Tanaka, E., Inui, K., Kida, T., & Kakigi, R. (2009). Common cortical responses evoked by appearance, disappearance and change of the human face. *BMC Neuroscience*, 10(1), 38. <https://doi.org/10.1186/1471-2202-10-38>
- Trepanier, M., Hopperton, K., Mizrahi, R., Mechawar, N., & Bazinet, R. (2016). Postmortem evidence of cerebral inflammation in schizophrenia: a systematic review. *Molecular Psychiatry*.
- Walther, S., Federspiel, A., Horn, H., Bianchi, P., Wiest, R., Wirth, M., Strik, W., Müller, T. J. (2009). Encoding deficit during face processing within the right fusiform face area in schizophrenia. *Psychiatry Research: Neuroimaging*, 172(3), 184–191. <https://doi.org/10.1016/j.pscychresns.2008.07.009>
- WHO. (2016, March 29). WHO | Statement on the 9th meeting of the IHR Emergency Committee regarding the Ebola outbreak in West Africa. Retrieved September 15, 2017, from <http://www.who.int/mediacentre/news/statements/2016/end-of-ebola-pheic/en/>
- Wichowicz, H., Ciszewski, S., Żuk, K., & Rybak-Korneluk, A. (2016). Hollow mask illusion - is it really a test for schizophrenia? *Psychiatria Polska*, 50(4), 741–745. <https://doi.org/10.12740/PP/60150>
- Widianto, S. (2016, April 14). 'Pasung' Is the Indonesian practice of shackling and imprisoning the mentally ill. Retrieved September 15, 2017, from [https://www.vice.com/en\\_au/article/nknx8/pasung-indonesian-shackling-imprisoning-suffering-mental-illness-schizophrenia](https://www.vice.com/en_au/article/nknx8/pasung-indonesian-shackling-imprisoning-suffering-mental-illness-schizophrenia)
- Yoon, J. H., D'Esposito, M., & Carter, C. S. (2006). Preserved function of the fusiform face area in schizophrenia as revealed by fMRI. *Psychiatry Research: Neuroimaging*, 148(2–3), 205–216. <https://doi.org/10.1016/j.pscychresns.2006.06.002>
- Zhang, J., Li, X., Song, Y., & Liu, J. (2012). The Fusiform Face Area Is Engaged in Holistic, Not Parts-Based, Representation of Faces. *PLoS ONE*, 7(7), e40390. <https://doi.org/10.1371/journal.pone.0040390>
- Zhen, Z., Fang, H., & Liu, J. (2013). The hierarchical brain network for face recognition. *PLoS ONE*, 8(3), e59886. <https://doi.org/10.1371/journal.pone.0059886>



UNIVERSITAS  
GADJAH MADA

**Resistance toward Hollow Mask Illusion as a Feature for Schizophrenia Spectrum Disorder**

WILLY KRISTIAN TO Y, Supra Wimbarti, MSc., Ph.D.

Universitas Gadjah Mada, 2018 | Diunduh dari <http://etd.repository.ugm.ac.id/>

Zillmer, E., Spiers, M., & Culbertson, W. C. (2008). *Principles of neuropsychology* (2nd ed). Australia ; United States: Thomson/Wadsworth.