

DAFTAR PUSTAKA

- Antonsson, B., Conti, F., Ciavatta, A., Montessuit, S., Lewis, S., Martinou, I., Bernasconi, L., Bernard, A., Mermod, J.J., Mazzei, G., Maundrell, K., Gambale, F., Sadoul, R., Martinou, J.C., 1997. Inhibition of Bax channel-forming activity by Bcl-2. *Science* 277, 370–372.
- Astuti, P. 2010. Peranan asetosal sebagai anti-trombotik terhadap metabolisme tromboksan A2 (TXA2) dan prostasiklin (PGI2). *Stigmatognatic*. 1(7): 51-55.
- Bakhda, R.N., 2016. Clinical study of fundus findings in pregnancy induced hypertension. *J. Fam. Med. Prim. Care* 5, 424–429. <https://doi.org/10.4103/2249-4863.192364>
- Bakhshi, A., Jensen, J.P., Goldman, P., Wright, J.J., McBride, O.W., Epstein, A.L., Korsmeyer, S.J., 1985. Cloning the chromosomal breakpoint of t(14;18) human lymphomas: clustering around JH on chromosome 14 and near a transcriptional unit on 18. *Cell* 41, 899–906.
- Cadavid, A.P., 2017. Aspirin: The Mechanism of Action Revisited in the Context of Pregnancy Complications. *Front. Immunol.* 8, 261. <https://doi.org/10.3389/fimmu.2017.00261>
- Chinnaiyan, A.M., O'Rourke, K., Lane, B.R., Dixit, V.M., 1997. Interaction of CED-4 with CED-3 and CED-9: a molecular framework for cell death. *Science* 275, 1122–1126.
- Conradt, B., Horvitz, H.R., 1998. The C. elegans protein EGL-1 is required for programmed cell death and interacts with the Bcl-2-like protein CED-9. *Cell* 93, 519–529.
- Dejana, E., Orsenigo, F., Lampugnani, M.G., 2008. The role of adherens junctions and VE-cadherin in the control of vascular permeability. *J. Cell Sci.* 121, 2115–2122. <https://doi.org/10.1242/jcs.017897>
- Facca, T., Kirsztajn, G.M., Pereira, A., Moreira, S., Teixeira, V., Nishida, S., Sass, N., 2012. Renal Evaluation in Women with Preeclampsia. *Nephron Extra* 2, 125–132. <https://doi.org/10.1159/000338271>
- Gavard, J., 2013. Endothelial permeability and VE-cadherin: a wacky comradeship. *Cell Adhes. Migr.* 7, 455–461. <https://doi.org/10.4161/cam.27330>
- Gerber, H.P., Dixit, V., Ferrara, N., 1998a. Vascular endothelial growth factor induces expression of the antiapoptotic proteins Bcl-2 and A1 in vascular endothelial cells. *J. Biol. Chem.* 273, 13313–13316.
- Gerber, H.P., McMurtrey, A., Kowalski, J., Yan, M., Keyt, B.A., Dixit, V., Ferrara, N., 1998b. Vascular endothelial growth factor regulates endothelial cell survival through the phosphatidylinositol 3'-kinase/Akt signal transduction pathway. Requirement for Flk-1/KDR activation. *J. Biol. Chem.* 273, 30336–30343.
- Hashemi, M., 2014. The Study of Pentoxifylline Drug Effects on Renal Apoptosis and BCL-2 Gene Expression Changes Following Ischemic Reperfusion Injury in Rat. *Iran. J. Pharm. Res. IJPR* 13, 181–189.

- Hladunewich, M., Karumanchi, S.A., Lafayette, R., 2007. Pathophysiology of the clinical manifestations of preeclampsia. *Clin. J. Am. Soc. Nephrol.* CJASN 2, 543–549. <https://doi.org/10.2215/CJN.03761106>
- Horvath, B., Hegedus, D., Szapary, L., Marton, Z., Alexy, T., Koltai, K., Czopf, L., Wittmann, I., Juricskay, I., Toth, K., Kesmarky, G., 2004. Measurement of von Willebrand factor as the marker of endothelial dysfunction in vascular diseases. *Exp. Clin. Cardiol.* 9, 31–34.
- Huang, D.C., Adams, J.M., Cory, S., 1998. The conserved N-terminal BH4 domain of Bcl-2 homologues is essential for inhibition of apoptosis and interaction with CED-4. *EMBO J.* 17, 1029–1039. <https://doi.org/10.1093/emboj/17.4.1029>
- Hunter, J.J., Parslow, T.G., 1996. A peptide sequence from Bax that converts Bcl-2 into an activator of apoptosis. *J. Biol. Chem.* 271, 8521–8524.
- Jeyabalan, A., 2013. Epidemiology of preeclampsia: Impact of obesity. *Nutr. Rev.* 71, 10.1111/nure.12055. <https://doi.org/10.1111/nure.12055>
- Karumanchi, S.A., Maynard, S.E., Stillman, I.E., Epstein, F.H., Sukhatme, V.P., 2005. Preeclampsia: a renal perspective. *Kidney Int.* 67, 2101–2113. <https://doi.org/10.1111/j.1523-1755.2005.00316.x>
- Keikkala, E., Koskinen, S., Vuorela, P., Laivuori, H., Romppanen, J., Heinonen, S., Stenman, U.-H., 2016. First trimester serum placental growth factor and hyperglycosylated human chorionic gonadotropin are associated with pre-eclampsia: a case control study. *BMC Pregnancy Childbirth* 16, 378. <https://doi.org/10.1186/s12884-016-1169-4>
- Knudson, C.M., Korsmeyer, S.J., 1997. Bcl-2 and Bax function independently to regulate cell death. *Nat. Genet.* 16, 358–363. <https://doi.org/10.1038/ng0897-358>
- Kopincová, J., Púzserová, A., Bernátová, I., 2012. L-NAME in the cardiovascular system - Nitric oxide synthase activator? *Pharmacol. Rep.* 64, 511–520. [https://doi.org/10.1016/S1734-1140\(12\)70846-0](https://doi.org/10.1016/S1734-1140(12)70846-0)
- LeFevre, M.L., U.S. Preventive Services Task Force, 2014. Low-dose aspirin use for the prevention of morbidity and mortality from preeclampsia: U.S. Preventive Services Task Force recommendation statement. *Ann. Intern. Med.* 161, 819–826. <https://doi.org/10.7326/M14-1884>
- Li, P., Nijhawan, D., Budihardjo, I., Srinivasula, S.M., Ahmad, M., Alnemri, E.S., Wang, X., 1997. Cytochrome c and dATP-dependent formation of Apaf-1/caspase-9 complex initiates an apoptotic protease cascade. *Cell* 91, 479–489.
- Liu, X., Kim, C.N., Yang, J., Jemmerson, R., Wang, X., 1996. Induction of apoptotic program in cell-free extracts: requirement for dATP and cytochrome c. *Cell* 86, 147–157.
- Maynard, S.E., Min, J.-Y., Merchan, J., Lim, K.-H., Li, J., Mondal, S., Libermann, T.A., Morgan, J.P., Sellke, F.W., Stillman, I.E., Epstein, F.H., Sukhatme, V.P., Karumanchi, S.A., 2003. Excess placental soluble fms-like tyrosine kinase 1 (sFlt1) may contribute to endothelial dysfunction, hypertension, and proteinuria in preeclampsia. *J. Clin. Invest.* 111, 649–658. <https://doi.org/10.1172/JCI200317189>

- Maynard, S.E., Venkatesha, S., Thadhani, R., Karumanchi, S.A., 2005. Soluble Fms-like tyrosine kinase 1 and endothelial dysfunction in the pathogenesis of preeclampsia. *Pediatr. Res.* 57, 1R–7R. <https://doi.org/10.1203/01.PDR.0000159567.85157.B7>
- Moore, K.L., Dalley, A.F., Agur, A.M.R., 2014. Clinically oriented anatomy, 7th ed. ed. Wolters Kluwer/Lippincott Williams & Wilkins Health, Philadelphia.
- Motta, C., Grosso, C., Zanuzzi, C., Molinero, D., Picco, N., Bellingeri, R., Alustiza, F., Barbeito, C., Vivas, A., Romanini, M.C., 2015. Effect of Sildenafil on Pre-Eclampsia-Like Mouse Model Induced By L-Name. *Reprod. Domest. Anim.* 50, 611–616. <https://doi.org/10.1111/rda.12536>
- Müller-Deile, J., Schiffer, M., 2011. Renal Involvement in Preeclampsia: Similarities to VEGF Ablation Therapy. *J. Pregnancy* 2011, 176973. <https://doi.org/10.1155/2011/176973>
- Murohara, T., Horowitz, J.R., Silver, M., Tsurumi, Y., Chen, D., Sullivan, A., Isner, J.M., 1998. Vascular endothelial growth factor/vascular permeability factor enhances vascular permeability via nitric oxide and prostacyclin. *Circulation* 97, 99–107.
- Oltvai, Z.N., Milliman, C.L., Korsmeyer, S.J., 1993. Bcl-2 heterodimerizes in vivo with a conserved homolog, Bax, that accelerates programmed cell death. *Cell* 74, 609–619.
- Osungbade, K.O., Ige, O.K., 2011. Public Health Perspectives of Preeclampsia in Developing Countries: Implication for Health System Strengthening. *J. Pregnancy* 2011, 481095. <https://doi.org/10.1155/2011/481095>
- Parker, R.I., Gralnick, H.R., 1989. Effect of aspirin on platelet-von Willebrand factor surface expression on thrombin and ADP-stimulated platelets. *Blood* 74, 2016–2021.
- Rachmi, A.S. 2016. Aspirin dosis rendah efektif menurunkan resistensi arteri uterina yang abnormal pada ibu hamil usia kehamilan 16–24 minggu. *Majalah Obstetri & Ginekologi.* 1(24): 25-30.
- Ribeiro, M.O., Antunes, E., de Nucci, G., Lovisolo, S.M., Zatz, R., 1992. Chronic inhibition of nitric oxide synthesis. A new model of arterial hypertension. *Hypertens. Dallas Tex* 1979 20, 298–303.
- Roberts, J.M., Taylor, R.N., Musci, T.J., Rodgers, G.M., Hubel, C.A., McLaughlin, M.K., 1989. Preeclampsia: an endothelial cell disorder. *Am. J. Obstet. Gynecol.* 161, 1200–1204.
- Rudel, T., 1999. Caspase inhibitors in prevention of apoptosis. *Herz* 24, 236–241.
- Sadek, S.A., Rashed, L.A., Bassam, A.M., Said, E.S., 2014. Effect of aliskiren, telmisartan and torsemide on cardiac dysfunction in l-nitro arginine methyl ester (l-NAME) induced hypertension in rats. *J. Adv. Res.* 6, 967–974. <https://doi.org/10.1016/j.jare.2014.11.003>
- Salakou, S., Kardamakis, D., Tsamandas, A.C., Zolota, V., Apostolakis, E., Tzelepi, V., Papathanasopoulos, P., Bonikos, D.S., Papapetropoulos, T., Petsas, T., Dougenis, D., 2007. Increased Bax/Bcl-2 ratio up-regulates caspase-3 and increases apoptosis in the thymus of patients with myasthenia gravis. *Vivo Athens Greece* 21, 123–132.

- Schiff, E., Peleg, E., Goldenberg, M., Rosenthal, T., Ruppin, E., Tamarkin, M., Barkai, G., Ben-Baruch, G., Yahal, I., Blankstein, J., 1989. The use of aspirin to prevent pregnancy-induced hypertension and lower the ratio of thromboxane A2 to prostacyclin in relatively high risk pregnancies. *N. Engl. J. Med.* 321, 351–356. <https://doi.org/10.1056/NEJM198908103210603>
- Schrör, K., 1997. Aspirin and platelets: the antiplatelet action of aspirin and its role in thrombosis treatment and prophylaxis. *Semin. Thromb. Hemost.* 23, 349–356. <https://doi.org/10.1055/s-2007-996108>
- Susin, S.A., Zamzami, N., Castedo, M., Daugas, E., Wang, H.G., Geley, S., Fassy, F., Reed, J.C., Kroemer, G., 1997. The central executioner of apoptosis: multiple connections between protease activation and mitochondria in Fas/APO-1/CD95- and ceramide-induced apoptosis. *J. Exp. Med.* 186, 25–37.
- Thakker, G.D., Hajjar, D.P., Muller, W.A., Rosengart, T.K., 1999. The role of phosphatidylinositol 3-kinase in vascular endothelial growth factor signaling. *J. Biol. Chem.* 274, 10002–10007.
- Tortora, G.J., Derrickson, B., 2012. Principles of anatomy & physiology, 13th ed. ed. Wiley, Hoboken, NJ.
- Tsujimoto, Y., 1998. Role of Bcl-2 family proteins in apoptosis: apoptosomes or mitochondria? *Genes Cells Devoted Mol. Cell. Mech.* 3, 697–707.
- Tsurumi, Y., Murohara, T., Krasinski, K., Chen, D., Witzenbichler, B., Kearney, M., Couffinhal, T., Isner, J.M., 1997. Reciprocal relation between VEGF and NO in the regulation of endothelial integrity. *Nat. Med.* 3, 879–886.
- Uzan, J., Carbonnel, M., Piconne, O., Asmar, R., Ayoubi, J.-M., 2011. Preeclampsia: pathophysiology, diagnosis, and management. *Vasc. Health Risk Manag.* 7, 467–474. <https://doi.org/10.2147/VHRM.S20181>
- van den Oever, I.A.M., Raterman, H.G., Nurmohamed, M.T., Simsek, S., 2010. Endothelial Dysfunction, Inflammation, and Apoptosis in Diabetes Mellitus. *Mediators Inflamm.* 2010, 792393. <https://doi.org/10.1155/2010/792393>
- van der Graaf, A.M., Toering, T.J., Faas, M.M., Lely, A.T., 2012. From preeclampsia to renal disease: a role of angiogenic factors and the renin-angiotensin aldosterone system? *Nephrol. Dial. Transplant. Off. Publ. Eur. Dial. Transpl. Assoc. - Eur. Ren. Assoc.* 27 Suppl 3, iii51-57. <https://doi.org/10.1093/ndt/gfs278>
- Vane, J.R., Botting, R.M., 2003. The mechanism of action of aspirin. *Thromb. Res.* 110, 255–258.
- Vestweber, D., 2008. VE-cadherin: the major endothelial adhesion molecule controlling cellular junctions and blood vessel formation. *Arterioscler. Thromb. Vasc. Biol.* 28, 223–232. <https://doi.org/10.1161/ATVBAHA.107.158014>
- Walsh, S.W., 1985. Preeclampsia: an imbalance in placental prostacyclin and thromboxane production. *Am. J. Obstet. Gynecol.* 152, 335–340.
- Warner, T.D., Nylander, S., Whatling, C., 2011. Anti-platelet therapy: cyclooxygenase inhibition and the use of aspirin with particular regard to dual

- anti-platelet therapy. *Br. J. Clin. Pharmacol.* 72, 619–633. <https://doi.org/10.1111/j.1365-2125.2011.03943.x>
- Yang, B., Johnson, T.S., Thomas, G.L., Watson, P.F., Wagner, B., Skill, N.J., Haylor, J.L., El Nahas, A.M., 2001. Expression of apoptosis-related genes and proteins in experimental chronic renal scarring. *J. Am. Soc. Nephrol.* JASN 12, 275–288.
- Yang, E., Korsmeyer, S.J., 1996. Molecular thanatopsis: a discourse on the BCL2 family and cell death. *Blood* 88, 386–401.
- Ylikorkala, O., Mäkilä, U.M., 1985. Prostacyclin and thromboxane in gynecology and obstetrics. *Am. J. Obstet. Gynecol.* 152, 318–329.
- Zachary, I., Gliki, G., 2001. Signaling transduction mechanisms mediating biological actions of the vascular endothelial growth factor family. *Cardiovasc. Res.* 49, 568–581. [https://doi.org/10.1016/S0008-6363\(00\)00268-6](https://doi.org/10.1016/S0008-6363(00)00268-6)
- Zoratti, M., Szabò, I., 1995. The mitochondrial permeability transition. *Biochim. Biophys. Acta* 1241, 139–176.
- Zou, H., Henzel, W.J., Liu, X., Lutschg, A., Wang, X., 1997. Apaf-1, a human protein homologous to *C. elegans* CED-4, participates in cytochrome c-dependent activation of caspase-3. *Cell* 90, 405–413.