

## DAFTAR PUSTAKA

- Adriaanse, M.C., Drewes, H.W., Heide, I.V.D., Struijs, J.N., & Baan, C.A., 2016, The impact of comorbid chronic conditions on quality of life in type 2 diabetes patients, *Qual Life Res* **25**, 175-181.
- Akinci, F., Yildirim, A., Gozu, H., Sargin, H., Orbay, E., & Sargin, M., 2008, Assessment of health-related quality of life (HRQoL) of patients with type 2 diabetes in Turkey, *Diabetes Research and Clinical Practice*, **79** (1), 117-123.
- Aklima, Kritpracha, C., & Thaniwattananon, P., 2013, Dietary Behaviours among Patients with Type 2 Diabetes Mellitus in Indonesia, *Journal of Nursing*, **3** (1), 499-509.
- Akter, S., Goto, A., & Mizoue, T., 2017, Smoking and the risk of type 2 diabetes in Japan: A systematic review and meta-analysis, *Journal of Epidemiology* **27**, 553-560.
- Al-Aboudi, I.S., Hassali, M.A., Shafie, A.A., Hassan, A., & Alrasheedy, A.A., 2015, A cross-sectional assessment of health-related quality of life among type 2 diabetes patients in Riyadh, Saudi Arabia, *SAGE Open Medicine* **3**, 1-5.
- Al-Haj Mohd, M.M.M., Phung, H., Sun, J., & Morisky, D.E., 2016, Improving adherence to medication in adults with diabetes in the United Arab Emirates, *BMC Public Health* **16** (857), 1-9.
- Al-Shehri, A.H., Taha, A.Z., Bahnassy, A.A., & Salah, M., 2008, Health-related quality of life in type 2 diabetic patients, *Ann Saudi Med* **28** (5), 352-357.
- Al Hayek, A.A., Robert, A.A., Al Saeed, A., Alzaid, A.A., & Al Sabaan, F.S., 2014, Factors Associated with Health-Related Quality of Life among Saudi Patients with Type 2 Diabetes Mellitus: A Cross-Sectional Survey, *Diabetes Metab J* **38**, 220-227.
- Amaliyah, R., 2016, Hubungan Tingkat Pengetahuan dan Tingkat Pendidikan dengan Kepatuhan Diet pada Pralansia Penderita Diabetes Mellitus, *Skripsi*, Fakultas Kesehatan Masyarakat, Universitas Airlangga, Surabaya.
- American Diabetes Association, 2002, Screening for diabetes, *Diabetes Care* **25**, (1), 21-24.

- American Diabetes Association*, 2018, Standards Of Medical Care In Diabetes-2018, *Diabetes Care* **41** (1), 517-522.
- American Diabetes Association*, 2018, Woman, diakses dari <http://www.diabetes.org/living-with-diabetes/treatment-and-care/women/>, 15 Mei 2018.
- Asif, M., 2014, The prevention and control the type-2 diabetes by changing lifestyle and dietary pattern, *Journal of Education and Health Promotion* **3** (1), 1-5.
- Butt, M., Ali, A.M., & Bakry, M.M., 2018, Health-Related Quality of Life in Poorly Controlled Type 2 Diabetes Patients-Association of Patients' Characteristics With EQ-5D Domains, Mean EQ-5D Score, and Visual Analog Scale Score, *Asian Journal of Pharmaceutical and Clinical Research* **11** (1), 1-6.
- Cardoso, A.F., Cruz, R., Queiros, P., Santiago, L., Ribeiro, C.F., & Ferreira, P.L., 2016, Assessment of Health-Related Quality of Life using the EQ-5D-3L in Individuals with Type 2 Diabetes Mellitus, *J Diabetes Metab Disord Control* **3** (2), 1-7.
- Chaliks, R., 2012, Kepatuhan Dan Kepuasan Terapi Dengan Antidiabetik Oral Pada Pasien Diabetes Melitus Tipe 2 Rawat Jalan Di RSUP Dr. Sardjito Yogyakarta, *Tesis*, Fakultas Farmasi, Universitas Gadjah Mada, Yogyakarta.
- Chentli, F., Azzoug, S., & Mahgoun, S., 2015, Diabetes mellitus in elderly, *Indoan Journal of Endocrinology and Metabolism* **19** (6), 744-752.
- Chyun, D.A., Melkus, G.D., Katten, D.M., Price, W.J., Davey, J.A., Grey, N., et al., 2006, The Association of Psychological Factors, Physical Activity, Neuropathy, and Quality of Life in Type 2 Diabetes, *Biological Research for Nursing* **7** (4), 279-288.
- Clarke P., Gray, A., Holman, R., 2002, Estimating utility values for health states of type 2 diabetic patients using the EQ-5D (UKPDS 62), *Med Decis Making* **22**, 340-349.
- Coffey, J.T., Brandle, M., Zhou, H., et al., 2002, Valuing health-related quality of life in diabetes, *Diabetes Care* **25**, 2238-2243.
- Colberg, S.R., Sigal, R.J., Fernhall, B., Regensteiner, J.G., Blissmer, B.J., Rubin, R.R., et al., 2010, Exercise and Type 2 Diabetes, *Diabetes Care* **33**, e147-e167.
- Cole, G., Leonard, B., Hammond, S., & Fridinger, F., 1998, Using "Stages of Behavioral Change" Constructs to Measure The Short-Term Effects

of a Worsite-Based Intervention to Increase Moderate Physical Activity', *Psychological Reports* **82**, 615-618.

Departemen Kesehatan RI, 2003, *Undang-undang Republik Indonesia nomor 20 tahun 2003 tentang Sistem Pendidikan Nasional*, Jakarta: Departemen Kesehatan RI.

Departemen Kesehatan RI, 2014, *Peraturan Menteri Kesehatan RI nomor 79 tahun 2014 tentang Penyelenggaraan Pelayanan Geriatri di Rumah Sakit*, Jakarta: Departemen Kesehatan RI.

Devlin, N.J., Krabbe, PF, 2013, The development of new research methods for the valuation of EQ-5D-5L, *The European Journal of Health Economics* **14**, 1.

Dinaryanti, P., Fudholi, A., & Andayani, T.M., 2012, Analisis Biaya dan Efektivitas Terapi Pasien Diabetes Melitus Tipe 2 Rawat Jalan di RSUD Sleman Yogyakarta, *Jurnal Manajemen dan Pelayanan Farmasi* **2** (1), 14-19.

DiPietro, L., 2001, Physical activity in aging: changes in patterns and their relationship to health and function, *The Journals of Gerontology* **56** (2), 13–22.

DiPiro, J. T. et al., 2009, *Pharmacotherapy Handbook*, 7<sup>th</sup> edition, 210, Mc Graw Hill, New York.

Ekwunife, O.I., Ezenduka, C.C., & Uzoma, B.E., 2016, Evaluating the sensitivity of EQ -5D in a sample of patients with type 2 diabetes mellitus in two tertiary health care facilities in Nigeria, *BMC Research Notes*, **9** (24), 1-4.

Fox, C.S., Sullivan, L., D'Agostino, R.B., & Wilson, P.W.F., 2004, The Significant Effect of Diabetes Duration on Coronary Heart Disease Mortality, *Diabetes Care* **27** (3), 704-708.

Frati, A.C., Iniestra, F., & Ariza, C.R., 1996, Acute Effect of Cigarette Smoking on Glucose Tolerance and Other Cardiovascular Risk Factors, *Diabetes Care* **19** (2), 1-7.

Funnell, M.M., 2008, Quality of life and insulin therapy in type 2 diabetes mellitus, *Insulin* **3**, 31–36.

Grandy, S., & Fox, K.M., 2008, EQ-5D visual analog scale and utility index values in individuals with diabetes and at risk for diabetes: Findings from the Study to Help Improve Early evaluation and management of risk factors Leading to Diabetes (SHIELD), *Health and Quality of Life Outcomes* **6** (18), 1-7.

- Grandy, S., Fox, K.M., *et al.*, 2012, Change in health status (EQ-5D) over 5 years among individuals with and without type 2 diabetes mellitus in the SHIELD longitudinal study, *Health and Quality of Life Outcomes* **10** (99), 1-6.
- Hayes, A., Arima, H., Woodward, M., Chalmers, J., Poulter, N., FMedSci, *et al.*, 2016, Changes in Quality of Life Associated with Complications of Diabetes: Results from the ADVANCE Study, *Value In Health* **19**, 36-41.
- Hendriks, S.H., Hartog, L.C., Groenier, K.H., Maas, A.H.E.M., Hateren, K.J.J., Kleefstra, N., *et al.*, 2016, Patient Activation in Type 2 Diabetes: Does It Differ between Men and Women?, *Journal of Diabetes Research* **2016**, 1-7.
- Hlatky, M.A., Chung, S.C., Escobedo, J., *et al.*, 2010, BARI 2D Study Group, The effect of obesity on quality of life in patients with diabetes and coronary artery disease. *AmHeart J* **159**, 292–300.
- Honish, A., Westerfield, W., Ashby, A., Momin, S., & Phillipi, R., 2006, Health-Related Quality of Life and Treatment Compliance with Diabetes Care, *Disease Management* **9** (4), 195-200.
- Hunger, M., Schunk, M., Meisinger, C., Peters, A., & Holle, R., 2012, Estimation of the relationship between body mass index and EQ-5D health utilities in individuals with type 2 diabetes: Evidence from the population-based KORA studies, *Journal of Diabetes and Its Complications* **26**, 413-417.
- Jacobson, A.M., Groot, M.D., & Samson, J.A., 1994, The Evaluation of Two Measures of Quality of Life in Patients With Type I and Type II Diabetes, *Diabetes Care* **17**(4), 267-273.
- Jannoo, Z., Wah, Y.B., Lazim, A.M., Hassali, M.A., 2017, Examining diabetes distress, medication adherence, diabetes self-care activities, diabetes-specific quality of life and health-related quality of life among type 2 diabetes mellitus patients, *Journal of Clinical & Translational Endocrinology* **9**, 48-54.
- Javanbakht, M., Abolhasani, F., Mashayekhi, A., Baradaran, H.R., & Jahangirinoudeh, Y., 2012, Health Related Quality of Life in Patients with Type 2 Diabetes Mellitus in Iran: A National Survey, diakses dari [www.plosone.org](http://www.plosone.org) pada 14 Maret 2018.
- Kementerian Kesehatan RI, 2011, *Diet Diabetes Melitus*, Jakarta:Kementerian Kesehatan.
- Kementerian Kesehatan RI, 2013, Riset Kesehatan Dasar, Jakarta: Kementerian Kesehatan.

- Kementerian Kesehatan RI, 2014, *Situasi dan Analisis Diabetes*, Pusat Data dan Informasi Kesehatan, Jakarta.
- Khotkar, K., Chaudari, S., Jadhav, P.R., Deshmukh, Y., 2017, Assesment of Medication Adherence in Type II Diabetic Patients: A Cross-sectional Study, *MGM Journal of Medical Science* **4** (2), 65-69.
- Kiadaliri, A.A., Najafi, B., & Mirmalek-Sani, M., 2013, Quality of life in people with diabetes:a systematic review of studies in Iran, *Journal of Diabetes & Metabolic Disorders* **12**(54), 1-8.
- Kiadaliri, A.A., Gerdtham, U-G., Eliasson, B., Gudbjornsdottir, S., Svensson, A-M., & Carlsson, K.S., 2014, Health Utilities of Type 2 Diabetes-Related Complications: A Cross-Sectional Study in Sweden, *Int. J. Environ. Res. Public Health* **11**, 4939-4949.
- Konsensus PERKENEI, 2015, *Penatalaksanaan DM Sesuai Konsensus PERKENEI 2015*.
- Kriska, A.M., 2000, Physical activity and the prevention of type 2 diabetes mellitus: how much for how long?, *Sports Medicine* **29** (3), 147-151.
- Lee, A.J., Morgan, C.L., Morrissey, M., Wittrup-Jensen, K.U., Kennedy-Martin, T., & Currie, C.J., 2005, Evaluation of the association between the EQ-5Dindex (health-related utility) and body mass index (obesity) in hospital-treated people with Type 1 diabetes, Type 2 diabetes and with no diagnosed diabetes, *Diabetic Medicine* **22**, 1482-1486.
- Lemeshow S., Hosmer D., Klar J., Lwanga S., 1990, *Adequacy of Sample Size in Health Studies*, John Wiley & Son.
- Li, Y., Xu, W., Liao, Z., Yao, B., Chen, X., Huang, Z., *et al.*, 2004, Induction of Long-term Glycemic Control in Newly Diagnosed Type 2 Diabetic Patients Is Associated With Improvement of  $\beta$ -Cell Function, *Diabetes Care*, **27** (11) :2601.
- Luk, A.O.Y., Zhang, Y., Ko, G.T.C., Brown, N., Ozaki, R., Tong, P.C.Y., *et al.*, 2014, Health-Related Quality of Life in Chinese Patients with Type 2 Diabetes: An Analysis of the Joint Asia Diabetes Evaluation (JADE) Program, *J Diabetes Metab* **5** (2), 1-7.
- Maggio, C.A., & Pi-Sunyer, F.X., 2003, Obesity and type 2 diabetes, *Endocrinol Metab Clin N Am* **32**, 805-822.
- Martinez, Y.V., Prado-Aguilar, C.A., Rascon-Pacheco, R.A., Valdivia-Martinez, J.J., 2008, Quality of life associated with treatment adherence in patients with type 2 diabetes: a cross-sectional study, *BMC Health Services Research* **8**, 1-8.

- Mooy, J.M., De Vries, H., Grootenhuis, P.A., Bouter, L.M., Heine, R.J., 2000, Major stressful life events in relation to prevalence of undetected type 2 diabetes: the Hoorn Study, *Diabetes Care* **23** (2), 197–201.
- Mulyani, R., Andayani, T.M., Pramantara, I.D.P., 2012, Kepatuhan Terapi Berbasis Insulin Pada Pasien Diabetes Melitus Tipe 2 di Poliklinik Endokrinologi RSUP Dr. Sardjito Yogyakarta, *Jurnal Manajemen dan Pelayanan Farmasi* **2** (2), 7 – 8.
- Nazir, S.U.R., Hassali, M.A., Saleem, F., Bashir, S., & Aljadhey, H., 2017, Does adherence to the therapeutic regimen associate with health related quality of life: Findings from an observational study of type 2 diabetes mellitus patients in Pakistan, *Pak. J. Pharm. Sci.* **30** (6), 2159-2164.
- Osterberg, L., & Blaschke, T., 2005, Adherence to Medication, *The new england journal of medicine* **353** (5), 487-494.
- Pemerintah DIY, 2017, Statistik Penduduk DIY, <http://www.kependudukan.jogjaprovo.go.id/olah.php?module=statistik&periode=9&jenisdata=penduduk&berdasarkan=pendidikan&prop=34&kab=04&kec=13>, 7 Juni 2018
- Pemerintah Kabupaten Sleman, 2017, Dashboard Data Kesehatan RSUD Sleman, <https://e-health.slemankab.go.id/main.php>, 22 September 2017.
- Purba F.D., Hunfeld J.A.M., Iskandarsyah A., Fitriana T.S., Sadarjoen S.S., Ramos-Goni J.M., et al., 2017, The Indonesian EQ-5D-5L Value Set, *PharmacoEconomics* DOI 10.1007/s40273-017-0538-9.
- Quah, J.H.M., Luo, N., Ng, W.Y., How, C.H., Tay, E.G., 2011, Health-related Quality of Life is Associated with Diabetic Complications, but not with Short-term Diabetic Control in Primary Care, *Ann Acad Med Singapore* **40**, 276-285.
- Redekop, W.K., Koopmanschap, M.A., Stolk, R.P., Rutten, G.E., Wolffenbuttel, B.H., Niessen, L.W., 2002, Health-related quality of life and treatment satisfaction in Dutch patients with type 2 diabetes, *Diabetes Care* **25**, 458–463.
- Rhee, M.K., Slocum, W., Ziemer, D.C., Culler, S.D., 2005, Patients Adherence Improves Glycemic Control, *Diabetes Educator* **31**(2): 240-50.
- Rhee, E.J., 2015, Diabetes in Asians, *Endocrinol Metab* **30** (3): 263-269.
- Ritter, et al., 2008, *A Textbook of Clinical Pharmacology and Therapeutics*, Fifth edition, 285 – 286, Hodden Education, London.
- Rubin, R.R., & Peyrot, M., 1999, Quality of Life and Diabetes, *Diabetes Metab Res Rev* **15**, 205-216.

- Rwegerera, G.M., Moshomo, T., Gaenamong, M., Oyewo, T.A., Gollakota, S., Rivera, Y.P., *et al.*, 2017, Health-related quality of life and associated factors among patients with diabetes mellitus in Botswana, *Alexandria Journal of Medicine* **1**, 1-6.
- Sakamaki, H., Ikeda, S., Ikegami, N., Uchigata, Y., Iwamoto, Y., Origasa, H., *et al.*, 2006, Measurement of HRQL Using EQ-5D in Patients with Type 2 Diabetes Mellitus in Japan, *International Society for Pharmacoeconomics and Outcomes Research* **9** (1), 47-52.
- Sastroasmoro, S., & Ismael, S., 2010, *Dasar-dasar Metodologi Penelitian Klinis, Edisi 3*, Jakarta: Sagung Seto.
- Solli, O., Stavem, K., Kristiansen, I.S., 2010, Health-related quality of life in diabetes: The associations of complications with EQ-5D scores, *Health and Quality of Life Outcomes* **8**, 18.
- Stipanovic, A.R., 2002, The Effects of Diabetes Education on Self-Efficacy and Self-Care of Adults with type 2 Diabetes, *Thesis*, Faculty of Nursing, University of Manitoba, Manitoba.
- Stratton, I.M., Adler, A.I., Neil, H.A.W., Matthews, D., Manley, S.E., Cull, C.A., *et al.*, 2000, Association of glycaemia with macrovascular and microvascular complications of type 2 diabetes (UKPDS 35): prospective observational study, *BMJ* **321** (1):407-410.
- Sullivan, P.W., and Ghuschyan, V.H., 2016, EQ-5D Scores for Diabetes-Related Comorbidities, *Value in Health* **19**, 1002 – 1008.
- Sultana, M., Sarker, A.R., Mahumud, R.A., Ahmed, S., Ahmed, W., Chakrovort, S., *et al.*, 2016, Inequalities in Health Status from EQ-5D Findings: A Cross-Sectional Study in Low-Income Communities of Bangladesh, *Int J Health Policy Manag* **5** (5), 301-307.
- Thiel, D.M., Al Sayah, F., Vallance, J.K., Johnson, S.T., Johnson, J.A., 2017, Association between Physical Activity and Health-Related Quality of Life in Adults with Type 2 Diabetes, *Can J Diabetes* **41**, 58-62.
- Timar, R., Velea, I., Timar, B., Lungeanu, D., Oancea, C. Roman, D., & Mazilu, O., 2016, Factors influencing the quality of life perception in patients with type 2 diabetes mellitus, *Patient Preference and Adherence* **10**, 2471-2476.
- Tucker, D.M.D., & Palmer, A.J., 2011, The cost-effectiveness of interventions in diabetes: A review of published economic evaluations in the UK setting, with an eye on the future, *Primary Care Diabetes* **5**, 9-16.

- Turner, R., 1998, Intensive blood-glucose control with sulphonylureas or insulin compared with conventional treatment and risk of complications in patients with type 2 diabetes (UKPDS 33), *The Lancet* **352**, 837-851.
- Utami, D.T., Karim, D., Agrina, 2014, Faktor-faktor yang Mempengaruhi Kualitas Hidup Pasien Diabetes Mellitus Dengan Ulkus Diabetikum, *Jom Psik* **1** (2), 1 – 6.
- Wasem J., *et al.*, 2013, Co-morbidity but not dysglycaemia reduces quality of life in patients with type-2 diabetes treated with oral mono- or dual combination therapy – an analysis of the DiaRegis registry, *Cardio Vascular Diabetology* **12**, 47.
- Wexler, D.J., Grant, R.W., Wittenberg, E., Bosch, J.L., Cagliero, E., Delahanty, L., *et al.*, 2006, Correlates of health-related quality of life in type 2 diabetes, *Diabetologia* **49**, 1489-1496.
- World Health Organization, 1982, *Privisional Guidelines On Standard International Age Classifications*, Departement of International Economic and Social Affairs, New York.
- World Health Organization, 2004, *BMI Classification*, diakses dari [http://www.who.int/bmi/index.jsp?introPage=intro\\_3.html](http://www.who.int/bmi/index.jsp?introPage=intro_3.html), 2 Februari 2018.
- World Health Organisation, 2011, *Global status report on non-communicable disease 2010*. Geneva: World Health Organisation.
- World Health Organization, 2014, Indonesia : Noncommunicable Disease, [http://www.who.int/nmh/countries/idn\\_en.pdf?ua=1](http://www.who.int/nmh/countries/idn_en.pdf?ua=1), 26 Oktober 2017.
- Wu, S-FV., Courtney, M., Edwards, H., McDowell, J., Shortridge-Bagget, L.M., Chang, P-J., 2007, Self-efficacy, outcome expectations and self-care behaviour in people with type 2 diabetes in Taiwan, *Journal of Clinical Nursing* **16**, 250-256.
- Xie, X., Liu, Q., Wu, J., & Wakui, M., 2009, Impact of cigarette smoking in type 2 diabetes development, *Acta Pharmacol Sin* **30** (6): 784-787.
- Yusra, A., 2011, Hubungan Antara Dukungan Keluarga Dengan Kualitas Hidup Pasien Diabetes Melitus Tipe 2 Di Poliklinik Penyakit Dalam Rumah Sakit Umum Pusat Fatmawati Jakarta, *Tesis*, Fakultas Ilmu Keperawatan, Universitas Indonesia, Depok.
- Zioga, E., Kazakos, K., Dimopoulos, E., Koutras, C., Marmara, K., Marmara, E., *et al.*, 2016, Adherence and Quality of Life in Patients With Type II Diabetes Mellitus in Northern Greece, *Mater Sociomed* **28** (4), 258-261.