

REFERENCES

- Aller., (1987). A Standardized System to Evaluate Ground Water Pollution Potential Using Hydrogeologic Settings. National Water Well Association. Retrieved from <http://rdn.bc.ca/cms/wpattachments/wpID3175atID5999.pdf>
- Firat Ersoy., (2013). DRASTIC-based methodology for assessing groundwater vulnerability in the Gmhaciky and Merzifon basin (Amasya, Turkey). *Earth Sciences Research Journal*, 17(1), 33–40. <https://doi.org/10.2118/1430-G>
- Foster, S., Hirata, R., Gomes, D., Elia, M., and Paris, M., (2002). *Groundwater Quality Protection: A Guide for Water Utilities, Municipal Authorities, and Environment Agencies*, World Bank, Washington, D.C.
- Gogu, R.C., and Dassargues, A., (2000). Current and Future trends and Future Challenges in Groundwater Vulnerability Assessment Using Overlay and Index Methods, *Environmental Geology*, 39: P 549 – 559pp.
- Langmuir D., (1972). Control on the amounts of pollutants in subsurface waters. *Earth and Mineral*.
- Lobbo-Ferraira, J.P and Olivereira.M.M., (2004). *Groundwater Vulnerability Assessment in Portugal*. *Journal of Geofisica International*, Vol. 43, Num4 p.541-550
- MacDonald. M., and Partners., (1984). Greater Yogyakarta Groundwater Resources Study Volume 3, Groundwater, Directorate General of Water Resources Development Groundwater Development project (P2AT), Ministry of Public Works, Government of the Republic of Indonesia.
- Panagopoulos, G. P., Antonakos, A. K., & Lambrakis, N. J., (2006). Optimization of the DRASTIC method for groundwater vulnerability assessment via the use of simple statistical methods and GIS. *Hydrogeology Journal*, 14(6), 894–911

- Putra, D. P. E., (2007). The Impact of Urbanization on Groundwater Quality: A Case Study in Yogyakarta City – Indonesia. Aachen, German: RWTH.
- Rahardjo, W., Sukandarrumidi., and Rosidi, H. M. D., (1995). Geologic Map of the Yogyakarta Quadrangle, Java, Geological Survey of Indonesia.
- Rahman, A., (2008). A GIS based DRASTIC model for assessing groundwater vulnerability in shallow aquifer in Aligarh, India. *Applied Geography*, 28(1), 32–53. <https://doi.org/10.1016/j.apgeog.2007.07.008>
- Santosa, L. W., (2010). The study of the genetic of reflux and its effect on hydrostratigraphic aquifers and hydrogeochemistry as geoindicators of the evolution of groundwater, southern part of Kulon Progo regency, Indonesia. Department of Geography, Gadjah Mada University, Indonesia.
- Sinha, M. K., Verma, M. K., Ahmad, I., Baier, K., Jha, R., & Azzam, R., (2016). Assessment of groundwater vulnerability using modified DRASTIC model in Kharun Basin, Chhattisgarh, India. *Arabian Journal of Geosciences*, 9(2), 1–22. <https://doi.org/10.1007/s12517-015-2180-1>
- Vrba, J., and Zaporozec, A. [Edts.] (1994). Guidebook on Mapping Groundwater Vulnerability: International Contributions to Hydrogeology. Vol16/1994.
- Van Bemmelen, V., (1949). The Geology of Indonesia Vol. IA General Geology of Indonesia.
- Zaporozec, A., (2002). Groundwater contamination inventory. IHP-VI, Series on Groundwater. International Hydrological Program, UNESCO, Paris. n. 2. Zaporozec, A. and Miller, JC (2000). Ground-Water Pollution. UNESCO, Paris. 24. 5. APPENDICES.