

DAFTAR PUSTAKA

- Ahmed, A. A., Ortashi, O. M., Abdelmagied, A. H., & Ramzy, M. S. (2004). Risk factors for recurrent miscarriage in Sudanese women. *Saudi Med J*, *25*(9), 1295–1297.
- American College of Obstetricians and Gynecologist. (2015). Early pregnancy loss. *Practice Bulletin*, *150*, 10.
- American College of Obstetricians and Gynecologists. (2016). FAQ: Repeated miscarriages. Retrieved December 25, 2017, from <https://www.acog.org/Patients/FAQs/Repeated-Miscarriages>
- American Society for Reproductive Medicine. (2005). Long-term use of oral contraceptives increases the risk of miscarriage. *Fertility and Sterility*, *83*(5), 2003–2005. <http://doi.org/10.1016/j.fertnstert.2004.11.085>
- American Society for Reproductive Medicine. (2012). Evaluation and treatment of recurrent pregnancy loss: a committee opinion. *Fertility and Sterility*, (11), 1–9. <http://doi.org/10.1016/j.fertnstert.2012.06.048>
- American Society for Reproductive Medicine. (2016). Fact Sheet: What is recurrent pregnancy loss (RPL)? Retrieved January 3, 2018, from <http://www.reproductivefacts.org/news-and-publications/patient-fact-sheets-and-booklets/fact-sheets-and-info-booklets/what-is-recurrent-pregnancy-loss-rpl/>
- Andersen, A. N., Wohlfahrt, J., Christens, P., Olsen, J., & Melbye, M. (2000). Maternal age and fetal loss: population based register linkage study. *BMJ*, *320*(June), 1708–1712.
- Aoki, K., Furukawa, T., Ogasawara, M., Hori, S., & Kitamura, T. (1998). Psychosocial factors in recurrent miscarriages. *Acta Obstetrica et Gynecologica Scandinavica*, *77*, 572–573.
- Bashiri, A., Ratzon, R., Amar, S., Serjienko, R., Mazor, M., & Shoham-wardi, I. (2012). Two vs. three or more primary recurrent pregnancy losses: are there any differences in epidemiologic characteristics and index pregnancy outcome? *Journal of Perinatal Medicine*, *40*, 365–371. <http://doi.org/10.1515/jpm-2011-0295>
- BPS, BKKBN, Kemenkes, & ICF International. (2013). *Indonesia Demographic and Health Survey 2012*. Jakarta, Indonesia. Retrieved from <http://dhsprogram.com/pubs/pdf/FR275/FR275.pdf>
- BPS, & Macro International. (2008). *Indonesia Demographic and Health Survey 2007*. Calverton, Maryland, USA. Retrieved from <http://dhsprogram.com/pubs/pdf/FR218/FR218.pdf>
- BPS, & ORC Macro. (2003). *Indonesia Demographic and Health Survey 2002-*

2003. Calverton, Maryland, USA. Retrieved from
<http://dhsprogram.com/pubs/pdf/FR147/FR147.pdf>

Bracken, M. B., Bryce-buchanan, C., Stlten, R., & Holford, T. (1985). Menarcheal age and habitual miscarriage: evidence for an association. *Annals of Human Biology*, 12(6), 525–531.

Bralic, I., Tahirovic, H., Matanic, D., Vrdoljak, O., Stojanovic-Spehar, S., Kovacic, V., & Blazekovic-Milakovic, S. (2012). Association of early menarche age and overweight/ obesity. *Journal of Pediatric Endocrinology*, 25(1–2), 57–62. <http://doi.org/10.1515/jpem-2011-0277>

CBS/ Indonesia, NFPCB/ Indonesia, Ministry of Health/ Indonesia, & Macro International. (1991). *Indonesia Demographic and Health Survey 1991*. Calverton, Maryland, USA. Retrieved from
<http://dhsprogram.com/pubs/pdf/FR18/FR18.pdf>

CBS/ Indonesia, NFPCB/ Indonesia, Ministry of Health/ Indonesia, & Macro International. (1995). *Indonesia Demographic and Health Survey 1994*. Calverton, Maryland, USA. Retrieved from
<http://dhsprogram.com/pubs/pdf/FR18/FR18.pdf>

CBS/ Indonesia, NFPCB/ Indonesia, Ministry of Health/ Indonesia, & Macro International. (1998). *Indonesia Demographic and Health Survey 1997*. Calverton, Maryland, USA. Retrieved from
<http://dhsprogram.com/pubs/pdf/FR95/FR95.pdf>

Christiansen, O. B., Pedersen, B., Nielsen, H. S., & Andersen, A. N. (2004). Impact of the sex of first child on the prognosis in secondary recurrent miscarriage. *Human Reproduction*, 19(12), 2946–2951.
<http://doi.org/10.1093/humrep/deh516>

Christiansen, O. B., Steffensen, R., & Nielsen, H. S. (2011). Anti-HY responses in pregnancy disorders. *American Journal of Reproductive Immunology*, 66, 93–100. <http://doi.org/10.1111/j.1600-0897.2011.01038.x>

Creswell, J. W. (2017). *Research design: pendekatan metode kualitatif, kuantitatif, dan campuran* (4th ed.). Yogyakarta, Indonesia: Pustaka Pelajar.

George, L., Granath, F., Johansson, A. L. V, Olander, B., Cnattingius, S., & George, L. (2006). Risks of repeated miscarriage. *Paediatric and Perinatal Epidemiology*, 20, 119–126.

Heijer, M. D. E. N., Thomas, C. M. G., & Eskes, T. O. M. K. A. B. (2000). Homocysteine and folate levels as risk factors for recurrent early pregnancy loss. *Obstetrics and Gynecology*, 95(4), 519–524.

Heikinheimo, O., Gissler, M., & Suhonen, S. (2008). Age, parity, history of abortion and contraceptive choices affect the risk of repeat abortion. *Contraception*, 78, 149–154.
<http://doi.org/10.1016/j.contraception.2008.03.013>

- Hemminki, E., & Forssas, E. (1999). Epidemiology of miscarriage and its relation to other reproductive events in Finland. *American Journal of Obstetric and Gynecology*, *181*(2), 396–401.
- Jellesen, R., Strandberg-larsen, K., Jørgensen, T., Olsen, J., Thulstrup, A. M., & Andersen, A. N. (2008). Maternal use of oral contraceptives and risk of fetal death. *Paediatric and Perinatal Epidemiology*, *22*, 334–340.
<http://doi.org/10.1111/j.1365-3016.2008.00942.x>
- Kesmodel, U., Wisborg, K., Olsen, S. F., Henriksen, T. B., & Secher, N. J. (2002). Moderate alcohol intake in pregnancy and the risk of spontaneous abortion. *Alcohol and Alcoholism*, *37*(1), 87–92.
- Kidd, S. A., Eskenazi, B., & Wyrobek, A. J. (2001). Effects of male age on semen quality and fertility: a review of the literature. *Fertility and Sterility*, *75*(2), 237–248.
- Kismiliansari, D. E., Nizomy, I. R., & Budiarti, L. Y. (2015). Hubungan antara usia ibu hamil dengan kejadian abortus habitualis di RSUD Ulin Banjarmasin periode tahun 2010 – 2013. *Berkala Kedokteran*, *11*(1), 73–83.
- Kleinbaum, D. G., Sullivan, K. M., & Barker, N. D. (2003). *ActivEpi companion textbook: A supplement for use with the ActivEpi CD-ROM*. New York, USA: Springer.
- Kupfermanc, M. J. (2003). Thrombophilia and pregnancy. *Reproductive Biology and Endocrinology*, *22*, 1–22.
- Lehmann, A., & Scheffler, C. (2016). What does the mean menarcheal age mean? — An analysis of temporal pattern in variability in a historical swiss population from the 19th and 20th centuries. *American Journal of Human Biology*, (August 2015), 1–9. <http://doi.org/10.1002/ajhb.22854>
- Lemeshow, S., Jr, D. W. H., Klar, J., & Lwanga, S. K. (1990). *Adequacy of sample size in health studies*. West Sussex, England: John Wiley & Sons Ltd.
- Li, J., Liu, L., Liu, B., Saravelos, S., & Li, T. (2014). Recurrent miscarriage and birth sex ratio. *European Journal of Obstetrics and Gynecology and Reproductive Biology*, *176*, 55–59.
<http://doi.org/10.1016/j.ejogrb.2014.02.031>
- Liestol, K. (1980). Menarcheal age and spontaneous abortion: a causal connection? *American Journal of Epidemiology*, *111*(6).
- Lindbohm, M., Sallmén, M., & Taskinen, H. (2002). Effects of exposure to environmental tobacco smoke on reproductive health by. *Scandinavian Journal of Work, Environment & Health*, *28*(2), 84–96.
- Madrigal, L. (1991). Menarcheal age and spontaneous abortion: Further evidence for a connection. *American Journal of Human Biology*, *628*.
- Metwally, M., Saravelos, S. H., Ledger, W. L., Phil, D., & Li, T. C. (2010). Body

- mass index and risk of miscarriage in women with recurrent miscarriage. *Fertility and Sterility*, 94(1), 290–295.
<http://doi.org/10.1016/j.fertnstert.2009.03.021>
- Mortensen, L. H., Nielsen, S., Cnattingius, S., & Andersen, A. N. (2011). Sex of the first-born and risk of preterm birth in the subsequent pregnancy. *Epidemiology*, 22(3). <http://doi.org/10.1097/EDE.0b013e31820e8600>
- Nielsen, H. S. (2011). Secondary recurrent miscarriage and H-Y immunity. *Human Reproduction Update*, 17(4), 558–574.
<http://doi.org/10.1093/humupd/dmr005>
- Nielsen, H. S., Andersen, A. N., Kolte, A. M., & Christiansen, O. B. (2008). A firstborn boy is suggestive of a strong prognostic factor in secondary recurrent miscarriage: a confirmatory study. *Fertility and Sterility*, 89(4), 907–911. <http://doi.org/10.1016/j.fertnstert.2007.04.029>
- Ooi, P. V., Russell, N., & O'Donoghue, K. (2011). Secondary recurrent miscarriage is associated with previous male birth. *Journal of Reproductive Immunology*, 88(1), 38–41. <http://doi.org/10.1016/j.jri.2010.10.004>
- Pokale, Y. S., & Khadke, P. (2016). Risk of recurrent miscarriage in India and the effect of paternal age and maternal age. *International Journal of Applied Sciences*, 03(03), 459–468.
<http://doi.org/http://dx.doi.org/10.21013/jas.v3.n3.p16>
- Poorolajal, J., Cheraghi, P., Cheraghi, Z., Ghahramani, M., & Irani, A. D. (2014). Predictors of miscarriage: a matched case-control study. *Epidemiology and Health*, 36(November), 1–7.
<http://doi.org/http://dx.doi.org/10.4178/epih/e2014031>
- Porter, T. F., & Scott, J. R. (2005). Evidence-based care of recurrent miscarriage. *Best Practice & Research Clinical Obstetrics and Gynaecology*, 19(1), 85–101. <http://doi.org/10.1016/j.bpobgyn.2004.11.005>
- PP HIFERI-POGI. (2011). *Panduan tata laksana keguguran berulang*. (A. Baziad, K. Sumapraja, & B. Santoso, Eds.). Jakarta.
- Puscheck, E. E., & Jeyendran, R. S. (2007). The impact of male factor on recurrent pregnancy loss. *Current Opinion in Obstetrics and Gynecology*, 222–228.
- Rai, R., & Regan, L. (2006). Recurrent miscarriage. *The Lancet*, 368(August), 601–611.
- Rao, V. R. C., Lakshmi, A., & Sadhnani, M. D. (2008). Prevalence of hypothyroidism in recurrent pregnancy loss in first trimester. *Indian Journal of Medical Sciences*, 62(9), 360–362.
- Regan, L., & Rai, R. (2000). Epidemiology and the medical causes of miscarriage. *Bailliere's Clinical Obstetric and Gynaecology*, 14(5), 839–854.

<http://doi.org/10.1053/beog.2000.0123>

- Say, L., Chou, D., Gemmill, A., Tunçalp, Ö., Moller, A., Daniels, J., ... Alkema, L. (2014). Global causes of maternal death: a WHO systematic analysis. *The Lancet*, (14), 1–11. [http://doi.org/10.1016/S2214-109X\(14\)70227-X](http://doi.org/10.1016/S2214-109X(14)70227-X)
- Scholl, T. O., & Johnson, W. G. (2000). Folic acid : influence on the outcome of pregnancy 1 – 4. *The American Journal of Clinical Nutrition*, 71.
- Shahine, L., & Lathi, R. (2015). Recurrent pregnancy loss: evaluation and treatment. *Obstetrics and Gynecology Clinics*, 42(1), 117–134. <http://doi.org/10.1016/j.ogc.2014.10.002>
- Shapira, E., Ratzon, R., Shoham-varadi, I., Serjienko, R., Mazor, M., & Bashiri, A. (2012). Primary vs. secondary recurrent pregnancy loss – epidemiological characteristics, etiology, and next pregnancy outcome. *Journal of Perinatal Medicine*, 40, 389–396. <http://doi.org/10.1515/jpm-2011-0315>
- Smith, M. L., & Schust, D. J. (2011). Endocrinology and recurrent early pregnancy loss. *Seminars in Reproductive Medicine*, 29.
- Stirrat, G. M. (1990). Recurrent miscarriage I: definition and epidemiology. *The Lancet*, (336), 673–675.
- Sugiura-ogasawara, M., Furukawa, T. A., Nakano, Y., Hori, S., Aoki, K., & Kitamura, T. (2002). Depression as a potential causal factor in subsequent miscarriage in recurrent spontaneous aborters. *Human Reproduction*, 17(10), 2580–2584.
- SurveyMETER. (2018). *Materi pelatihan STATA untuk pengolahan data IFLS: Kelas STATA dasar – lanjutan*. Yogyakarta, Indonesia: SurveyMETER.
- Thomas, F., Renaud, F., Benefice, E., & Meeüs, T. D. E. (2001). International variability of ages at menarche and menopause: patterns and main determinants. *Human Biology*, 73(2), 271–290.
- Wijaya, G. P. (2017). *Pengalaman traumatis pada wanita yang mengalami keguguran berulang*. Universitas Katolik Widya Mandala Surabaya.
- Wildschut, H. I. J. (1994). Sociodemographic factors: age, parity, social class, and ethnicity. In D. K. James, P. J. Steer, C. P. Weiner, & B. Gonik (Eds.), *High Risk Pregnancy: Management Options*. England: W. B. Saunders Company Ltd.
- Wisborg, K., Kesmodel, U., Bech, B. H., Hedegaard, M., & Henriksen, T. B. (2003). Maternal consumption of coffee during pregnancy and prospective study. *BMJ*, 326(February), 1–4.
- World Health Organization. (1977). WHO: Recommended definitions, terminology and format for statistical tables related to the perinatal period and use of a new certificate for cause for perinatal deaths. *Acta Obstetrica et Gynecologica Scandinavica*, 56, 247–253.

World Health Organization. (2016). Congenital anomalies. Retrieved January 2, 2018, from <http://www.who.int/mediacentre/factsheets/fs370/en/>

Zhang, B., Wei, Y., Niu, J., Li, Y., Miao, Z., & Wang, Z. (2010). Risk factors for unexplained recurrent spontaneous abortion in a population from southern China. *International Journal of Gynecology and Obstetrics*, *108*(2), 135–138. <http://doi.org/10.1016/j.ijgo.2009.09.019>