



DAFTAR PUSTAKA

- Aminiahidashti, H., Shafiee, S., Hosseininejad, S.M., Firouzian, A., Barzegarnejad, A., Kiasari, A.Z., Kerigh, B.F., Bozorgi, F., Shafizad, M. & Geraeeli, A. 2018. Comparison of the combination of propofol–fentanyl with combination of propofol–ketamine for procedural sedation and analgesia in patients with trauma. *The American Journal of Emergency Medicine*. Tersediadit <http://linkinghub.elsevier.com/retrieve/pii/S0735675718300937> [Accessed 17 April 2018].
- Avidan, M.S., Zhang, L., Burnside, B.A., Finkel, K.J., Searleman, A.C., Selvidge, J.A., Saager, L., Turner, M.S., Rao, S., Bottros, M., Hantler, C., Jacobsohn, E. & Evers, A.S. 2008. Anesthesia Awareness and the Bispectral Index. *New England Journal of Medicine*, 358(11): 1097–1108.
- Banister, R., Mishra, R. & Kaye, A.D. 2012. Sedation for interventional pain management procedures. R.D. Urman & A.D. Kaye, ed., *Moderate and Deep Sedation in Clinical Practice*. Cambridge: Cambridge University Press, hal.291–299. Tersedia di https://www.cambridge.org/core/product/identifier/9781139084000%23c40045-23-1/type/book_part [Accessed 17 April 2018].
- Barash, P.G. ed., 2013. *Clinical anesthesia ; [Includes interactive web and mobile versions with hundreds of videos and regular updates!]*. 7. ed ed. Anesthesiology. Philadelphia, PA: Wolters Kluwer, Lippincott Williams & Wilkins.
- Bestas, A., Erhan, O. & Ezici, M. 2013. As an analgesic ketamine versus fentanyl for total intravenous anesthesia. *Saudi Journal of Anaesthesia*, 7(1): 107.
- Butterworth, J.F., Mackey, D.C. & Wasnick, J.D. 2013. *Morgan & Mikhail's Clinical Anesthesiology*. Fifth ed. McGraw Hill Education.
- Chernik, D.A., Gillings, D., Laine, H., Hendler, J., Silver, J.M., Davidson, A.B., Schwam, E.M. & Siegel, J.L. 1990. Validity and reliability of the Observer's Assessment of Alertness/Sedation Scale: study with intravenous midazolam. *Journal of Clinical Psychopharmacology*, 10(4): 244–251.
- Chidambaran, V., Costandi, A. & D'Mello, A. 2015. Propofol: A Review of its Role in Pediatric Anesthesia and Sedation. *CNS Drugs*, 29(7): 543–563.



Chun, E.H., Han, M.J., Baik, H.J., Park, H.S., Chung, R.K., Han, J.I., Lee, H.J. & Kim, J.H. 2015. Dexmedetomidine-ketamine versus Dexmedetomidine-midazolam-fentanyl for monitored anesthesia care during chemoport insertion: a Prospective Randomized Study. *BMC Anesthesiology*, 16(1). Tersedia di <http://bmcanesthesiol.biomedcentral.com/articles/10.1186/s12871-016-0211-4> [Accessed 17 April 2018].

Chung, H.S. 2014. Awareness and recall during general anesthesia. *Korean Journal of Anesthesiology*, 66(5): 339.

Cole, D.J. & Schlunt, M. 2004. *Adult perioperative anesthesia*. 1st ed ed. The requisites in anesthesiology series. Philadelphia: Elsevier/Mosby.

Dal, T., Sazak, H., Tunç, M., Sahin, S. & Yılmaz, A. 2014. A comparison of ketamine-midazolam and ketamine-propofol combinations used for sedation in the endobronchial ultrasound-guided transbronchial needle aspiration: a prospective, single-blind, randomized study. *Journal of Thoracic Disease*, 6(6): 742–751.

Das, S. & Ghosh, S. 2015. Monitored anesthesia care: An overview. *Journal of Anaesthesiology Clinical Pharmacology*, 31(1): 27.

Driest, SLV, Marshall, M.D., Hachey, B., Beck, C., Crum, K., Owen, J., Smith, A.H., Kannankeril, P.J., Woodworth, A., Caprioli, R.M., Choi, L., 2016, Pragmatic Pharmacology: Population Pharmacokinetic Analysis of Fentanyl Using Reman Samples from Children after Cardiac Surgery, *Br J of Clinical Pharmacology* (81): 1165-1174

Eshghi, A., Mohammadpour, M., Kaviani, N., Tahirian, D. & Akhlaghi, N. 2016. Comparative evaluation of bispectral index system after sedation with midazolam and propofol combined with remifentanil versus ketamine in uncooperative during dental procedures. *Dental Research Journal*, 13(1): 1.

Gelen, S.A., Sarper, N., Demirsoy, U., Zengin, E. & Çakmak, E. 2015. The Efficacy and Safety of Procedural Sedoanalgesia with Midazolam and Ketamine in Pediatric Hematology. *Turkish Journal of Hematology*, 32(4): 351–354.

Godambe, S.A., Elliot, V., Matheny, D. & Pershad, J. 2003. Comparison of propofol/fentanyl versus ketamine/midazolam for brief orthopedic



procedural sedation in a pediatric emergency department. *Pediatrics*, 112(1 Pt 1): 116–123.

Hollman, G.A., Schultz, M.M., Eickhoff, J.C. & Christenson, D.K. 2008. Propofol-fentanyl versus propofol alone for lumbar puncture sedation in children with acute hematologic malignancies: Propofol dosing and adverse events: *Pediatric Critical Care Medicine*, 9(6): 616–622.

Johnson, P.N., Miller, J.L. & Hagemann, T.M. 2012. Sedation and Analgesia in Critically Ill Children: *AACN Advanced Critical Care*, 23(4): 415–434.

Kaplan, R.F., Cravero, J.P., Yaster, M. & Coté, C.J. 2009. Sedation for Diagnostic and Therapeutic Procedures Outside the Operating Room. *A Practice of Anesthesia for Infants and Children*. Elsevier, hal.1023–1048. Tersedia di http://www.crossref.org/deleted_DOI.html [Accessed 17 April 2018].

Kaul, H.L. & Bharti, N. 2002. Monitoring Depth Of Anaesthesia. *Indian Journal of Anaesthesia*, 46(4): 323–332.

Khajavi, M., Emami, A., Etezadi, F., Safari, S., Sharifi, A. & Shariat Moharari, R. 2013. Conscious Sedation and Analgesia in Colonoscopy: Ketamine/Propofol Combination has Superior Patient Satisfaction Versus Fentanyl/Propofol. *Anesthesiology and Pain Medicine*, 3(1): 208–13.

Laurenza, C. 2014. Evaluation of Sedation for the Unconscious Patient: Are We Still Far Away? *Journal of Anesthesia & Critical Care: Open Access*, 1(2). Tersedia di <http://medcraveonline.com/JACCOA/JACCOA-01-00007.php> [Accessed 18 April 2018].

Lysakowski, C., Dumont, L., Pellegrini, M., Clergue, F. & Tassonyi, E. 2001. Effects of fentanyl, alfentanil, remifentanil and sufentanil on loss of consciousness and bispectral index during propofol induction of anaesthesia. *British Journal of Anaesthesia*, 86(4): 523–527.

Malviya, S., Galinkin, J.L., Bannister, C.F., Burke, C., Zuk, J., Popenhagen, M., Brown, S. & Voepel-Lewis, T. 2009. The Incidence of Intraoperative Awareness in Children: Childhood Awareness and Recall Evaluation: *Anesthesia & Analgesia*, 109(5): 1421–1427.

Mandal, M., Basu, S. & Bagchi, D. 2014. Arousal time from sedation during spinal anaesthesia for elective infraumbilical surgeries: Comparison between propofol and midazolam. *Indian Journal of Anaesthesia*, 58(4): 403.



- Mandal, M., Basu, S., Khutia, S. & Das, S. 2012. Intravenous infusion of ketamine-propofol can be an alternative to intravenous infusion of fentanyl-propofol for deep sedation and analgesia in paediatric patients undergoing emergency short surgical procedures. *Indian Journal of Anaesthesia*, 56(2): 145.
- Mantadakis, E., Katzilakis, N., Foundoulaki, E. & Kalmanti, M. 2009. Moderate Intravenous Sedation With Fentanyl and Midazolam for Invasive Procedures in Children With Acute Lymphoblastic Leukemia. *Journal of Pediatric Oncology Nursing*, 26(4): 217–222.
- Meyer, S., Aliani, S., Graf, N. & Gottschling, S. 2004. Inter- and Intraindividual Variability in Ketamine Dosage in Repetitive Invasive Procedures in Children with Malignancies. *Pediatric Hematology and Oncology*, 21(2): 161–166.
- Migita, R.T., Klein, E.J. & Garrison, M.M. 2006. Sedation and Analgesia for Pediatric Fracture Reduction in the Emergency Department: A Systematic Review. *Archives of Pediatrics & Adolescent Medicine*, 160(1): 46.
- Murray, A. 2011. Awareness During Anesthesia. *Anesthesia Secrets*. Elsevier, hal.207–209. Tersedia di <http://linkinghub.elsevier.com/retrieve/pii/B9780323065245000301> [Accessed 18 April 2018].
- Musizza, B. & Ribaric, S. 2010. Monitoring the Depth of Anaesthesia. *Sensors*, 10(12): 10896–10935.
- Orser, B.A., Mazer, C.D. & Baker, A.J. 2008. Awareness during anesthesia. *Canadian Medical Association Journal*, 178(2): 185–188.
- Overly, F.L., Wright, R.O., Connor, F.A., Jay, G.D. & Linakis, J.G. 2005. Bispectral analysis during deep sedation of pediatric oral surgery patients. *Journal of Oral and Maxillofacial Surgery*, 63(2): 215–219.
- Pacifici, G.M., 2014, Clinical Pharmacology of Fentanyl in Preterm Infants. A Review, Elsevier, hal 143-148. Tersedia di <http://www.pediatr.neonatol.com> [Accessed 19 April 2018]
- Phelan, L., Stargatt, R. & Davidson, A.J. 2009. Long-term posttraumatic effects of intraoperative awareness in children. *Pediatric Anesthesia*, 19(12): 1152–1156.



- Sadhasivam, S., Ganesh, A., Robison, A., Kaye, R. & Watcha, M.F. 2006. Validation of the Bispectral Index Monitor for Measuring the Depth of Sedation in Children: *Anesthesia & Analgesia*, 102(2): 383–388.
- Samuelsson, P., Brudin, L. & Sandin, R.H. 2007. Late Psychological Symptoms after Awareness among Consecutively Included Surgical Patients. *Anesthesiology*, 106: 26–32.
- Sastroasmoro, S. & Ismael, S. 2014. *Dasar-dasar Metodologi Penelitian Klinis*. Binarupa Aksara.
- Sen, J. & Sen, B. 2014. A comparative study on monitored anesthesia care. *Anesthesia: Essays and Researches*, 8(3): 313.
- Sinha, P.K. & Koshy, T. 2007. Monitoring Devices for Measuring the Depth of Anaesthesia – An Overview. *Indian Journal of Anaesthesia*, 51(5): 365–381.
- Stoelting, R.K. & Hillier, S.C. 2006. Nonbarbiturate intravenous anesthetic drugs. *Pharmacology and Physiology in Anesthetic Practice*, 4th ed. Lippincott Williams & Wilkins, hal.155–178.
- Szostakiewicz, K.M., Tomaszewski, D., Rybicki, Z. & Rychlik, A. 2014. Śródoperacyjny powrót świadomości podczas znieczulenia ogólnego — wyniki obserwacyjnego badania ankietowego. *Anestezjologia Intensywna Terapia*, 46(1): 23–28.
- Urman, R.D. & Kaye, A.D. ed., 2012. *Moderate and deep sedation in clinical practice*. Cambridge ; New York: Cambridge University Press.
- Vuyk, J. 1997. Pharmacokinetic and Pharmacodynamic Interactions Between Opioids and Propofol. *Journal of Clinical Anesthesia*, 9(6): 23S–26S.
- Whitwam, J.G. ed., 1998. *Principles and practice of sedation*. 2. ed ed. Oxford: Blackwell Science.