

DAFTAR PUSTAKA

- [1] H. A. Quigley, "Number of people with glaucoma worldwide," *Br. J. Ophthalmol.*, vol. 80, no. 5, pp. 389–393, 1996.
- [2] A. Jackson and S. Radhakrishnan, "Understanding and Living with Glaucoma," *Glaucoma Res. Found.*, pp. 1–33, 2016.
- [3] H. Quigley and A. T. Broman, "The number of people with glaucoma worldwide in 2010 and 2020," *Br. J. Ophthalmol.*, vol. 90, no. 3, pp. 262–267, 2006.
- [4] E. S. Affandi, M. Siholita, and W. Artini, "The reduction of intraocular pressure after instillation of travoprost compared with timolol in chronic primary angle-closure glaucoma," *Med J Indones*, vol. 15, no. 4, pp. 242–245, 2006.
- [5] R. Chrástek *et al.*, "Automated segmentation of the optic nerve head for diagnosis of glaucoma," *Med. Image Anal.*, vol. 9, no. 4, pp. 297–314, 2005.
- [6] R. R. a Bourne, "The optic nerve head in glaucoma," *Community Eye Heal.*, vol. 19, no. 59, pp. 44–45, 2006.
- [7] J. Nayak, R. Acharya U., P. S. Bhat, N. Shetty, and T. C. Lim, "Automated diagnosis of glaucoma using digital fundus images," *J. Med. Syst.*, vol. 33, no. 5, pp. 337–346, 2009.
- [8] R. Acharya U, C. K. Chua, E. Y. K. Ng, W. Yu, and C. Chee, "Application of higher order spectra for the identification of diabetes retinopathy stages," *J. Med. Syst.*, vol. 32, no. 6, pp. 481–488, 2008.
- [9] M. M. Hermann *et al.*, "Interobserver variability in confocal optic nerve analysis (HRT)," *Int. Ophthalmol.*, vol. 26, no. 4–5, pp. 143–149, 2005.
- [10] M. M. Hermann, I. Theofylaktopoulos, N. Bangard, C. Jonescu-Cuypers, S. Coburger, and M. Diestelhorst, "Optic nerve head morphometry in healthy

- adults using confocal laser scanning tomography,” *Br. J. Ophthalmol.*, vol. 88, no. 6, pp. 761–765, 2004.
- [11] G. J. Jaffe and J. Caprioli, “Optical coherence tomography to detect and manage retinal disease and glaucoma,” *Am. J. Ophthalmol.*, vol. 137, no. 1, pp. 156–169, 2004.
- [12] M. Fingeret, F. A. Medeiros, R. Susanna, and R. N. Weinreb, “Five rules to evaluate the optic disc and retinal nerve fiber layer for glaucoma,” *Optometry*, vol. 76, no. 11, pp. 661–668, 2005.
- [13] S. Y. Lee *et al.*, “Automated quantification of retinal nerve fiber layer atrophy in fundus photograph,” *Conf. Proc. IEEE Eng. Med. Biol. Soc.*, vol. 2, pp. 1241–1243, 2004.
- [14] R. Kolar and J. Jan, “Detection of Glaucomatous Eye via Color Fundus Images Using Fractal Dimensions,” *Radioengineering*, vol. 17, no. 3, pp. 109–114, 2008.
- [15] P. . Prageeth, D. J, and a S. Kumar, “Early detection of retinal nerve fiber layer defects using fundus image processing,” *2011 IEEE Recent Adv. Intell. Comput. Syst.*, pp. 930–936, 2011.
- [16] G. D. Joshi, J. Sivaswamy, R. Prashanth, and S. R. Krishnadas, “Detection of peri-papillary atrophy and RNFL defect from retinal images,” *Lect. Notes Comput. Sci. (including Subser. Lect. Notes Artif. Intell. Lect. Notes Bioinformatics)*, vol. 7325 LNCS, no. PART 2, pp. 400–407, 2012.
- [17] M. V Wyawahare and U. I. Nagar, “Feature selection and classification for automatic detection of retinal nerve fibre layer thinning in retinal fundus images,” *Int. J. Biomed. Eng. Technol.*, vol. 19, no. 3, pp. 205–219, 2015.
- [18] R. Panda, N. B. Puan, A. Rao, D. Padhy, and G. Panda, “Recurrent neural network based retinal nerve fiber layer defect detection in early glaucoma,” *Proc. - Int. Symp. Biomed. Imaging*, pp. 692–695, 2017.
- [19] A. Issac, M. Partha Sarathi, and M. K. Dutta, “An adaptive threshold based

- image processing technique for improved glaucoma detection and classification,” *Comput. Methods Programs Biomed.*, vol. 122, no. 2, pp. 229–244, 2015.
- [20] T. Khalil, S. Khalid, and A. M. Syed, “Review of Machine Learning techniques for glaucoma detection and prediction,” *Proc. 2014 Sci. Inf. Conf. SAI 2014*, pp. 438–442, 2014.
- [21] R. Gayathri, P. V. Rao, and S. Aruna, “Automated glaucoma detection system based on wavelet energy features and ANN,” in *2014 International Conference on Advances in Computing, Communications and Informatics, ICACCI*, 2014, pp. 2808–2812.
- [22] M. Pahlitzsch *et al.*, “Significance of the disc damage likelihood scale objectively measured by a non-mydratic fundus camera in preperimetric glaucoma,” *Clin. Ophthalmol.*, vol. 9, pp. 2147–2158, 2015.
- [23] G. M. A. Budai, J. Hornegger, “Multiscale Approach for Blood Vessel Segmentation on Retinal Fundus Images,” *Invest. Ophthalmol. Vis. Sci.*, vol. 50, no. 13, p. 325, 2009.
- [24] W. T. Andrew, “Primary Open-Angle Glaucoma: A Transforming Growth Factor- β Pathway-Mediated Disease,” *Am. J. Pathol.*, vol. 180, no. 6, pp. 2201–2204, 2012.
- [25] D. Krawitz, “WHAT IS GLAUCOMA?” [Online]. Available: <https://www.huntingtoneyecare.com/what-is-glaucoma/>. [Accessed: 20-Jul-2018].
- [26] R. R. Allingham, *Shields Textbook of Glaucoma, 6th Edition*, 6th ed. Philadelphia: Lippincott Williams & Wilkins, 2011.
- [27] A. Kadir and A. Susanto, *Pengolahan Citra: Teori dan Aplikasi*, 1st ed. Yogyakarta: Penerbit Andi, 2013.
- [28] C. Solomon and T. Breckon, *Fundamentals of Digital Image Processing: A Practical Approach with Examples in Matlab*, 1st ed. Wiley-Blackwell,

2011.

- [29] S. S. Al-amri, N. V Kalyankar, and S. D. Khamitkar, "Linear and Non-linear Contrast Enhancement Image," *J. Comput. Sci.*, vol. 10, no. 2, pp. 139–143, 2010.
- [30] P. Darma, *Pengolahan Citra Digital*, 1st ed. Yogyakarta: Penerbit Andi, 2010.
- [31] A. M. Reza, "Realization of the contrast limited adaptive histogram equalization (CLAHE) for real-time image enhancement," *J. VLSI Signal Process. Syst. Signal Image. Video Technol.*, vol. 38, no. 1, pp. 35–44, 2004.
- [32] R. C. Gonzalez and R. E. Woods, *Digital Image Processing (3rd Edition)*. 2007.
- [33] X. Bai and F. Zhou, "Multi Scale Top-hat Transform Based Algorithm for Image Enhancement," *ICSP2010 Proc.*, pp. 797–800, 2010.
- [34] V. Barrile, G. M. Meduri, and D. Cuzzocrea, "Automatic Recognition of Road Signs by Hough Transform: Road-GIS," *J. Earth Sci. ...*, vol. 2, pp. 42–50, 2012.
- [35] N. Jain and N. Jain, "Coin Recognition Using Circular Hough Transform," *Int. J. Electron. Commun. Comput. Technol.*, vol. 2, no. 3, pp. 101–104, 2012.
- [36] M. Smereka and I. Dulęba, "Circular object detection using a modified Hough transform," *Int. J. Appl. Math. Comput. Sci.*, vol. 18, no. 1, pp. 85–91, 2008.
- [37] M. Kass, a. Witkin, and D. Terzopoulos, "Snakes: Active contour models," *Int. J. Comput. Vis.*, vol. 1, no. 4, pp. 321–331, 1988.
- [38] V. Caselles, R. Kimmel, and G. Sapiro, "Geodesic active contours," *IEEE Int. Conf. Comput. Vis.*, vol. 22, no. 1, pp. 694–699, 1995.
- [39] I. Maglogiannis, "Detection of irregular edges in skin lesions using parametric active contours for dermatological digital images," *J. Inf.*

Technol. Healthc., vol. 5, no. 2, 2007.

- [40] E. Alimudin, “Klasifikasi Citra Mamogram Digital untuk Skrining Mamografi,” Universitas Gadjah Mada, 2015.
- [41] W. Han, J. Dong, Y. Guo, M. Zhang, and J. Wang, “Identification of masses in digital mammogram using an optimal set of features,” in *Proc. 10th IEEE Int. Conf. on Trust, Security and Privacy in Computing and Communications, TrustCom 2011, 8th IEEE Int. Conf. on Embedded Software and Systems, ICESS 2011, 6th Int. Conf. on FCST 2011*, 2011, pp. 1763–1768.
- [42] F. Esposito, D. Malerba, G. Semeraro, and J. Kay, “A comparative analysis of methods for pruning decision trees,” *IEEE Trans. Pattern Anal. Mach. Intell.*, vol. 19, no. 5, pp. 476–491, 1997.

