

Daftar Pustaka

- Adcock, IM, 2016. HDAC inhibitors as anti-inflammatory agents. *Br J Pharmacol* 150(7): 829–831.
- Agostino, L, Martinon, F, Burns, K, McDermott, MF, Hawkins, PN, Tschopp, J, 2004. NALP3 forms an IL-1 processing inflammasome with increased activity in Muckle–Wells autoinflammatory disorder. *Immunity* 20:319–325.
- Angelantonio, DE, Gao, P, Pennells, L, Kaptoge, S, Caslake, M, Thompson, A, et al, 2012. Lipid related markers and cardiovascular disease prediction. *JAMA* 307:2499–2506.
- Armutcu, F, Coskun, O, Gürel, A, Kanter, M, Can, M, Ucar, F, et al, 2005. Thymosin alpha 1 attenuates lipid peroxidation and improves fructose-induced steatohepatitis in rats. *Clin Biochem* 38:540–547.
- Bauernfeind, F, Horvath, G, Stutz, A, Alnemri, ES, MacDonald, K, Speert, D, et al, 2009. NF- κ B activating pattern recognition and cytokine receptors license NLRP3 inflammasome activation by regulating NLRP3 expression. *J Immunol* 183(2):787–791.
- Besten, GD, Eunen, KV, Groen, AK, Venema, K, Reijngoud, DJ, Bakker, BM, 2016. The role of short-chain fatty acids in the interplay between diet, gut microbiota, and host energy metabolism. *Gut Microbes* 7(3): 189–200.
- Boekholdt, SM, Arsenault, BJ, Mora, S, Pedersen, TR, LaRosa, JC, Nestel, PJ, et al, 2012. Association of LDL cholesterol, nonHDL cholesterol, and apolipoprotein B levels with risk of cardiovascular events among patients treated with statins: a meta-analysis. *JAMA* 307:1302–1309.
- Brough, D, and Rothwell, NJ, 2007. Caspase-1-dependent processing of pro-interleukin-1 is cytosolic and precedes cell death. *J Cell Sci* 120:772–781.
- Campbell, JM, Fahey, GC, Wolf, BW, 1997. Selected indigestible oligosaccharides affect large bowel mass, cecal and fecal short-chain fatty acids, pH and microflora in rats. *American Society Nutrition Sc* 10(2): 133–137.
- Canani, RB, Costanzo, MD, Leone, L, Pedata, M, Meli, R, Calignano, A, 2011. Potential beneficial effects of butyrate in intestinal and extraintestinal diseases. *World J Gastroenterol* 17(12):1519–1528.
- Carta, S, Lavieri, R, Rubartelli, A, 2013. Different members of the il-1 family come out in different ways: DAMPs vs. Cytokines. *Front Immunol* 4:123.
- Chang, PV, Hao, L, Offermanns, S, Medzhitov, R, 2014. The microbial metabolite butyrate regulates intestinal macrophage function via histone deacetylase inhibition. *Proc Natl Acad Sci* 111: 2247 –2252.
- Church, LD, Cook, GP, McDermott, MF, 2008. Inflammasomes and interleukin-1 in inflammatory disorders. *Nat Clin Pract Rheumatol* 4(1):34–42.

- Clark, AKD, Aquisto, F, Gentry, C, Marchand, F, McMahon, SB, Malcangio, M, 2006. Rapid co-release of interleukin 1 and caspase 1 in spinal cord inflammation. *J Neurochem* 99:868–880.
- Coelho, DF, Pereira-Lancha, LO, Chaves, DS, Diwan, D, Ferraz, R, Campos-Ferraz, PL, Poortmans JR, et al, 2011. High-fat diets on body composition, lipid metabolism and insulin sensitivity, and the role of exercise on these parameters. *Braz J Med Biol Res* 44(10):966-972
- Comalada, M, Bailon, E, de-Haro, O, Lara-Villoslada, F, Xaus, J, Zarzuelo, A, et al, 2006. The effects of short-chain fatty acids on colon epithelial proliferation and survival depend on the cellular phenotype. *J Cancer Res Clin Oncol* 132:487–497
- Dinarello, CA, 2004. Therapeutic strategies to reduce IL-1 activity in treating local and systemic inflammation. *Curr Opin Pharmacol* 4:378–385.
- Dinarello, CA, 2011. Interleukin-1 in the pathogenesis and treatment of inflammatory diseases. *Blood* 117(14):3720-3732.
- Dinarto, M, 2000. Nutrisi pada penyakit vaskular aterosklerotik. Di dalam: Pegangan Penatalaksanaan Nutrisi Pasien. Jakarta: *PDGMI* 2:86-91.
- Donath, Y, Dalmas, E, Sauter, NS, 2013. Inflammation in Obesity and Diabetes: Islet dysfunction and therapeutic opportunity. *Cell metabolism* 17(6):860-872
- Donohoe, DR, Collins, LB, Wali, A, Bigler, R, Sun, W, Bultman, SJ, 2012. The warburg effect dictates the mechanism of butyrate-mediated histone acetylation and cell proliferation. *Mol Cell* 48:612–626.
- Duangnin, N, Klangjorhor, J, Tipparat, P, Pinmanee, S, Phitak, T, Pothacharoen, P, et al, 2017. Anti-inflammatory effect of methanol extracts of hemp leaf in IL-1 -induced synovitis. *Trop J Pharm Res* 16(7):1553-1563.
- Evans, K, Burdge, GC, Wootton, SA, Clark, ML, Frayn, KN, 2002. Regulation of dietary fatty acid entrapment in subcutaneous adipose tissue and skeletal muscle. *Diabetes* 51:2684–2690.
- Ferreira, CM, Vieira, AT, Vinolo, MA, Oliveira, FA, Curi, R, Martins FS, 2014. The central role of the gut microbiota in chronic inflammatory diseases. *J Immunol Res* 10:689-692.
- Gattorno, M, Tassi, S, Carta, S, Delfino, L, Ferlito, F, Pelagatti, MAD, et al, 2007. Pattern of interleukin-1beta secretion in response to lipopolysaccharide and ATP before and after interleukin-1 blockade in patients with CIAS1 mutations. *Arthritis Rheum* 56(9):3138–3148.
- Greten, FR, Arkan, MC, Bollrath, J, Hsu, LC, Goode, J, Miething, C, et al, 2007. NF-kB Is a negative regulator of IL-1b secretion as revealed by genetic and pharmacological Inhibition of IKKb. *Cell* 130(5):918–931.
- Grundy, SM, 2012. Use of emerging lipoprotein risk factors in assessment of cardiovascular risk. *JAMA* 307:2540–2542.

- He, Y, Hara, H, Nunez, G, 2016. Mechanism and regulation of NLRP3 inflammasome activation. *Cell* 41(12):1012–1021.
- Hemphill, LC, 2010. Familial hypercholesterolemia: Current treatment options and patient selection for low-density lipoprotein apheresis. *J Clin Lipid* 4:346–349.
- Huang, S, Rutkowsky, JM, Snodgrass, RG, Moore, KDO, Schneider, DA, Newman, JA, 2012. Saturated fatty acids activate TLR-mediated proinflammatory signaling pathways. *J Lipid Res* 53(9):2002–2013.
- Hurst, NR, Kendig, DM, Murthy, KS, Grider, JR, 2014. The short chain fatty acids, butyrate and propionate, have differential effects on the motility of the guinea pig colon. *Neurogastroenterol Motil* 26(11):1586-1596.
- Inoue, A, Ikoma, K, Morioka, N, Kumagai, K, Hashimoto, T, Hide, I, et al, 1999. Interleukin-1beta induces substance P release from primary afferent neurons through the cyclooxygenase-2 system. *J Neurochem* 73(5):2206–2213.
- Iraporda, C, Errea, A, Romanin, DE, Cayet, D, Pereyra, E, Pignataro O, et al, 2015. Lactate and short chain fatty acids produced by microbial fermentation downregulate proinflammatory responses in intestinal epithelial cells and myeloid cells. *Immunobiol* 220(10):1161-9.
- Israel, A, 2010. The IKK complex, a central regulator of NF- B activation. *Cold Spring Harb Perspect Biol* 2(3):149-158.
- Kamada, N, Seo, SU, Chen, GY, Nunez, G, 2013. Role of the guts microbiota in immunity and inflammatory disease. *Nat Rev Immunol* 13: 321-335.
- Kawasaki, Y, Zhang, L, Cheng, JK, Ji, RR, 2008. Cytokine mechanisms of central sensitization: distinct and overlapping role of interleukin-1beta, interleukin-6, and tumor necrosis factor-alpha in regulating synaptic and neuronal activity in the superficial spinal cord. *J Neurosci* 28:5189–5194.
- Khitan, Z, Kim, DH, 2013. Fructose: a key factor in the development of metabolic syndrome and hypertension. *J Nutr Metab* 10(3):673-685.
- Kim, MH, Kang, SG, Park, JH, Yanagisawa, M, Kim, CH, 2013. Short-chain fatty acids activate GPR41 and GPR43 on intestinal epithelial cells to promote inflammatory responses in mice. *Gastroenterol* 145:396 –406.
- Klop, B, Elte, JWF, Cabezas, MC, 2013. Dyslipidemia in obesity: mechanisms and potential targets. *Nutrients* 5(4):1218-1240.
- LeMaitre, CL, dan Hoyland, JA, 2007. Catabolic cytokine expression in degenerate and herniated human intervertebral discs: IL-1beta and TNF alpha expression profile. *Arthritis Res Ther* 9(4):7-10.
- Livak, KJ, dan Schmittgen, TD, 2001. Analysis of relative gene expression data using real-time quantitative PCR and the 2(- CT) Method. *PubMed* 25(4):402-408.
- Lottenberg, AMP, Fan, PLT, Buonacorso, P, 2006. Effects of dietary fiber intake on inflammation in chronic diseases. *J Clin Gastroenterol* 40(3):235-43.

- Luo, ZL, Ren, JD, Huang, Z, Wang, T, Xiang, K, Cheng, L, et al, 2017. The role of exogenous hydrogen sulfide in free fatty acids induced inflammation in macrophages. *Cell Physiol Biochem* 42(4):1635-1644.
- Lynn, DA, dan Brown, GR, 2009. The ontogeny of exploratory behavior in male and female adolescent rats (*Rattus norvegicus*). *Dev Psychobiol* 51(6): 513–520.
- Maa, MC, Chang, MY, Hsieh, MY, Chen, YJ, Yang, CJ, Chen, ZC, et al, 2010. Butyrate reduced lipopolysaccharide-mediated macrophage migration by suppression of Src enhancement and focal adhesion kinase activity. *J Nutr Biochem* 21:1186–1192.
- Macia, L, Tan, J, Vieira, AT, Leach, K, Stanley, D, Luong, S, et al, 2015. Metabolite-sensing receptors GPR43 and GPR109A facilitate dietary fibre-induced gut homeostasis through regulation of the inflammasome. *Nat Commun* 6:6734.
- Maithilikarpagaselvi, M, Sridhar, MG, Swaminathan, RP, Sripradha, R, Badhe, B, 2016. Curcumin inhibits hyperlipidemia and hepatic fat accumulation in high-fructose-fed male Wistar rats. *Pharm Biol* 54(12):2857-2863.
- Maki, KC, Dicklin, MR, Schild, AL, Kling, D, Davidson, MH, 2014. Plasma Fatty Acids as Predictors of Triglyceride and Non-HDL Cholesterol Responses to Omega-3 Free Fatty Acid Therapy in Hypertriglyceridemia. *J Clin Lipid* 8(3):341–342.
- Mariathasan, S, dan Monack, D, 2007. Inflammasome adaptors and sensors: intracellular regulators of infection and inflammation. *Nat Immunol* 7:31–40.
- Martinon, F, dan Tschopp, J, 2007. Inflammasome caspases and inflammasomes: master switches of inflammation. *Cell Death Differ* 14:10–22.
- Masi, LN, Martins, AR, Crisma, AR, Amaral, CR, Davanzo, MR, Serdan, TDA, et al, 2017. Combination of a high-fat diet with sweetened condensed milk exacerbates inflammation and insulin resistance induced by each separately in mice. *Nature* 7:3937.
- Maslowski, KM, Vieira, AT, Ng, A, Kranich, J, Sierro, F, Yu, D, 2009. Regulation of inflammatory responses by gut microbiota and chemoattractant receptor GPR43. *Nature* 461:1282-1286.
- McGillicuddy, FC, Harford, KA, Reynolds, CM, Oliver, E, Claessens, M, Mills, KHG, et al, 2011. Lack of interleukin-1 receptor i (IL-1RI) protects mice from high-fat diet-induced adipose tissue inflammation coincident with improved glucose homeostasis. *Diabetes* 60(6): 1688-1698.
- McKenna, S, Butler, B, Jatana, K, Ghosh, S, Wright, CJ, 2017. Inhibition of I B /NF B signaling prevents LPS-induced IL1 expression without increasing apoptosis in the developing mouse lung. *Pediatric Research* 82:1064–1072.

- Mirmonsef, P, Zariffard, MR, Gilbert, D, Makinde, H, Landay, AL, Spear, GT, 2012. Short- chain fatty acids induce pro-inflammatory cytokine production alone and in combination with toll-like receptor ligands. *Am J Reprod Immunol* 67:391 –400.
- Morrison, DJ, dan Preston, T, 2016. Formation of short chain fatty acids by the gut microbiota and their impact on human metabolism. *J Gut Microbes* 7(3):189-200.
- Nelson, RH, 2013. Hyperlipidemia as a Risk Factor for Cardiovascular Disease. *J Prim Care* 40(1):195–211.
- Nies, LK, Cymbala, AA, Kasten, SL, Lamprecht, DG, Olson, KL, 2006. Complementary and alternative therapies for the management of dyslipidemia. *Ann Pharmacother* 40:1984–1992.
- Ning, B, Wang, X, Yu, Y, Waqar, AB, Yu, Q, Koike, T, et al, 2015. High-fructose and high-fat diet-induced insulin resistance enhances atherosclerosis in Watanabe heritable hyperlipidemic rabbits. *Nutr Metab* 12:30-36.
- Ochoa, EC, Ortega, KH, Ferrera, P, Morimoto, S, Arias, C, 2014. Short-term high-fat-and-fructose feeding produces insulin signaling alterations accompanied by neurite and synaptic reduction and astroglial activation in the rat hippocampus. *J Cereb Blood Flow Metab* 34(6):1001–1008.
- Ohira, H, Fujioka, Y, Katagiri, C, Yano, M, Mamoto, R, Aoyama, M, et al, 2012. Butyrate enhancement of inteleukin-1 production via activation of oxidative stress pathways in lipopolysaccharide-stimulated THP-1 cells. *J Clin Biochem Nutr* 50(1): 59–66.
- Oosterveer, DM, Versmissen, J, Yazdanpanah, M, Hamza, TH, Sijbrands, EJ, 2009. Differences in characteristics and risk of cardiovascular disease in familial hypercholesterolemia patients with and without tendon xanthomas: a systematic review and meta-analysis. *Atherosclerosis* 207:311–317.
- Oprea, A, Kress, M, 2000. Involvement of the proinflammatory cytokines Tumor Necrosis Factor- α , IL-1 β , and IL-6 but not IL-8 in the development of heat hyperalgesia: effects on heat-evoked Calcitonin Gene-Related Peptide release from rat skin. *J Neurosci* 20(16):6289–6293.
- Ostos, MA, Recalde, D, Baroukh, N, Callejo, A, Rouis, M, Castro, G, et al, 2002. Fructose Intake Increases Hyperlipidemia and Modifies Apolipoprotein Expression in Apolipoprotein AI-CIII-AIV Transgenic Mice. *J Nutr* 132: 918–923.
- Panchal, SK, Poudyal, H, Iyer, A, Nazer, R, Alam, MA, Diwan, V, et al, 2011. High- carbohydrate, high-fat diet-induced metabolic syndrome and cardiovascular remodeling in rats. *J Cardiovasc Pharmacol* (5):611-624.
- Park, J, Kim, M, Kang, SG, Jannasch, AH, Cooper, B, Patterson, J, et al, 2015. Short-chain fatty acids induce both effector and regulatory T cells by

- suppression of histone deacetylases and regulation of the mTOR-S6K pathway. *Mucosal Immunol* 8: 80–93.
- Park, JS, Lee, EJ, Lee, JC, Kim, WK, Kim, HS, 2007. Anti-inflammatory effects of short chain fatty acids in IFN-gamma-stimulated RAW 264.7 murine macrophage cells: involvement of NF-kappaB and ERK signaling pathways. *Int Immunopharmacol* 7(1):70-77.
- Park, MJ, dan Sohrabji, F, 2016. The histone deacetylase inhibitor, sodium butyrate, exhibits neuroprotective effects for ischemic stroke in middle-aged female rats. *J Neuroinflam* 13:300.
- Pektas, MB, Koca, HB, Sadi, G, Akar, F, 2016. Dietary Fructose Activates Insulin Signaling and Inflammation in Adipose Tissue: Modulatory Role of Resveratrol. *Bio Med Research Int* 10(2):247-252.
- Ren, K, dan Torres, R, 2009. Role of interleukin-1 during pain and inflammation. *Brain Res* 60(1): 57–64.
- Safieh-Garabedian, B, Poole, S, Allchorne, A, Winter, J, Woolf, CJ, 1995. Contribution of interleukin-1 to the inflammation-induced increase in nerve growth factor levels and inflammatory hyperalgesia. *J Pharmacol* 115:1265–1275.
- Sandra, MCF dan Budiman, I, 2011. Efek fruktosa dan glukosa terhadap kadar trigliserida plasma. 11(1): 39-47.
- Santos, SHS, Fernandes, LR, Pereira, CS, Guimarães, ALS, de Paula, AMB, Santos, MJC, et al, 2012. Increased circulating angiotensin-(1–7) protects white adipose tissue against development of a proinflammatory state stimulated by a high-fat diet. *Reg Peptides* 178(3):64-70.
- Schnetzler BM, Boller S, Debray S, Bouzakri K, Meier DT, Prazak R, et al, 2009. Free fatty acids induce a proinflammatory response in islets via the abundantly expressed interleukin-1 receptor I. *Endocrinology* 150(12):5218-5229.
- Segain, J, de la Bletiere, DR, Bourreille, A, Leray, V, Gervois, N, Rosales, C, et al, 2000. Butyrate inhibits inflammatory responses through NF B inhibition: implications for Crohn's disease. *Gut* 47(3): 397–403.
- Shamash S, Reichert F, Rotshenker ST, 2002. The cytokine network of Wallerian degeneration: Tumor Necrosis Factor- , Interleukin-1 , and Interleukin-1 . *J Neurosci* 22(8):3052–3060.
- Sharma, M, Ansari, MT, Abou-Setta, AM, Soares-Weiser, K, Ooi, TC, Sears, M, et al, 2009. Systematic review: comparative effectiveness and harms of combination therapy and monotherapy for dyslipidemia. *Ann Intern Med* 151:622–630.
- Sina, C, Gavrilova, O, Forster, M, Till, A, Derer, S, Hildebrand, F, 2009. G protein-coupled receptor 43 is essential for neutrophil recruitment during intestinal inflammation. *J Immunol* 183: 7514 –7522.

- Stevinson, C, Pittler, MH, Ernst, E, 2000. Garlic for treating hypercholesterolemia. A meta- analysis of randomized clinical trials. *Ann Intern Med* 133:420–429.
- Sumiyoshi, M, Sakanaka, M, Kimura, Y, 2006. Chronic intake of high-fat and high-sucrose diets differentially affects glucose intolerance in mice. *J Nutr* 136(3):582-587.
- Tang, Q, Feng, Z, Tong, M, Xu, J, Zheng, G, Shen, L, et al, 2017. Piceatannol inhibits the IL-1 -induced inflammatory response in human osteoarthritic chondrocytes and ameliorates osteoarthritis in mice by activating Nrf2. *Food Funct* 8(11):3926-3937.
- Tedelind, S, Westberg, F, Kjerrulf, M, Vidal, A, 2007. Anti-inflammatory properties of the shortchain fatty acids acetate and propionate: a study with relevance to inflammatory bowel disease. *World J Gastroenterol* 13: 2826 –2832.
- Tranchida F, Tchiakpe L, Rakotoniaina Z, Deyris V, Ravion O, Hiol A, 2012. Long-term high fructose and saturated fat diet affects plasma fatty acid profile in rats. *J Zhejiang Univ Sci B* 13(4):307-17.
- Tripathy, D, Mohanty, P, Dhindsa, S, Syed, T, Ghanim, H, Aljada, A, et al, 2003. Elevation of free fatty acids induces inflammation and impairs vascular reactivity in healthy subjects. *Diabetes* 52(12):2882-2887.
- Tsalissavrina, I, Wahono, D, Handayani, D, 2006. Pengaruh Pemberian Diet Tinggi Karbohidrat Dibandingkan Diet Tinggi Lemak Terhadap Kadar Trigliserida Dan Hdl Darah Pada Rattus Novergicus Galur Wistar. *JKB* 22(2):80-89
- Vieira, AT, Macia, L, Vinolo, MA, Tavares L, Sousa, LP, Mackay, CR, 2016. Dietary fiber and the short-chain fatty acid acetate promote resolution of neutrophilic inflammation in a model of gout in mice. *J Leuko Biol* 101(1): 133-139.
- Vinolo, MAR, Ferguson, GJ, Kulkarni, S, Damoulakis, G, Anderson, K, Bohlooly, YM, et al, 2007. SCFAs induce mouse neutrophil chemotaxis through the GPR43 receptor. *PLoS ONE* 6: 212-216.
- Vinolo, MAR, Rodrigues, HG, Hatanaka, E, Hebeda, CB, Farsky, SH, Curi, R, 2009. Short-chain fatty acids stimulate the migration of neutrophils to inflammatory sites. *Clin Sci* 117: 331 –338.
- Vinolo, MAR, Rodrigues, HG, Nachbar, RT, Curi, R, 2011. Regulation of inflammation by short chain Fatty acids. *Nutrients* 3: 858 –876.
- Wider, B, Pittler, MH, Thompson-Coon, J, Ernst, E, 2009. Artichoke leaf extract for treating hypercholesterolaemia. *Cochrane Database Syst Rev* 7(4):133-135.
- Winzell, MS, Ahrén, B, 2004. The high-fat diet-fed mouse: a model for studying mechanisms and treatment of impaired glucose tolerance and type 2 diabetes. *Diabetes* 53:215-219.

- Wong, JM, de Souza, R, Kendall, CW, Emam, A, Jenkins, DJ, 2006. Colonic health: fermentation and short chain fatty acids. *J Clin Gastroenterol* 40(3):235-43.
- Xu,MX, Wang,M, Yang, WW, 2017. Gold-quercetin nanoparticles prevent metabolic endotoxemia-induced kidney injury by regulating Tlr4/NF-kB signaling and Nrf2 pathway in high fat diet fed mice. *Int J Nanomed* 12:327–345
- Yeboah, J, McClelland, RL, Polonsky, TS, Burke, GL, Sibley, CT, O'Leary, D, et al, 2012. Comparison of novel risk markers for improvement in cardiovascular risk assessment in intermediate-risk individuals. *JAMA* 308:788–795.
- Yoshida, M, Yoshimura, N, Hangai, M, Tanihara, H, Honda, Y, 2004. Interleukin-1 alpha, interleukin-1 beta, and tumor necrosis factor gene expression in endotoxin-induced uveitis. *Invest Ophthalmol Vis Sci* 35(3):1107-1113.
- Zhang, JM, dan Jianxiong, AN, 2007. Cytokines, Inflammation and Pain. *Int Anesthesiol Clin* 45(2):27–37.
- Zheng, G, Lin, L, Zhong, S, Zhang, Q, Li, D, 2015. Effects of Puerarin on Lipid Accumulation and Metabolism in High-Fat Diet-Fed Mice. *PLoS ONE* 10(3): e0122925.
- Zhou, Q, Wu, J, Tang, J, Wang, J, Lu, CH, Wang, P, 2015. Beneficial effect of higher dietary fiber intake on plasma HDL-C and TC/HDL-C ratio among chinese rural-to-urban migrant workers. *Int J Environ Res Public Health* 12(5): 4726–4738.
- Zhou, Z, Hui, X, Li, H, Lam, KS, Xiao, Y, Wu, D, et al, 2010. Adipocyte fatty acid-binding protein modulates inflammatory responses in macrophages through a positive feedback loop involving c-Jun NH2-terminal kinases and activator protein-1. *J Biol Chem* 285(14):273-280.