



REFERENCES

- Deodhar, M., Wood, K., Black, D.S., and Kumar, N., 2012, Synthesis of oxygenated 4-arylisoﬂavans and 4-arylﬂavans, *Tetrahedron Lett.*, 53, 6697–6700.
- Gharpure, S.J., Sathiyarayanan, A.M., and Jonnalagadda, P., 2008, o-Quinone methide based approach to isoﬂavans: application to the total syntheses of equol, 3'-hydroxyequol and vestitol, *Tetrahedron Lett.*, 49, 2974–2978.
- Guan, L.P. and Liu, B.Y., 2016, Antidepressant-like effects and mechanisms of ﬂavonoids and related analogues, *Eur. J. Med. Chem.*, 121, 47–57.
- Iskander, G., Wood, K., Eiffe, E., Yee, E.M.H., Black, D.S., and Kumar, N., 2015, Ritter reactions of isoﬂavanols: A facile route to 4-amidoisoﬂavans, *Tetrahedron Lett.*, 56, 1941–1943.
- Kang, M.R., Park, K.H., Oh, S.J., Yun, J., Lee, C.W., Lee, M.Y., et al., 2015, Cardiovascular protective effect of glabridin: Implications in LDL oxidation and inflammation, *Int. Immunopharmacol.*, 29, 914–918.
- Karabin, M., Hudcova, T., Jelinek, L., and Dostalek, P., 2014, Biotransformations and biological activities of hop ﬂavonoids, *Biotechnol. Adv.*, 33, 1063–1090.
- Liu, J., Wang, X., Yong, H., Kan, J., and Jin, C., 2018, Recent advances in ﬂavonoid-grafted polysaccharides: Synthesis, structural characterization, bioactivities and potential applications, *Int. J. Biol. Macromol.*, 116, 1011–1025.
- Lu, Z.H., Gu, X.J., Shi, K.Z., Li, X., Chen, D.D., and Chen, L., 2017, Accessing anti-human lung tumor cell line (A549) potential of newer 3,5-disubstituted pyrazoline analogs, *Arab. J. Chem.*, 10, 624–630.
- Mazimba, O. and Keroletswe, N., 2015, Flavans: Synthetic Strategies: A Review, *Int. Res. J. Pure Appl. Chem.*, 8, 112–146.
- Moon, Y.J., Wang, X., and Morris, M.E., 2006, Dietary ﬂavonoids: Effects on xenobiotic and carcinogen metabolism, *Toxicol. Vitro.*, 20, 187–210.
- Mourouti, N. and Panagiotakos, D.B., 2013, Soy food consumption and breast cancer, *Maturitas*, 76, 118–122.
- Nwachukwu, I.D., Luciano, F.B., and Udenigwe, C.C., 2013, The inducible soybean glyceollin phytoalexins with multifunctional health-promoting properties, *Food Res. Int.*, 54, 1208–1216.
- Raffa, D., Maggio, B., Raimondi, M.V., Plescia, F., and Daidone, G., 2017, Recent discoveries of anticancer ﬂavonoids, *Eur. J. Med. Chem.*, 142, 213–228.



- Saito, K., Yonekura-Sakakibara, K., Nakabayashi, R., Higashi, Y., Yamazaki, M., Tohge, T., and Fernie, A.R., 2013, The flavonoid biosynthetic pathway in Arabidopsis: Structural and genetic diversity, *Plant Physiol. Biochem.*, 72, 21–34.
- Sebastian, R.S., Wilkinson Enns, C., Goldman, J.D., Steinfeldt, L.C., Martin, C.L., Clemens, J.C., Murayi T., and Moshfegh, A. J., 2017, New, publicly available flavonoid data products: Valuable resources for emerging science, *J. Food Compos. Anal.*, 64, 68–72.
- Sghaier, M.B., Skandrani, I., Nasr, N., Franca, M.G.D., Chekir-Ghedira, L., and Ghedira, K., 2011, Flavonoids and sesquiterpenes from *Teucrium ramosissimum* promote antiproliferation of human cancer cells and enhance antioxidant activity: A structure-activity relationship study, *Environ. Toxicol. Pharmacol.*, 32, 336–348.
- Shamsuzzaman, Khanam, H., Dar, A.M., Siddiqui, N., and Rehman, S., 2016, Synthesis, characterization, antimicrobial and anticancer studies of new steroidal pyrazolines, *J. Saudi Chem. Soc.*, 20, 7–12.
- Takashima, Y., Kaneko, Y., and Kobayashi, Y., 2010, Synthetic access to optically active isoflavans by using allylic substitution, *Tetrahedron*, 66, 197–207.
- Takekuma, S.I., Kaibara, M., Minematsu, T., and Takekuma, H., 2011, Reaction of azulene with 1,2-bis[4-(dimethylamino)phenyl]-1,2-ethanediol in a mixed solvent of methanol and acetonitrile in the presence of hydrochloric acid: A facile one-pot synthesis and properties of new triarylethylenes possessing an azulene-1-yl group, *Tetrahedron*, 67, 4780–4792.
- Tilley, A.J., Zanatta, S.D., Qin, C.X., Kim, I.K., Seok, Y.M., Stewart, A., Woodman O. L., and Williams S. J., 2012, 2-Morpholinoisoflav-3-enes as flexible intermediates in the synthesis of phenoxodiol, isophenoxodiol, equol and analogues: Vasorelaxant properties, estrogen receptor binding and Rho/RhoA kinase pathway inhibition, *Bioorganic Med. Chem.*, 20, 2353–2361.
- Wang, T. yang, Li, Q., and Bi, K. shun, 2017, Bioactive flavonoids in medicinal plants: Structure, activity and biological fate, *Asian J. Pharm. Sci.*, 13, 12–23.
- Won, D., Shin, B.K., Kang, S., Hur, H.G., Kim, M., and Han, J., 2008, Absolute configurations of isoflavan-4-ol stereoisomers, *Bioorganic Med. Chem. Lett.*, 18, 1952–1957.
- Xia, J.B., Zhu, C., and Chen, C., 2013, Visible light-promoted metal-free C-H activation: Diarylketone-catalyzed selective benzylic mono- and difluorination, *J. Am. Chem. Soc.*, 135, 17494–17500.



Xiao, Z., Waters, N.C., Woodard, C.L., Li, Z., and Li, P.K., 2001, Design and synthesis of Pfmrk inhibitors as potential antimalarial agents, *Bioorganic Med. Chem. Lett.*, 11, 2875–2878.