

DAFTAR PUSTAKA

- Arfian, N., Muflikhah, K., Soeyono, S. K., Sari, D. C. R., Tranggono, U., Anggorowati, N., 2016. Vitamin D Attenuates Kidney Fibrosis Via Reducing Fibroblast Expansion, Inflammation, and Epithelial Cell Apoptosis. *Kobe J Med Sci.* 62(2):E38-E44.
- Baek, J. H., Birchmeier, C., Zenke, M., Hieronymus, T., 2012. The HGF receptor/Met tyrosine kinase is a key regulator of dendritic cell migration in skin immunity. *J Immunol.* 189(4):1699-1707.
- Barnes, J. L. & Gorin, Y., 2011. Myofibroblast differentiation during fibrosis: role of NAD(P)H oxidases. *Kidney Int.* 79(9):944–956.
- Boor, P., Ostendorf, T., Floege, J., 2010. Renal fibrosis: novel insights into mechanisms and therapeutic targets. *Nat Rev Nephrol.* 6(11):643–656.
- Bottinger, E. P. & Bitzer, M., 2002. TGF- β signaling in renal development and disease. *J Am Soc Nephrol.* 13:2600-2610.
- Boutet, A., De Frutos, C. A., Maxwell, P. H., Mayol, M. J., Romero, J., Nieto, M. A., 2006. Snail activation disrupts tissue homeostasis and induces fibrosis in the adult kidney. *EMBO J.* 25(23):5603-5613.
- Celic, T., Spanjol, J., Grskovic, A., Markic, D., Prebilib, I., Fuckar, Z., *et al.*, 2011. Bone morphogenetic protein-7 reduces cold ischemic injury in rat kidney. *Transplant Proc.* 43(7):2505-2509.
- Chevalier, R. L., Forbes, M. S., Thornhill, B. A., 2009. Ureteral obstruction as a model of renal interstitial fibrosis and obstructive nephropathy. *Kidney Int.* 75(11):1145–1152.
- Chung, A. C. & Lan H. Y., 2011. Chemokines in renal injury. *J Am Soc Nephrol.* 22(5):802-809.
- Clifford, M. N., 1999. Chlorogenic acids and other cinnamates—nature, occurrence and dietary burden. *J Sci Food Agric.* 79(3):362–372.
- Dai, C. & Liu, Y., 2004. Hepatocyte growth factor antagonizes the profibrotic action of TGF- β 1 in mesangial cells by stabilizing Smad transcriptional corepression TGIF. 15(6):1402-1412.
- Domitrović, R., Cvijanovic, O., Susnic, V., Katalinic N., 2014. Renoprotective mechanisms of chlorogenic acid in cisplatin-induced kidney injury. *Toxicology.* 324:98–107.
- Duffield, J. S., 2010. Macrophages and immunologic inflammation of the kidney. *Semin Nephrol.* 30(3):234-254.
- Eddy, A., A., Lopez, G. J. M., Okamura, D. M., Yamaguchi, I., 2012. Investigating mechanisms of chronic kidney disease in mouse models. *Pediatr Nephrol.* 27(8):1233-1247.
- Efstratiadis, G., Divani, M., Katsioulis, E., Vergoulas, G., 2009. Renal fibrosis. *Hippokratia.* 13(4):224–229.
- Gong, R., Rifai, A., Dworkin, L. D., 2006. Anti-inflammatory effect of hepatocyte growth factor in chronic kidney disease: Targeting the inflamed Vascular Endothelium. *J Am Soc Nephrol.* 17:2464-2473.

- Grande, M. T. & López-Novoa, J. M., 2009. Fibroblast activation and myofibroblast generation in obstructive nephropathy. *Nat Rev Nephrol.* 5(6):319–328.
- Hewitson, T. D., 2009. Renal tubulointerstitial fibrosis: common but never simple. *Am J Physiol Renal Physiol.* 296(6):1239-1244.
- Hruska, K. A., Guo, G., Wozniak, M., Martin, D., Miller, S., Liapis, H., *et al.*, 2000. Osteogenic protein-1 prevents renal fibrogenesis associated with ureteral obstruction. *Am J Physiol Renal Physiol.* 279(1):F130-F143.
- Hwang, S. J., Kim, Y. W., Park, Y., Lee, H. J., Kim, K. W., 2014. Anti-inflammatory effects of chlorogenic acid in lipopolysaccharide-stimulated RAW 264.7 cells. *Inflamm Res.* 63(1):81–90.
- Jha, V., 2009. Current status of chronic kidney disease care in southeast Asia. *Semin Nephrol.* 29(5):487–496.
- Ji, L., Jiang, P., Lu, B., Sheng, Y., Wang, X., Wang, Z., *et al.*, 2013. Chlorogenic acid, a dietary polyphenol, protects acetaminophen-induced liver injury and its mechanism. *J Nutr Biochem.* 24(11):1911–1919.
- Kaissling, B. & LeHir, M., 2008. The renal cortical interstitium: morphological and functional aspects. *Histochem Cell Biol.* 130(2):247–262.
- Kalluri, R. & Zeisberg, M., 2003. Exploring the connection between chronic renal fibrosis and bone morphogenic protein-7. *Histol Histopathol.* 18(1):217-224.
- Kementrian Kesehatan RI, 2013. Situasi Penyakit Ginjal Kronis. *InfoDATIN.*
- Kim, J., Lee, S., Shim, J., Kim, H. W., Kim, J., Jang, Y. J., *et al.*, 2012. Caffeinated coffee, decaffeinated coffee, and the phenolic phytochemical chlorogenic acid up-regulate NQO1 expression and prevent H₂O₂-induced apoptosis in primary cortical neurons. *Neurochem Int.* 60(5):466–474.
- Ko, G. J., Boo, C. S., Jo, S. K., Cho, W. Y., Kim, H. K., 2008. Macrophages contribute to the development of renal fibrosis following ischemia/reperfusion-induced acute kidney injury. *Nephrol Dial Transplant.* 23:842-852.
- Kriz, W., Kaissling, B., LeHir, M., 2011. Epithelial-mesenchymal transition (EMT) in kidney fibrosis: fact or fantasy? *J Clin Invest.* 121(2):468–474.
- LeBleu, V. S., Taduri, G., O’Connell, J., Teng, Y., Cooke, V. G., Woda, C., *et al.*, 2013. Origin and function of myofibroblasts in kidney fibrosis. *Nat Med.* 19(8):1047–1053.
- Levey, A. S. & Coresh, J., 2012. Chronic kidney disease. *Lancet.* 379(9811): 165–180.
- Li, R. X., Yiu, W. H., Wu, H. J., Wong, D. W., Chan, L. Y., Lin, M., *et al.*, 2015. BMP7 reduces inflammation and oxidative stress in diabetic tubulopathy. *Clin Sci (Lond).* 128(4):269-280.
- Lin, J., Patel, S. R., Cheng, X., Cho, E. A., Levitan, I., Ullenbruch, M., 2005. Kielin/chordin-like protein, a novel enhancer of BMP signaling, attenuates renal fibrotic disease. *Nat Med.* 11(4):387-393.
- Lin, S. L., Castano, A. P., Nowlin, B. T., Lupher, M. L., Duffield, J. S., 2009. Bone marrow Ly6Chigh monocytes are selectively recruited to injured kidney and differentiate into functionally distinct populations. *J Immunol.* 183(10):6733-6743.

- Liu, Y., 2004. Hepatocyte growth factor in kidney fibrosis: therapeutic potential and mechanisms of action. *Am J Physiol Renal Physiol*. 287(1):F7-16.
- Liu, Y., 2011. Cellular and molecular mechanisms of renal fibrosis. *Nat Rev Nephrol*. 7(12):684-696.
- López-Novoa, J. M. & Nieto, M. A., 2009. Inflammation and EMT: an alliance towards organ fibrosis and cancer progression. *EMBO Mol Med*. 1(6-7):303-314.
- Manson, S. R., Austin, P. F., Guo, Q., Moore, K. H., 2015. BMP-7 Signaling and its Critical Roles in Kidney Development, the Responses to Renal Injury, and Chronic Kidney Disease. *Vitam Horm*. 99:91-144.
- Manson, S. R., Niederhoff, R. A., Hruska, K. A., Austin, P. F., 2011. The BMP-7 Smad1/5/8 pathway promotes kidney repair after obstruction induced renal injury. *J Urol*. 185(6):2523-2530.
- Massague, J. & Wotton, D., 2000. Transcriptional control by the TGF-beta/Smad signaling system. *EMBO J*. 19(8):1745-1754.
- Matsumoto, K. & Nakamura, T., 2001. Hepatocyte growth factor: renotropic role and potential therapeutics for renal disease. *Kidney Int*. 59(6):2023-2038.
- Matsumoto, K. & Nakamura, T., 2002. Renotropic role and therapeutic potential of HGF in the kidney. *Nephrol Dial Transplant*. 17 Suppl 9:59-61.
- Meng, X. M., Chung, A. C., Lan, H. Y., 2013. Role of the TGF-β/BMP-7/Smad pathways in renal diseases. *Clin Sci (Lond)*. 123(4):243-254.
- Meng, X. M., Nikolic-Paterson, D. J., Lan, H. Y., 2016. TGF-β: the master regulator of fibrosis. *Nat Rev Nephrol*. 12(6):325-338.
- Meran, S. & Steadman, R., 2011. Fibroblasts and myofibroblasts in renal fibrosis. *Int J Exp Pathol*. 92(3):158-167.
- Micallef, L., Vedrenne, N., Billet, F., Coulomb, B., Darby, I. A., Desmouliere, A., 2012. The myofibroblast, multiple origins for major roles in normal and pathological tissue repair. *FTR*. 5(1):S5.
- Misaki, T., Yamamoto, T., Suzuki, S., Fukasawa, H., Togawa, A., Ohashi, N., *et al.*, 2009. Decrease in tumor necrosis factor-alpha receptor-associated death domain results from ubiquitin-dependent degradation in obstructive renal injury in rats. *Am J Pathol*. 175(1):74-83.
- Mizuno, S., Matsumoto, K., Kurosawa, T., Horikawa, Y. M., Nakamura, T., 2000. Reciprocal balance of hepatocyte growth factor and transforming growth factor-beta 1 in renal fibrosis in mice. *Kidney Int*. 57(3):937-948.
- Monteiro, M., Farah, A., Perrone, D., Trugo, L. C., Donangelo, C., 2007. Chlorogenic acid compound from coffee are differentially absorbed and metabolized in human. *J Nutr*. 137(10):2196-2201.
- Multazam, A., 2017. *Efek protektif asam klorogenat terhadap ginjal pada mencit dengan cedera iskemia reperfusi* [tesis]. Yogyakarta, Universitas Gadjah Mada.
- Nakamura, T., & Mizuno, S., 2010. The discovery of hepatocyte growth factor (HGF) and its significance for cell biology, life science and clinical medicine. *Proc Jpn Acad Ser B Phys Biol Sci*. 86(6):588-610.
- Nathan, C. & Ding A., 2010. Nonresolving inflammation. *Cell*. 140(6):871-882.

- Onakpoya, I. J., Spencer, E. A., Thompson, M. J., Heneghan, C. J., 2015. The effect of chlorogenic acid on blood pressure: a systematic review and meta-analysis of randomized clinical trials. *J Hum Hypertens.* 29(2):77-81.
- Phanish, M. K., Winn, S. K., Dockrell, M. E., 2010. Connective tissue growth factor-(CTGF, CCN2)-a marker, mediator and therapeutic target for renal fibrosis. *Nephron Exp Nephrol.* 114(3):83-92.
- Pusat Data dan Informasi Kemenkes RI., 2013. *Situasi Penyakit Ginjal Kronis.* Jakarta: Kementerian Kesehatan RI.
- Ricardo, S. D., 2008. Macrophage diversity in renal injury and repair. *J Clin Invest.* 118(11):3522-3530.
- Rowe, R. G., Li, X. Y., Hu, Y., Saunders, T. L., Virtanen, I., De Herreros, G., *et al.*, 2009. Mesenchymal cells reactivate Snail1 expression to drive three-dimensional invasion programs. *J Cell Biol.* 184(3):399-408.
- Sato, Y., Itagaki, S., Kurokawa, T., Ogura, J., Kobayashi, M., Hirano, T., *et al.*, 2011. In vitro and in vivo antioxidant properties of chlorogenic acid and caffeic acid. *Int J Pharm.* 403(1-2):136-138.
- Schrimpf, C., Duffield, J. S., 2011. Mechanisms of fibrosis: the role of the pericyte. *Curr Opin Nephrol Hypertens.* 20(3):297-305.
- Schrimpf, C., Teebken, O. E., Wilhelmi, M., Duffield, J. S., 2014. The role of pericyte detachment in vascular rarefaction. *J Vasc Res.* 51(4):247-258.
- Schroder, K. & Tschopp, J., 2010. The inflammasomes. *Cell.* 140(6):821-832.
- Shao, D. D., Suresh, R., Vakil, V., Gomer, R. H., Pilling, D., 2008. Pivotal Advance: Th-1 cytokines inhibit, and Th-2 cytokines promote fibrocyte differentiation. *J Leukoc Biol.* 83(6):1323-1333.
- Shi, H., Dong, L., Jiang, J., Zhao, J., Zhao, G., Dang, X., *et al.*, 2013. Chlorogenic acid reduces liver inflammation and fibrosis through inhibition of toll-like receptor 4 signaling pathway. *Toxicology.* 303:107-114.
- Shi, H., Dong, L., Jiang, J., Zhao, J., Zhao, G., Dang, X., *et al.*, 2009. Chlorogenic acid against carbon tetrachloride-induced liver fibrosis in rats. *Eur J Pharmacol.* 623(1-3):119-124.
- Shin, H. S., Satsu, H., Bae, M. J., Zhao, Z., Ogiwara, H., Totsuka, M., *et al.*, 2015. Anti-inflammatory effect of chlorogenic acid on the IL-8 production in Caco-2 cells and the dextran sulphate sodium-induced colitis symptoms in C5BL/6 mice. *Food Chem.* 168:167-175.
- Strutz, F. & Müller, G. A., 2006. Renal fibrosis and the origin of the renal fibroblast. *Nephrol Dial Transplant.* 21(12):3368-3370.
- Tanaka, M., Endo, S., Yanagita, M., Economides, A. N., Valenzuela, D. M., Murphy, A. J., *et al.*, 2008. Expression of BMP-7 and USAG-1 (a BMP antagonist) in kidney development and injury. *Kidney Int.* 73(2):181-191.
- Thiery, J. P., Aclogue, H., Huang, R. Y., Nieto, M. A., 2009. Epithelial-mesenchymal transitions in development and disease. *Cell.* 139(5):871-890.
- Thomas, R., Kanso, A., Sedor, J. R., 2008. Chronic Kidney Disease and its Complications. *Prim care.* 35(2):329-344.
- Tomasek, J. J., Gabiani, G., Hinz, B., Chaponnier, C., Brown, Robert, A., 2002. Myofibroblasts and mechano-regulation of connective tissue remodelling. *Nat Rev Mol Cell Biol.* 3(5):349-363.

- Ucero, A. C., Benito, M. A., Fuentes, C. I., Santamaria, B., Blanco, J., Lopez, N. J.M., 2013. TNF-relates weak inducer of apoptosis (TWEAK) promotes kidney fibrosis and Ras-dependent proliferation of cultured rena fibroblast. *Biochim Biophys Acta*. 1832(10):1744-1755.
- Ucero, A. C., Benito, M. A., Izquierdo, M. C., Maria, D., Nino, S., Sanz, A. B., *et al.*, 2014. Unilateral ureteral obstruction: beyond obstruction. *Int Urol Nephrol*. 46:765-776.
- Vernon, M. A., Mylonas, K. J., Hughes, J., 2010. Macrophages and renal fibrosis. *Semin Nephrol*. 30(3):302-317.
- Vielhauer, V., Kulkarni, O., Reichel, C. A., Anders, H. J., 2010. Targeting the recruitment of monocytes and macrophages in renal disease. *Semin Nephrol*. 30(3):318-333.
- Vukicevic, S., Basic, V., Rogic, D., Basic, N., Shih, M. S., Shepard, A., *et al.*, 1998. Osteogenic protein-1 (bone morphogenetic protein-7) reduces severity of injury after ischemic acute renal failure in rat. *J Clin Invest*. 102(1):202-214.
- Wang, S. & Hirschberg, R., 2003. BMP7 antagonizes TGF-beta-dependent fibrogenesis in mesangial cells. *Am J Physiol Renal Physiol*. 284(5):1006-1013.
- Wang, Y. & Harris, D. C., 2011. Macrophages in renal disease. *J Am Soc Nephrol*. 22(1):21-27.
- Wu, Y., Deng, J., Rychahou, P. G., Qiu, S., Evers, B. M., Zhou, B. P., 2009. Stabilization of snail by NF-kappaB is required for inflammation-induced cell migration and invasion. *Cancer Cell*. 15(5):416-428.
- Wynn, T. A., 2008. Cellular and molecular mechanisms of fibrosis. *J Pathol*. 214(2):199-210.
- Yamada, S., Nakamura, J., Asada, M., Takase, M., Matsusaka, T., Iguchi, T., 2014. Twisted gastrulation, a BMP antagonist, exacerbates podocyte injury. *PLoS One*. 9(2):e89135.
- Yanagita, M., Okuda, T., Endo, S., Tanaka, M., Takahashi, K., Sugiyama, F., *et al.*, 2006. Uterine sensitization-associated gene-1 (USAG-1), a novel BMP antagonist expressed in the kidney, accelerates tubular injury. *J Clin Invest*. 116(1):70-79.
- Yang, F., Luo, L., Zhu, Z., Zhou, X., Wang, Y., Xue, J., 2017. Chlorogenic Acid Inhibits Liver Fibrosis by Blocking the miR-21-Regulated TGF- β 1/Smad7 Signaling Pathway *in Vitro* and *in Vivo*. *Front Pharmacol*. 8:929.
- Yang, J., Dai, C., Liu, Y., 2003. Hepatocyte growth factor supresses renal interstitial myofibroblast activation and intercepts Smad signal transduction. *Am J Pathol*. 163(2):621-632.
- Yo, Y., Morishita, R., Yamamoto, K., 2008. Action of hepatocyte growth factor as a local modulator in the kidney: Potential role in pathogenesis of renal disease. *Kidney Int*. 53:50-58.
- Yoo, K. H., Thornhill, B. A., Forbes, M. S., Coleman, C. M., Marcinko, E. S., Liaw, L., *et al.*, 2006. Osteopontin regulates renal apoptosis and interstitial fibrosis in neonatal chronic unilateral ureteral obstruction. *Kidney Int*. 70(10):1735-1741.

- Zeisberg, M. & Neilson, E. G., 2010. Mechanisms of tubulointerstitial fibrosis. *J. Am Soc Nephrol.* 21(11):1819–1834.
- Zeisberg, M., Hanai, J., Sugimoto, H., Mammoto, T., Charytan, D., Strutz, F., *et al.*, 2003. BMP-7 counteracts TGF-beta1-induced epithelial-to-mesenchymal transition and reverses chronic renal injury. *Nat Med.* 9(7):964–968.
- Zeisberg, M., Shah, A. A., Kalluri, R., 2005. Bone morphogenic protein-7 induces mesenchymal to epithelial transition in adult renal fibroblasts and facilitates regeneration of injured kidney. *J Biol Chem.* 280(9):8094-8100.
- Zhang, X. & Liu, Y., 2003. Supression of HGF receptor gene expression by oxidative stress is mediated through the iterplay between Sp1 and Egr-1. *Am J Physiol Renal Physiol.* 284:F1216-F1225.
- Zhang, X., Li, Y., Dai, C., Yang, J., Munder, P., Liu, Y., 2003. Sp1 and Sp3 transcription factor synergistically regulate HGF receptor gene expression in kidney. *Am J Physiol Renal Physiol.* 284(11):620-628.