

DAFTAR PUSTAKA

- American Academy of Ophthalmology. 2016. *Diagnostic Approach to Retinal Disease in Fundamentals and Principles of Ophthalmology*, San Francisco, 49–51.
- American Academy of Ophthalmology. 2016. *The Eye in Fundamentals and Principles of Ophthalmology*, San Francisco, 122–126.
- Ahn, H.C., Son, H.W., Kim, J.S. & Lee, J.H. 2005. Quantitative Analysis of Retinal Nerve Fiber Layer Thickness of Normal Children and Adolescents. *Korean J. Ophthalmol.*, 19: 195-200.
- Alhaddad, C., Barikian, A., Jaroudi, M., Massoud, V., Tamim, H. & Nouredin, B. 2014 Spectral domain optical coherence tomography in children: normative data and biometric correlations. *B. M. C.*, 14: 1471-2415.
- Alhaddad, C., Antonius, R., Tamim, H., Nouredin, B., 2014. Interocular symmetry in retinal and optic nerve parameters in children as measured by spectral domain optical coherence tomography. *Br. J. Ophthalmol.*, 98(4):502-506.
- Alasil, Wang, Keane, Lee, Baniasadi, Boer, D. & Chen. 2013. Analysis of normal retinal nerve fiber layer thickness by age, sex, and race using spectral domain optical coherence tomography. *J. Glaucoma.*, 22: 532-541.
- Allingham. 2011. *Optic Nerve, Retina, and Choroid, in: Shields Textbook of Glaucoma*. Lippincott Williams & Wilkins, Philadelphia.
- Altemir, I., Pueyo, V., Elia, N., Polo, V., Larrosa, J.M. & Oros, D. 2013. Reproducibility of optical coherence tomography measurements in children. *Am. J. Ophthalmol.*, 155: 171-176.
- Barrio-Barrio, J., 2013. Multicenter spanish study of spectral-domain tomography in normal children. *Acta Ophthalmol.*, 91(1):56-63.
- Brancato, R., Lumbroso, B. 2004. *Guide to Optical Coherence Tomography Interpretation*. Innovation-News-Communication, Italy
- Cook, R. and Glasscock, R. 1951. Refractive and Ocular Findings in the Newborn, *Am. J. Ophthalmol.*, 34(10): 1407–13.
- Cheng, S, C. K., Lam, C. Y. M., Yap, M. K. H. 2010. Retinal thickness in myopic and non myopic eyes. *Ophthalmic. Physiol. Opt.*, 30:776-784.
- Choi, S. W., Lam, S., 2006. Thickness changes in fovea and peripapillary retinal nerve fiber layer depend on the degree of myopia, *Korean J. Ophthalmol.*, 20(4):215-219.
- Curtin, B.J., 1977. The posterior staphyloma of pathologic myopia. *Trans. Am. Ophthalmol. Soc.*, 75:67-86
- Dubis, M. K., 2009. Reconstructing foveal pit morphology from optical coherence tomography imaging. *Br. J. Ophthalmol.*, 93(9):1223-1227.
- Elder, D. 1970. *Ophthalmic Optics and Refraction*, The CV Mosby Company. 300-351
- Fercher, A.F., 1996. Optical coherence tomography. *Journal of Biomedical Optics*, 1(2), pp.157-174.
- Fujiwara, T., Imamura, Y., Margolis, R. 2009. Enhanced depth imaging optical coherence tomography of the choroid in highly myopic eyes. *Am. J. Ophthalmol.*, 148:445-50

- Garcia, G.E., 1989. Handbook of Refraction. Little, Brown and Company. 32-33
- Gao, L.Q., Liu, W., Liang Y.B., 2011. Prevalence and characteristics of myopic retinopathy in a rural Chinese adult population: the Handan Eye Study. *Arch. Ophthalmol.*, 129:1199-204
- Gerardo, M., Ohlsson, J., Abrahamsson, M., Sjostrom, A. and Sjostrand, J. 2000. Myopisation: The Refractive Tendency in Teenagers. Prevalence of Myopia Among Young Teenagers in Sweden, *Acta Ophthalmol.*, 78(2):177-181.
- Gordon, R. and Donzis, P., 1985. Refractive Development of the Human Eye, *Arch. Ophthalmol.*, 103(6):785-9.
- Goss, D.A. 1987. Myopia. In Diagnosis and Management in Vision Care. Butterworth, USA, 121-162
- Gurses-Ozden, R., Teng, C., Vessani, R., Zafar, S., Liebmann, J. M., Ritch, R. 2004. Macular and retinal nerve fiber layer thickness measurement reproducibility using optical coherence tomography (OCT-3). *J. Glaucoma*, 13: 238-244
- Hee, M. R., Puliafito, C. A., Wong, C., Duker, J. S., Reichel, E., Rutledge, B., Schuman, J. S., 1995. Optical coherence tomography of human retina. *Arch. Ophthalmol.*, 113(3):325-332.
- Hendrianto, A., Agni, A.N, Supartoto, A. 2016. Angka Normal Ketebalan Nerve Fiber Layer Peripapil dan Makula pada Anak Sehat di Yogyakarta. Tesis. (unpublished)
- Hendrickson, A.E. and Yuodelis, C., 1984. The morphological development of the human fovea. *Ophthalmology*, 91(6), pp.603-612.
- Huang, D., Swanson, E.A., Lin, C.P., Schuman, J.S., Stinson, W.G., Chang, W., Hee, M.R., Flotte, T., Gregory, K. and Puliafito, C.A., 1991. Optical coherence tomography. *Science*, 254(5035), pp.1178-1181.
- Hwuang, Y. H., Kim. Y., 2012. Macular thickness and volume of myopic eyes measured using spectral domain coherence tomography. *Clin. Exp. Opt.*, 95(5):492-498.
- Ikuno, Y., Ohji, M. 2018 'High Myopia and Vitreoretinal complications', in *Ryan's Retina*, 2:2146-2154.
- Kobayashi, K., Ohno-Matsui, K., Kojima, M., 2005. Fundus Characteristic of High Myopia in Children. *Jpn. J. Ophthalmol.*, 49:306-11
- Lam, D.S., Leung, K. M. S., Mohamed, S., Chan, W. M., Palanivelu, M. S., Cheung, C. Y., Li, E. Y. 2007. Regional variations in the relationship between macular thickness measurements and myopia. *Invest. Ophthalmol. Vis. Sci.*, 48:376-382.
- Lee, J. W. Y., Yau, G. S. K., Woo, T. T. Y., Wong, R. L. M., & Wong, I. Y. H. 2015. Central Macular Thickness in Children with Myopia, Emmetropia, and Hyperopia: An Optical Coherence Tomography Study. *BioMed Res. Int.*, 29: 200-205
- Liang, H., Crewther, D. P., Crewther, S. G., Barila, A. M., 2000. A role of photoreceptor outer segments in the induction of deprivation myopia in tree shrews. *Vision Res*, 40:2369-2377.
- Lim, M.C., Hoh, S.T., Foster, P.J., Lim, T.H., Chew, S.J., Scah, S.K., 2005. Use of optical coherence tomography to assess variations in macular retinal thickness in myopia. *Invest. Ophthalmol. Vis. Sci.*, 46: 974-978

- Liu, H.H. 2010. Prevalence and progression of myopic retinopathy in Chinese adults: the Beijing Eye Study. *Ophthalmology*, 117:1763-68
- Luo, H. D., Gazzard, G., Fong, A., Aung, T., Hoh, S. T., Loon, S. C. 2006. Myopia, axial length, and OCT characteristics of macula in Singaporean children. *Invest. Ophthalmol. Vis. Sci.*, 47:2773-2781.
- McBrien, N.A., Gentle, A. 2003. Role of the sclera in the development and pathological complications of myopia. *Prog. Retin. Eye. Res.* 22:307-38
- Morgan, I. G., Ohno-Matsui, K. and Mei Saw, S. 2012. Myopia. *Lancet*, 379: 1739–48.
- Ohno-Matsui, K., Shimada, N., Yasuzumi, K. 2011. Long-term development of significant visual field defects in highly myopic eyes. *Am. J. Ophthalmol.*, 152: 256-65
- Pados, S.M., Yanoff, M., 1994. Spectacle Lens. In: Textbook of Ophthalmology. London, Mosby-Year Book Europe, Ltd. 9: 102-109
- Pang, Y., Goodfellow, G. W., Allison, C., Block, S., Franz, K. A., 2011. A prospective study of macular thickness in amblyopic children with unilateral high myopia. *Invest. Ophthalmol. Vis. Sci.*, 52:2444-2449.
- Panazzo G, M. A., 2004. Optical coherence tomography findings in myopic traction maculopathy. *Arch. Ophthalmol.*, 122:1455-1460.
- Salchow, D.J., Oleynikov, Y.S., Chiang, M.F., Salchow, S.E.K., Langton, K., Tsai, J.C. & Al-Aswad, L.A. 2006. Retinal nerve fiber layer thickness in normal children measured with optical coherence tomography. *Am. J. Ophthalmol.*, 113, 786-791.
- Samarawickrama, C., Pai, A., Huynh, S. C., Burlustky, G., Wong. T.Y., Mitchell, P., 2010. Influence of OCT signal strength on Macular, Optic Nerve Head, and Retinal Nerve Fiber Layer Parameters. *Invest. Ophthalmol. Vis. Sci.*, 51: 4471-4475.
- Saxena, S., Meredith, T. A., 2006. *Optical Coherence Tomography in Retinal Diseases*. Jaypee Brothers Medical Publisher. New Delhi.
- Saw, S. M., Chua, W.H., Hong, C.Y., *et al.*, 2002. Nearwork in early onset myopia. *Invest. Ophthalmol. Vis. Sci.*, 43, 332-339.
- Schachat, A., Wilkinson, C. and Hinton, D., 2017. *Myopic Macular Degeneration, in Ryan's Retina*: 1326–30.
- Song W. K., Lee, S. C., Lee, E. S., Kim, C. Y., Kim, S. S., 2010. Macular thickness variations with sex, age, and axial length in healthy subjects: a spectral domain optical coherence tomography study. *Invest. Ophthalmol. Vis. Sci.*, 51(8):3913-3918.
- Sugar, E.A., Jabs, D.A., Altaweel, M.M., Lightman, S., Acharya, N., Vitale, A.T., Thorne, J.E., 2011. Identifying a Clinically Meaningful Threshold for Change in Uveitic Macular Edema Evaluated by Optical Coherence Tomography. *Am. J. Ophthalmol.* 152(6):1044-1052
- Turk A., Ceylan O. M., Arici C., *et al.* 2012. Evaluation of the nerve fiber layer and macula in the eyes of healthy children using spectral-domain optical coherence tomography. *Am. J. Ophthalmol.* 153(3):552–559
- Wakitani, Y., Sasoh, M., Sugimoto, M., Ito, Y., Ido, M. and Uji, Y., 2003. Macular Thickness Measurements in Healthy Subjects with Different Axial Lengths

Using Optical Coherence Tomography, *Retina*, 23: 177–182.

Wu, P.C., Chen, Y.J., Chen, C.H., Chen, Y.H., Shin, S.J., Yang, H.J., Kuo, H.K., 2008. Assessment of macular retinal thickness and volume in normal eyes and highly myopic eyes with third-generation optical coherence tomography. *Eye (Lond)*, 2008;22(4):551–555

Ziylan, S., Kiziloglu, O. Y., Yenerel, N. M., Gokce, B., Ciftci, F., 2015. Macular thickness in highly myopic children aged 3 to 7 years. *J. Pediatr. Ophthalmol. Strabismus*, 52(4):502-506.

Zhao, M., Wu, Q., Hu, P. and Jia, L., 2015. Macular Thickness Assessed with Optical Coherence Tomography in Young Chinese Myopic Patients, *J. Ophthalmol.*, 29: 43-54