

## KEPUSTAKAAN

- Allen, S.J., Watson, J.J., Shoemark, D.K., Barua, N.U., Patel, N.K., 2013. GDNF, NGF and BDNF as therapeutic options for neurodegeneration. *Pharmacol. Ther.* 138, 155–175. doi:10.1016/j.pharmthera.2013.01.004
- Aranha, A.M.F., Zhang, Z., Neiva, K.G., Costa, C.A.S., Hebling, J., Nör, J.E., 2010. Hypoxia enhances the angiogenic potential of human dental pulp cells. *J. Endod.* 36, 1633–1637. doi:10.1016/j.joen.2010.05.013
- Barzilai, A., Yamamoto, K.I., 2004. DNA damage responses to oxidative stress. *DNA Repair (Amst)*. 3, 1109–1115. doi:10.1016/j.dnarep.2004.03.002
- Bathina, S., Das, U.N., 2015. Brain-derived neurotrophic factor and its clinical Implications. *Arch. Med. Sci.* 11, 1164–1178. doi:10.5114/aoms.2015.56342
- Baziyar, Y., Edalatmanesh, M.A., Hosseini, S.A., Zar, A., 2016. The effects of endurance training and gallic acid on BDNF and TNF- $\alpha$  in male rats with Alzheimer. *Int. J. Appl. Exerc. Physiol.* 5, 45–54.
- Block, F., 1999. Global ischemia and behavioural deficits. *Prog. Neurobiol.* 58, 279–295. doi:10.1016/S0301-0082(98)00085-9
- Bono, F., Lamarche, I., Bornia, J., Savi, P., Valle, G. Della, Herbert, J.-M., 1999. Nerve growth factor (NGF) exerts its pro-apoptotic effect via the P75 NTR receptor in a cell cycle-dependent manner. *FEBS Lett.* 457, 93–97.
- Bouet, V., Freret, T., Ankri, S., Bezault, M., Renolleau, S., Boulouard, M., Jacotot, E., Chauvier, D., Schumann-Bard, P., 2010. Predicting sensorimotor and memory deficits after neonatal ischemic stroke with reperfusion in the rat. *Behav. Brain Res.* 212, 56–63. doi:10.1016/j.bbr.2010.03.043
- Boyce, R.W., Dorph-Petersen, K.A., Lyck, L., Gundersen, H.J.G., 2010. Design-based stereology: introduction to basic concepts and practical approaches for estimation of cell number. *Toxicol. Pathol.* 38, 1011–1025. doi:10.1177/0192623310385140
- Bromley-Brits, K., Deng, Y., Song, W., 2011. Morris water maze test for learning and memory deficits in Alzheimer's disease model mice. *J. Vis. Exp.* 2–6. doi:10.3791/2920
- Buravkova, L.B., Andreeva, E.R., Gogvadze, V., Zhivotovsky, B., 2014. Mesenchymal stem cells and hypoxia: Where are we? *Mitochondrion* 19, 105–112. doi:10.1016/j.mito.2014.07.005
- Busletta, C., Novo, E., Valfrè Di Bonzo, L., Povero, D., Paternostro, C., Ievolella, M., Mareschi, K., Ferrero, I., Cannito, S., Compagnone, A., Bandino, A., Colombatto, S., Fagioli, F., Parola, M., 2011. Dissection of the biphasic nature of hypoxia-induced motogenic action in bone marrow-derived human mesenchymal stem cells. *Stem Cells* 29, 952–963. doi:10.1002/stem.642
- Cantinieux, D., Quertainmont, R., Blacher, S., Rossi, L., Wanet, T., Noël, A., Brook, G., Schoenen, J., Franzen, R., 2013. Conditioned medium from bone marrow-derived mesenchymal stem cells improves recovery after spinal cord injury in rats: An original strategy to avoid cell transplantation. *PLoS One* 8, e69515. doi:10.1371/journal.pone.0069515
- Ceccariglia, S., D'Altocolle, A., Del Fa', A., Pizzolante, F., Caccia, E., Michetti, F., Gangitano, C., 2011. Cathepsin D plays a crucial role in the trimethyltin-induced hippocampal neurodegeneration process. *Neuroscience* 174, 160–

170. doi:10.1016/j.neuroscience.2010.11.024
- Chacko, S.M., Ahmed, S., Selvendiran, K., Kuppusamy, M.L., Khan, M., Kuppusamy, P., 2010. Hypoxic preconditioning induces the expression of pro-survival and proangiogenic markers in mesenchymal stem cells. *Am. J. Physiol. Cell Physiol.* 299, C1562–C1570. doi:10.1152/ajpcell.00221.2010
- Chang, C.-P., Chio, C.-C., Cheong, C.-U., Chao, C.-M., Cheng, B.-C., Lin, M.-T., 2013. Hypoxic preconditioning enhances the therapeutic potential of the secretome from cultured human mesenchymal stem cells in experimental traumatic brain injury. *Clin. Sci.* 124, 165–76. doi:10.1042/CS20120226
- Chang, W., Song, B.W., Lim, S., Song, H., Chi, Y.S., Cha, M.J., Dong, H.A., Jung, Y.G., Lee, D.H., Ji, H.C., Choi, K.D., Lee, S.K., Chung, N., Lee, S.K., Jang, Y., Hwang, K.C., 2009. Mesenchymal stem cells pretreated with delivered Hsp-1-Hsp70 protein are protected from hypoxia-mediated cell death and rescue heart functions from myocardial injury. *Stem Cells* 27, 2283–2292. doi:10.1002/stem.153
- Charan, J., Kantharia, N.D., 2013. How to calculate sample size in animal studies? *J. Pharmacol. Pharmacother.* 4, 303–6. doi:10.4103/0976-500X.119726
- Corvino, V., Geloso, M.C., Cavallo, V., Guadagni, E., Passalacqua, R., Florenzano, F., Giannetti, S., Molinari, M., Michetti, F., 2005. Enhanced neurogenesis during trimethyltin-induced neurodegeneration in the hippocampus of the adult rat. *Brain Res. Bull.* 65, 471–477. doi:10.1016/j.brainresbull.2005.02.031
- Crisostomo, P.R., Wang, Y., Markel, T.A., Wang, M., Lahm, T., Meldrum, D.R., 2008. Human mesenchymal stem cells stimulated by TNF- $\alpha$ , LPS, or hypoxia produce growth factors by an NF $\kappa$ B- but not JNK-dependent mechanism. *Am. J. Physiol. Cell Physiol.* 294, C675–C682. doi:10.1152/ajpcell.00437.2007
- Cunha, C., Brambilla, R., Thomas, K.L., 2010. A simple role for BDNF in learning and memory? *Front. Mol. Neurosci.* 3, 1–14. doi:10.3389/neuro.02.001.2010
- da Silva Meirelles, L., Fontes, A.M., Covas, D.T., Caplan, A.I., Meirelles, L.D.S., Fontes, A.M., Covas, D.T., Caplan, A.I., Da, L., Meirelles, S., Fontes, A.M., Covas, D.T., Caplan, A.I., 2009. Mechanisms involved in the therapeutic properties of mesenchymal stem cells. *Cytokine Growth Factor Rev.* 20, 419–427. doi:10.1016/j.cytogfr.2009.10.002
- Dalous, J., Larghero, J., Baud, O., 2012. Transplantation of umbilical cord-derived mesenchymal stem cells as a novel strategy to protect the central nervous system: technical aspects, preclinical studies, and clinical perspectives. *Pediatr. Res.* 71, 482–490. doi:10.1038/pr.2011.67
- Dember, W.N., Fowler, H., 1958. Spontaneous alternation behavior. *Psychol. Bull.* 55, 412–428. doi:10.1007/BF02686644
- Deschepper, M., Oudina, K., David, B., Myrtil, V., Collet, C., Bensidhoum, M., Logeart-Avramoglou, D., Petite, H., 2011. Survival and function of mesenchymal stem cells (MSCs) depend on glucose to overcome exposure to long-term, severe and continuous hypoxia. *J. Cell. Mol. Med.* 15, 1505–1514. doi:10.1111/j.1582-4934.2010.01138.x

- Dominici, M., Le Blanc, K., Mueller, I., Slaper-Cortenbach, I., Marini, F.C., Krause, D.S., Deans, R.J., Keating, A., Prockop, D.J., Horwitz, E.M., 2006. Minimal criteria for defining multipotent mesenchymal stromal cells. The International Society for Cellular Therapy position statement. *Cytotherapy* 8, 315–317. doi:10.1080/14653240600855905
- Drago, D., Cossetti, C., Iraci, N., Gaude, E., Musco, G., Bachi, A., Pluchino, S., 2013. The stem cell secretome and its role in brain repair. *Biochimie* 95, 2271–2285. doi:10.1016/j.biochi.2013.06.020
- Drela, K., Sarnowska, A., Siedlecka, P., Szablowska-Gadomska, I., Wielgos, M., Jurga, M., Lukomska, B., Domanska-Janik, K., 2014. Low oxygen atmosphere facilitates proliferation and maintains undifferentiated state of umbilical cord mesenchymal stem cells in an hypoxia inducible factor-dependent manner. *Cytotherapy* 16, 881–892. doi:10.1016/j.jcyt.2014.02.009
- Dyachok, J., Earnest, S., Iturraran, E.N., Cobb, M.H., Ross, E.M., 2016. Amino acids regulate mTORC1 by an obligate two-step mechanism. *J. Biol. Chem.* 291, 22414–22426. doi:10.1074/jbc.M116.732511
- Ebert, R., Ulmer, M., Zeck, S., Meissner-Weigl, J., Schneider, D., Stopper, H., Schupp, N., Kassem, M., Jakob, F., 2006. Selenium supplementation restores the antioxidative capacity and prevents cell damage in bone marrow stromal cells in vitro. *Stem Cells* 24, 1226–1235. doi:10.1634/stemcells.2005-0117
- Efimenko, A., Starostina, E., Kalinina, N., Stolzing, A., 2011. Angiogenic properties of aged adipose derived mesenchymal stem cells after hypoxic conditioning.pdf 1–13.
- Eichenbaum, H., Stewart, C., Morris, R.G.M., 1990. Hippocampal representation in place learning. *J. Neurosci.* 10, 3531–3542. doi:10.1523/jneurosci.1068-06.2006
- Ernfors, P., Ibáñez, C.F., Ebendal, T., Olson, L., Persson, H., 1990. Molecular cloning and neurotrophic activities of a protein with structural similarities to nerve growth factor: developmental and topographical expression in the brain. *Proc. Natl. Acad. Sci. U. S. A.* 87, 5454–5458. doi:10.1073/pnas.87.14.5454
- Fan, C.-G., Zhang, Q., Zhou, J., 2011. Therapeutic potentials of mesenchymal stem cells derived from human umbilical cord. *Stem Cell Rev. Reports* 7, 195–207. doi:10.1007/s12015-010-9168-8
- Forsythe, J.O.A., Jiang, B., Iyer, N. V., Agani, F., Leung, S.W., 1996. Activation of vascular endothelial growth factor gene transcription by hypoxia-inducible factor 1. *Mol. Cell. Biol.* 16, 4604–4612. doi:10.1128/MCB.16.9.4604
- Friedenstein, A.J., Piatetzky-Shapiro, I.I., Petrakova, K.V., 1966. Osteogenesis in transplants of bone marrow cells. *J. Embryol. Exp. Morphol.* 16, 381–390.
- Gage, F.H., 2000. Mammalian neural stem cells. *Science* (80-. ). 287, 1433–1438. doi:10.1126/science.287.5457.1433
- Galati, G., Pelle, G., Berthoz, A., Comitteri, G., 2010. Multiple reference frames used by the human brain for spatial perception and memory. *Exp. Brain Res.* 206, 109–120. doi:10.1007/s00221-010-2168-8
- Gasparova, Z., Janega, P., Stara, V., Ujhazy, E., Ujhazy, A.E., 2012. Early and late stage of neurodegeneration induced by trimethyltin in hippocampus and

- cortex of male Wistar rats. *Neuroendocrinol. Lett.* 33, 689–696.
- Geloso, M.C., Corvino, V., Michetti, F., 2011. Trimethyltin-induced hippocampal degeneration as a tool to investigate neurodegenerative processes. *Neurochem. Int.* 58, 729–738. doi:10.1016/j.neuint.2011.03.009
- Ghosh, A., Carnahan, J., Greenberg, M.E., 1994. Requirement for BDNF in activity-dependent survival of cortical neurons. *Science* (80-. ). 263, 1618–1623. doi:10.1126/science.7907431
- Gögel, S., Gubernator, M., Minger, S.L., 2011. Progress and prospects: Stem cells and neurological diseases. *Gene Ther.* 18, 1–6. doi:10.1038/gt.2010.130
- Gómez-Palacio-Schjetnan, A., Escobar, M.L., 2013. Neurotrophins and synaptic plasticity. *Curr. Top. Behav. Neurosci.* 15, 117–36. doi:10.1007/7854
- Govindasamy, V., Ronald, V.S., Totey, S., Binti Din, S., Mustafa, W.M.B.W., Totey, S., Zakaria, Z., Bhonde, R.R., 2010. Micromanipulation of culture niche permits long-term expansion of dental pulp stem cells-an economic and commercial angle. *Vitr. Cell. Dev. Biol. - Anim.* 46, 764–773. doi:10.1007/s11626-010-9332-0
- Grayson, W.L., Zhao, F., Bunnell, B., Ma, T., 2007. Hypoxia enhances proliferation and tissue formation of human mesenchymal stem cells. *Biochem. Biophys. Res. Commun.* 358, 948–953. doi:10.1016/j.bbrc.2007.05.054
- Gundersen, H.J.G., 1977. Notes on the estimation of numerical density of arbitrary profiles: the edge effect. *J. Microsc.* 111, 219–223.
- Guzy, R.D., Hoyos, B., Robin, E., Chen, H., Liu, L., Mansfield, K.D., Simon, M.C., Hammerling, U., Schumacker, P.T., 2005. Mitochondrial complex III is required for hypoxia-induced ROS production and cellular oxygen sensing. *Cell Metab.* 1, 401–408. doi:10.1016/j.cmet.2005.05.001
- Guzy, R.D., Schumacker, P.T., 2006. Oxygen sensing by mitochondria at complex III: The paradox of increased reactive oxygen species during hypoxia. *Exp. Physiol.* 91, 807–819. doi:10.1113/expphysiol.2006.033506
- Ha, J.S., Jin, D.E., Park, S.K., Park, C.H., Seung, T.W., Bae, D.-W., Kim, D.-O., Heo, H.J., 2015. Antiamnesic effect of *Actinidia arguta* extract intake in a mouse model of TMT-induced learning and memory dysfunction. *Evidence-Based Complement. Altern. Med.* 2015, 1–13.
- Hagmann, S., Moradi, B., Frank, S., Dreher, T., Kämmerer, P.W., Richter, W., Gotterbarm, T., 2013. Different culture media affect growth characteristics, surface marker distribution and chondrogenic differentiation of human bone marrow-derived mesenchymal stromal cells. *BMC Musculoskelet. Disord.* 14, 1–7. doi:10.1186/1471-2474-14-223
- Hayashi, M., Mitsunaga, F., Ohira, K., Shimizu, K., Yamashita, A., 1999. Development of full-length Trk B-immunoreactive structures in the hippocampal formation of the macaque monkey. *Anat. Embryol.* 199, 529–537.
- Hofer, M., Pagliusi, S.R., Hohn, A., Leibrock, J., Barde, Y.-A., 1990. Regional distribution of brain-derived neurotrophic factor mRNA in the adult mouse brain-derived neurotrophic factor/hippocampus/ in situ hybridization/nerve growth factor/trophic interactions. *EMBO J.* 9, 2459–2464.

- Holmes, D.I.R., Zachary, I., 2005. The vascular endothelial growth factor (VEGF) family: angiogenic factors in health and disease. *Genome Biol.* 6, 1–10. doi:10.1186/gb-2005-6-2-209
- Hsieh, J.-Y., Wang, H.-W., Chang, S.-J., Liao, K.-H., Lee, I.-H., Lin, W.-S., Wu, C.-H., Lin, W.-Y., Cheng, S.-M., 2013. Mesenchymal stem cells from human umbilical cord express preferentially secreted factors related to neuroprotection, neurogenesis, and angiogenesis. *PLoS One* 8, 1–11. doi:10.1371/journal.pone.0072604
- Hsieh, P.-S., Bochinski, D.J., Lin, G.T., Nunes, L., Lin, C.S., Lue, T.F., 2003. The effect of vascular endothelial growth factor and brain-derived neurotrophic factor on cavernosal nerve regeneration in a nerve-crush rat model. *BJU Int.* 92, 470–475. doi:10.1046/j.1464-410X.2003.04373.x
- Jin, K., Zhu, Y., Sun, Y., Mao, X.O., Xie, L., Greenberg, D.A., 2002. Vascular endothelial growth factor (VEGF) stimulates neurogenesis in vitro and in vivo. *Proc Natl Acad Sci U S A* 99, 11946–11950. doi:10.1073/pnas.182296499
- Joerger-Messerli, M.S., Marx, C., Oppliger, B., Mueller, M., Surbek, D. V., Schoeberlein, A., 2016. Mesenchymal stem cells from Wharton's jelly and amniotic fluid. *Best Pract. Res. Clin. Obstet. Gynaecol.* 31, 30–44. doi:10.1016/j.bpobgyn.2015.07.006
- Johnson, E., Adamo-Villani, N., 2010. A study of the effects of immersion on short-term spatial memory. *World Acad. Sci. Eng. Technol.* 71, 582–587.
- Jung, E.-Y., Lee, M.-S., Ahn, C.J., Cho, S.-H., Bae, H., Shim, I., 2013. The neuroprotective effect of gugijihwang-tang on trimethyltin-induced memory dysfunction in the rat. *Evidence-Based Complement. Altern. Med.* 2013, 1–6. doi:10.1155/2013/542081
- Kang, H., Schuman, E.M., 1995. Long-lasting neurotrophin-induced enhancement of synaptic transmission in the adult hippocampus. *Science* (80-. ). 267, 1658–1662.
- Kang, J.Y., Park, S.K., Guo, T.J., Ha, J.S., Lee, D.S., Kim, J.M., Lee, U., Kim, D.O., Heo, H.J., 2016. Reversal of trimethyltin-induced learning and memory deficits by 3, 5-dicaffeoylquinic acid. *Oxid. Med. Cell. Longev.* 2016, 1–13.
- Ke, X., Zhang, J., 2013. Changes in HIF-1  $\alpha$ , VEGF, NGF and BDNF levels in cerebrospinal fluid and their relationship with cognitive impairment in patients with cerebral infarction. *J. Huazhong Univ. Sci. Technol. [Medical Sci.* 33, 433–437.
- Kesner, R.P., Hopkins, R.O., 2006. Mnemonic functions of the hippocampus: A comparison between animals and humans. *Biol. Psychol., Translational Research: Parallels of Human and Animal Research in Biological Psychology* 73, 3–18. doi:10.1016/j.biopsycho.2006.01.004
- Kessels, R.P.C., Feijen, J., Postma, A., 2005. Implicit and explicit memory for spatial information in Alzheimer's disease. *Dement. Geriatr. Cogn. Disord.* 20, 184–191. doi:10.1159/000087233
- Kim, J.W., Tchernyshyov, I., Semenza, G.L., Dang, C. V., 2006. HIF-1-mediated expression of pyruvate dehydrogenase kinase: A metabolic switch required



- for cellular adaptation to hypoxia. *Cell Metab.* 3, 177–185. doi:10.1016/j.cmet.2006.02.002
- Kim, S.G., Buel, G.R., Blenis, J., 2013. Nutrient regulation of the mTOR complex 1 signaling pathway. *Mol. Cells* 35, 463–473. doi:10.1007/s10059-013-0138-2
- Koda, T., Kuroda, Y., Imai, H., 2008. Protective effect of rutin against spatial memory impairment induced by trimethyltin in rats. *Nutr. Res.* 28, 629–634. doi:10.1016/j.nutres.2008.06.004
- Korte, M., Carroll, P., Wolf, E., Brem, G., Thoenen, H., Bonhoeffer, T., 1995. Hippocampal long-term potentiation is impaired in mice lacking brain-derived neurotrophic factor. *Proc. Natl. Acad. Sci. U. S. A.* 92, 8856–8860. doi:10.1073/pnas.92.19.8856
- Korte, M., Staiger, V., Griesbeck, O., Thoenen, H., Bonhoeffer, T., 1996. The involvement of brain-derived neurotrophic factor in hippocampal long-term potentiation revealed by gene targeting experiments. *J. Physiol. Paris* 90, 157–164. doi:10.1016/S0928-4257(97)81415-5
- Kovalchuk, Y., Hanse, E., Kafitz, K.W., Konnerth, A., 2002. Postsynaptic induction of BDNF-mediated long-term potentiation. *Science* (80-. ). 295, 1729–1734. doi:10.1126/science.1067766
- Lavoie, J.R., Rosu-Myles, M., 2013. Uncovering the secrets of mesenchymal stem cells. *Biochimie* 95, 2212–2221. doi:10.1016/j.biochi.2013.06.017
- Lavrentieva, A., Majore, I., Kasper, C., Hass, R., 2010. Effects of hypoxic culture conditions on umbilical cord-derived human mesenchymal stem cells. *Cell Commun. Signal.* 8, 18. doi:10.1186/1478-811X-8-18
- Lee, B., Sur, B., Cho, S.-G., Yeom, M., Shim, I., Lee, H., Hahm, D.-H., 2016. Wogonin attenuates hippocampal neuronal loss and cognitive dysfunction in trimethyltin-intoxicated rats. *Biomol. Ther.* 24, 328–337. doi:10.4062/biomolther.2015.152
- Lee, C.Y., Kang, J.Y., Lim, S., Ham, O., Chang, W., Jang, D.-H., 2016. Hypoxic conditioned medium from mesenchymal stem cells promotes lymphangiogenesis by regulation of mitochondrial-related proteins. *Stem Cell Res. Ther.* 7, 38. doi:10.1186/s13287-016-0296-1
- Lee, E., Son, H., 2009. Adult hippocampal neurogenesis and related neurotrophic factors. *BMB Rep.* 42, 239–244. doi:10.5483/BMBRep.2009.42.5.239
- Lee, H.J., Ryu, J.M., Jung, Y.H., Oh, S.Y., Lee, S.J., Han, H.J., 2015. Novel pathway for hypoxia-induced proliferation and migration in human mesenchymal stem cells: Involvement of HIF-1 $\alpha$ , FASN, and mTORC1. *Stem Cells* 33, 2182–2195. doi:10.1002/stem.2020
- Lee, S.C., Kim, K.H., Kim, O.H., Lee, S.K., Hong, H.E., Won, S.S., Jeon, S.J., Choi, B.J., Jeong, W., Kim, S.J., 2017. Determination of optimized oxygen partial pressure to maximize the liver regenerative potential of the secretome obtained from adipose-derived stem cells. *Stem Cell Res. Ther.* 8, 1–12. doi:10.1186/s13287-017-0635-x
- Lee, S.H., Lee, Y.J., Song, C.H., Ahn, Y.K., Han, H.J., 2010. Role of FAK phosphorylation in hypoxia-induced hMSCS migration: involvement of VEGF as well as MAPKS and eNOS pathways. *Am. J. Physiol. Cell Physiol.*

298, C847–C856. doi:10.1152/ajpcell.00418.2009

- Levine, E.S., Dreyfus, C.F., Black, I.B., Plummer, M.R., 1995. Brain-derived neurotrophic factor rapidly enhances synaptic transmission in hippocampal neurons via postsynaptic tyrosine kinase receptors. *Proc. Natl. Acad. Sci.* 92, 8074–8077. doi:10.1073/pnas.92.17.8074
- Li, X., Bai, J., Ji, X., Li, R., Xuan, Y., Wang, Y., 2014. Comprehensive characterization of four different populations of human mesenchymal stem cells as regards their immune properties, proliferation and differentiation. *Int. J. Mol. Med.* 34, 695–704. doi:10.3892/ijmm.2014.1821
- Lin, C., McGough, R., Aswad, B., Block, J., Terek, R., 2004. Hypoxia induces HIF-1- $\alpha$  and VEGF expression in chondrosarcoma cells and chondrocytes. *J. Orthop. Res.* 22, 1175–1181. doi:10.1016/j.orthres.2004.03.002
- Lindholm, D., Carroll, P., Tzimagiorgis, G., Thoenen, H., 1996. Autocrine-paracrine regulation of hippocampal neuron survival by IGF-1 and the neurotrophins BDNF, NT-3 and NT-4. *Eur. J. Neurosci.* 8, 1452–1460. doi:10.1111/j.1460-9568.1996.tb01607.x
- Lindvall, O., Kokaia, Z., 2010. Stem cells in human neurodegenerative disorders — time for clinical translation? *J. Clin. Invest.* 120, 29–40. doi:10.1172/JCI40543.patients
- Liu, L., Gao, J., Yuan, Y., Chang, Q., Liao, Y., Lu, F., 2013. Hypoxia preconditioned human adipose derived mesenchymal stem cells enhance angiogenic potential via secretion of increased VEGF and bFGF. *Cell Biol. Int.* 37, 551–560. doi:10.1002/cbin.10097
- Lowenstein, D.H., Arsenault, L., 1996. The effects of growth factors on the survival and differentiation of cultured dentate gyrus neurons. *J. Neurosci.* 16, 1759–1769.
- Lührs, M.-L., Dammhahn, M., Kappeler, P.M., Fichtel, C., 2009. Spatial memory in the grey mouse lemur (*Microcebus murinus*). *Anim. Cogn.* 12, 599–609. doi:http://dx.doi.org/10.1007/s10071-009-0219-y
- Ma, Y.L., Wang, H.L., Wu, H.C., Wei, C.L., Lee, E.H., 1998. Brain-derived neurotrophic factor antisense oligonucleotide impairs memory retention and inhibits long-term potentiation in rats. *Neuroscience* 82, 957–967. doi:10.1016/S0306-4522(97)00325-4
- Majumdar, D., Bhonde, R., Datta, I., 2013. Influence of ischemic microenvironment on human Wharton's jelly mesenchymal stromal cells. *Placenta* 34, 642–649. doi:10.1016/j.placenta.2013.04.021
- Maltman, D.J., Hardy, S. a., Przyborski, S. a., 2011. Role of mesenchymal stem cells in neurogenesis and nervous system repair. *Neurochem. Int.* 59, 347–356. doi:10.1016/j.neuint.2011.06.008
- Martens, L.K., Kirschner, K.M., Warnecke, C., Scholz, H., 2007. Hypoxia-inducible factor-1 (HIF-1) is a transcriptional activator of the TrkB neurotrophin receptor gene. *J. Biol. Chem.* 282, 14379–14388. doi:10.1074/jbc.M609857200
- McAllister, A.K., Katz, L.C., Lo, D.C., 1999. Neurotrophins and synaptic plasticity neurotrophins and synaptic plasticity. *Annu. Rev. Neurosci.* 22, 295–318. doi:10.1007/7854

- McKenzie, S., Eichenbaum, H., 2011. Consolidation and reconsolidation: Two lives of memories? *Neuron* 71, 224–233. doi:<http://dx.doi.org/10.1016/j.neuron.2011.06.037>
- Meirelles, L. da S., Chagastelles, P.C., Nardi, N.B., 2006. Mesenchymal stem cells reside in virtually all post-natal organs and tissues. *J. Cell Sci.* 119, 2204–2213. doi:10.1242/jcs.02932
- Messaoudi, E., Bårdsen, K., Srebro, B., Bramham, C.R., 1998. Acute intrahippocampal infusion of BDNF induces lasting potentiation of synaptic transmission in the rat dentate gyrus. *J. Neurophysiol.* 79, 496–499. doi:10.1152/jn.1998.79.1.496
- Mignini, F., Nasuti, C., Artico, M., Giovannetti, F., Fumagalli, L., Iannetti, G., Pompili, E., Fabrizi, C., 2012. Effects of trimethyltin on hippocampal dopaminergic markers and cognitive behaviour. *Int. J. Immunopathol. Pharmacol.* 25, 1107–1119. doi:10.1177/039463201202500428
- Miki, T., Satriotomo, I., Li, H.-P., Matsumoto, Y., Gu, H., Yokoyama, T., Lee, K.-Y., Bedi, K.S., Takeuchi, Y., 2005. Application of the physical disector to the central nervous system: Estimation of the total number of neurons in subdivisions of the rat hippocampus. *Anat. Sci. Int.* 80, 153–162. doi:10.1111/j.1447-073x.2005.00121.x
- Ming, G., Song, H., 2011. Adult neurogenesis in the mammalian brain: Significant answers and significant questions. *Neuron* 70, 687–702. doi:10.1016/j.neuron.2011.05.001
- Mizuno, M., Yamada, K., Olariu, A., Nawa, H., Nabeshima, T., 2000. Involvement of brain-derived neurotrophic factor in spatial memory formation and maintenance in a radial arm maze test in rats. *J. Neurosci.* 20, 7116–21. doi:10.1523/JNEUROSCI.2018-00.2000 [pii]
- Mohyeldin, A., Garzón-Muvdi, T., Quiñones-Hinojosa, A., 2010. Oxygen in stem cell biology: A critical component of the stem cell niche. *Cell Stem Cell* 7, 150–161. doi:10.1016/j.stem.2010.07.007
- More, S., Kumar, H., Cho, D.-Y., Yun, Y.-S., Choi, D.-K., 2016. Toxin-induced experimental models of learning and memory impairment. *Int. J. Mol. Sci.* 17, 1–34. doi:10.3390/ijms17091447
- Morris, R.G.M., 1981. Spatial localization does not require the presence of local cues. *Learn. Motiv.* 12, 239–260. doi:[http://dx.doi.org/10.1016/0023-9690\(81\)90020-5](http://dx.doi.org/10.1016/0023-9690(81)90020-5)
- Nakatomi, H., Kuriu, T., Okabe, S., Yamamoto, S. ichi, Hatano, O., Kawahara, N., Tamura, A., Kirino, T., Nakafuku, M., 2002. Regeneration of hippocampal pyramidal neurons after ischemic brain injury by recruitment of endogenous neural progenitors. *Cell* 110, 429–441. doi:10.1016/S0092-8674(02)00862-0
- Nakazawa, K., Sun, L.D., Quirk, M.C., Rondi-Reig, L., Wilson, M.A., Tonegawa, S., 2003. Hippocampal CA3 NMDA receptors are crucial for memory acquisition of one-time experience. *Neuron* 38, 305–315. doi:[http://dx.doi.org/10.1016/S0896-6273\(03\)00165-X](http://dx.doi.org/10.1016/S0896-6273(03)00165-X)
- Neha, Sodhi, R.K., Jaggi, A.S., Singh, N., 2014. Animal models of dementia and cognitive dysfunction. *Life Sci.* 109, 73–86. doi:10.1016/j.lfs.2014.05.017



- Nekanti, U., Dastidar, S., Venugopal, P., Totey, S., Ta, M., 2010. Increased proliferation and analysis of differential gene expression in human Wharton's jelly-derived mesenchymal stromal cells under hypoxia. *Int. J. Biol. Sci.* 6, 499–512. doi:10.7150/ijbs.6.499
- Nie, Y., Han, B.M., Liu, X. Bin, Yang, J.J., Wang, F., Cong, X.F., Chen, X., 2011. Identification of micrnas involved in hypoxia- and serum deprivation-induced apoptosis in mesenchymal stem cells. *Int. J. Biol. Sci.* 7, 762–768. doi:10.7150/ijbs.7.762
- Nishina, A., Kimura, H., Kozawa, K., Sommen, G., Nakamura, T., Heimgartner, H., Koketsu, M., Furukawa, S., 2011. A superoxide anion-scavenger, 1,3-selenazolidin-4-one suppresses serum deprivation-induced apoptosis in PC12 cells by activating MAP kinase. *Toxicol. Appl. Pharmacol.* 257, 388–395. doi:10.1016/j.taap.2011.09.022
- Nyengaard, J.R., 1999. Stereologic methods and their application in kidney research. *J. Am. Soc. Nephrol.* 10, 1100–1123.
- Ogita, K., Nishiyama, N., Sugiyama, C., Higuchi, K., Yoneyama, M., Yoneda, Y., 2005. Regeneration of granule neurons after lesioning of hippocampal dentate gyrus: Evaluation using adult mice treated with trimethyltin chloride as a model. *J. Neurosci. Res.* 82, 609–621. doi:10.1002/jnr.20678
- Palomäki, S., Pietilä, M., Laitinen, S., Pesälä, J., Sormunen, R., Lehenkari, P., Koivunen, P., 2013. HIF-1 $\alpha$  is upregulated in human mesenchymal stem cells. *Stem Cells* 31, 1902–1909. doi:10.1002/stem.1435
- Papandreou, I., Cairns, R.A., Fontana, L., Lim, A.L., Denko, N.C., 2006. HIF-1 mediates adaptation to hypoxia by actively downregulating mitochondrial oxygen consumption. *Cell Metab.* 3, 187–197. doi:10.1016/j.cmet.2006.01.012
- Park, H.-J., Shim, H.S., Choi, W.K., Kim, K.S., Bae, H., Shim, I., 2011. Neuroprotective effect of *Lucium chinense* fruit on trimethyltin-induced learning and memory deficits in the rats. *Exp. Neurobiol.* 20, 137–143. doi:10.5607/en.2011.20.3.137
- Park, H.J., Lee, M.S., Shim, H.S., Lee, G.R., Chung, S.Y., Kang, Y.M., Lee, B.J., Seo, Y.B., Kim, K.S., Shim, I., 2016. Fermented *Saccharina japonica* (Phaeophyta) improves neuritogenic activity and TMT-induced cognitive deficits in rats. *Algae* 31, 73–84. doi:10.4490/algae.2016.31.11.10
- Park, I.H., Kim, K.H., Choi, H.K., Shim, J.S., Whang, S.Y., Hahn, S.J., Kwon, O.J., Oh, I.H., 2013. Constitutive stabilization of hypoxia-inducible factor alpha selectively promotes the self-renewal of mesenchymal progenitors and maintains mesenchymal stromal cells in an undifferentiated state. *Exp. Mol. Med.* 45, e44. doi:10.1038/emm.2013.87
- Park, S., Koh, S.E., Maeng, S., Lee, W.D., Lim, J., Lee, Y.J., 2011. Neural progenitors generated from the mesenchymal stem cells of first-trimester human placenta matured in the hypoxic-ischemic rat brain and mediated restoration of locomotor activity. *Placenta* 32, 269–276. doi:10.1016/j.placenta.2010.12.027
- Partadiredja, G., Bedi, K.S., 2010. Undernutrition during the gestation and suckling periods does not cause any loss of pyramidal neurons in the CA2-

- CA3 region of the rat hippocampus. *Nutr. Neurosci.* 13, 102–108. doi:10.1179/147683010X12611460764165
- Patterson, S.L., Abel, T., Deuel, T.A.S., Martin, K.C., Rose, J.C., Kandel, E.R., 1996. Recombinant BDNF rescues deficits in basal synaptic transmission and hippocampal LTP in BDNF knockout mice. *Neuron* 16, 1137–1145. doi:10.1016/S0896-6273(00)80140-3
- Paul, G., Anisimov, S. V., 2013. The secretome of mesenchymal stem cells: Potential implications for neuroregeneration. *Biochimie* 95, 2246–2256. doi:10.1016/j.biochi.2013.07.013
- Peterson, K.M., Aly, A., Lerman, A., Lerman, L.O., Rodriguez-Porcel, M., 2011. Improved survival of mesenchymal stromal cell after hypoxia preconditioning: Role of oxidative stress. *Life Sci.* 88, 65–73. doi:10.1016/j.lfs.2010.10.023
- Poldrack, R.A., Clark, J., Pare-Blagoev, E.J., Shohamy, D., 2001. Interactive memory systems in the human brain. *Nature* 414, 546–550. doi:http://dx.doi.org/10.1038/35107080
- Pulungan, Z.S.A., Sofro, Z.M., Partadiredja, G., 2016. Sodium fluoride does not affect the working memory and number of pyramidal cells in rat medial prefrontal cortex. *Anat. Sci. Int.* 93, 1–11. doi:10.1007/s12565-016-0384-4
- Puzzo, D., Lee, L., Palmeri, A., Calabrese, G., Arancio, O., 2014. Behavioral assays with mouse models of Alzheimer's disease: practical considerations and guidelines. *Biochem. Pharmacol.* 88, 450–467. doi:10.1016/j.bcp.2014.01.011
- Razban, V., Lotfi, A.S., Soleimani, M., Ahmadi, H., Massumi, M., Khajeh, S., Ghaedi, M., Arjmand, S., Najavand, S., Khoshdel, A., 2012. HIF-1 $\alpha$  overexpression induces angiogenesis in mesenchymal stem cells. *Biores. Open Access* 1, 174–183. doi:10.1089/biores.2012.9905
- Ribeiro, C.A., Fraga, J.S., Grãos, M., Neves, N.M., Reis, R.L., Gimble, J.M., Sousa, N., Salgado, A.J., 2012. The secretome of stem cells isolated from the adipose tissue and Wharton jelly acts differently on central nervous system derived cell populations. *Stem Cell Res. Ther.* 3, 1–7. doi:10.1186/scrt109
- Romanczyk, T.B., Weickert, C.S., Webster, M.J., Herman, M.M., Akil, M., Kleinman, J.E., 2002. Alterations in trkB mRNA in the human prefrontal cortex throughout the lifespan. *Eur. J. Neurosci.* 15, 269–280. doi:10.1046/j.0953-816x.2001.01858.x
- Rosova, I., Dao, M., Capoccia, B., Link, D., Nolte, J. a., 2008. Hypoxic preconditioning results in increased motility and improved therapeutic potential of human mesenchymal stem cells. *Stem Cells* 26, 2173–2182. doi:10.1634/stemcells.2007-1104.Hypoxic
- Sakhaie, M.H., Soleimani, M., Pirhajati, V., Soleimani Asl, S., Madjd, Z., Mehdizadeh, M., 2016. Coenzyme Q10 ameliorates trimethyltin chloride neurotoxicity in experimental model of injury in dentate gyrus of hippocampus: A histopathological and behavioral study. *Iran. Red Crescent Med. J.* 18, e30297. doi:10.5812/ircmj.30297
- Saki, N., Jalalifar, M.A., Soleimani, M., Hajizamani, S., Rahim, F., 2013. Adverse effect of high glucose concentration on stem cell therapy. *Int. J. Hematol.*

- Stem Cell Res. 7, 33–39.
- Salgado, A.J., Fraga, J.S., Mesquita, A.R., Neves, N.M., Reis, R.L., Sousa, N., 2010. Role of human umbilical cord mesenchymal progenitors conditioned media in neuronal/glial cell densities, viability, and proliferation. *Stem Cells Dev.* 19, 1067–1074. doi:10.1089/scd.2009.0279
- Schäbitz, W.-R., Steigleder, T., Cooper-Kuhn, C.M., Schwab, S., Sommer, C., Schneider, A., Kuhn, H.G., 2007. Intravenous brain-derived neurotrophic factor enhances poststroke sensorimotor recovery and stimulates neurogenesis. *Stroke* 38, 2165–2172. doi:10.1161/STROKEAHA.106.477331
- Semenza, G.L., 2004. Hydroxylation of HIF-1: Oxygen sensing at the molecular level. *Physiology* 19, 176–182. doi:10.1152/physiol.00001.2004
- Semenza, G.L., Jiang, B.-H., Leung, S.W., Passantino, R., Concorde, J.-P., Maire, P., Giallongo, A., 1996. Hypoxia response elements in the aldolase A, enolase 1, and lactate dehydrogenase A gene promoters contain essential binding sites for hypoxia-inducible factor 1. *J. Biol. Chem.* 271, 32529–32537. doi:10.1074/jbc.271.51.32529
- Silhol, M., Arancibia, S., Maurice, T., Tapia-Arancibia, L., 2007. Spatial memory training modifies the expression of brain-derived neurotrophic factor tyrosine kinase receptors in young and aged rats. *Neuroscience* 146, 962–973. doi:10.1016/j.neuroscience.2007.02.013
- Silhol, M., Bonnichon, V., Rage, F., Tapia-Arancibia, L., 2005. Age-related changes in brain-derived neurotrophic factor and tyrosine kinase receptor isoforms in the hippocampus and hypothalamus in male rats. *Neuroscience* 132, 613–624. doi:10.1016/j.neuroscience.2005.01.008
- Simon, H.-U., Haj-Yehia, A., Levi-Schaffer, F., 2000. Role of reactive oxygen species (ROS) in apoptosis induction. *Apoptosis* 5, 415–418. doi:10.1023/A:1009616228304
- Spence, K.W., Lippitt, R., 1946. An experimental test of the sign-gestalt theory of trial and error learning. *J. Exp. Psychol.* 36, 491–502. doi:10.1037/h0062419
- Sterio, D.C., 1984. The unbiased estimation of number and sizes of arbitrary particles using the disector. *J. Microsc.* 134, 127–136. doi:10.1111/j.1365-2818.1984.tb02501.x
- Stolzing, A., Coleman, N., Scutt, A., 2006. Glucose-induced replicative senescence in mesenchymal stem cells. *Rejuvenation Res.* 9, 31–35. doi:10.1089/rej.2006.9.31
- Teixeira, F.G., Carvalho, M.M., Neves-Carvalho, A., Panchalingam, K.M., Behie, L.A., Pinto, L., Sousa, N., Salgado, A.J., 2015. Secretome of mesenchymal progenitors from the umbilical cord acts as modulator of neural/glial proliferation and differentiation. *Stem Cell Rev. Reports* 11, 288–297. doi:10.1007/s12015-014-9576-2
- Tsai, M.-J., Tsai, S.-K., Hu, B.-R., Liou, D.-Y., Huang, S.-L., Huang, M.-C., Huang, W.-C., Cheng, H., Huang, S.-S., 2014. Recovery of neurological function of ischemic stroke by application of conditioned medium of bone marrow mesenchymal stem cells derived from normal and cerebral ischemia rats. *J. Biomed. Sci.* 21, 5. doi:10.1186/1423-0127-21-5

- Uccelli, A., 2013. Mesenchymal stem cells exert a remarkable regenerative effect requiring minimal CNS integration. Commentary on: "Mesenchymal stem cells protect CNS neurons against glutamate excitotoxicity by inhibiting glutamate receptor expression and function" by Voulg. *Exp. Neurol.* 247, 292–295. doi:10.1016/j.expneurol.2013.01.028
- Ullman, M.T., 2001. The declarative/procedural model of lexicon and grammar. *J. Psycholinguist. Res.* 30, 37–69. doi:<http://dx.doi.org/10.1023/A:1005204207369>
- Vizoso, F.J., Eiro, N., Cid, S., Schneider, J., Perez-Fernandez, R., 2017. Mesenchymal stem cell secretome: Toward cell-free therapeutic strategies in regenerative medicine. *Int. J. Mol. Sci.* doi:10.3390/ijms18091852
- Vorhees, C. V., Williams, M.T., 2014. Assessing spatial learning and memory in rodents. *ILAR J.* 55, 310–332. doi:10.1093/ilar/ilu013
- Vorhees, C. V., Williams, M.T., 2006. Morris water maze: procedures for assessing spatial and related forms of learning and memory. *Nat. Protoc.* 1, 848–858. doi:10.1038/nprot.2006.116
- Wang, G.L., Semenza, G.L., 1993. General involvement of hypoxia-inducible factor 1 in transcriptional response to hypoxia. *Genetics* 90, 4304–4308. doi:10.1073/pnas.90.9.4304
- Wang, H.-S., Hung, S.-C., Peng, S.-T., Huang, C.-C., Wei, H.-M., Guo, Y.-J., Fu, Y.-S., Lai, M.-C., Chen, C.-C., 2004. Mesenchymal stem cells in the Wharton's jelly of the human umbilical cord. *Stem Cells* 22, 1330–1337. doi:10.1634/stemcells.2004-0013
- Weickert, C.S., Hyde, T.M., Lipska, B.K., Herman, M.M., Weinberger, D.R., Kleinman, J.E., 2003. Reduced brain-derived neurotrophic factor in prefrontal cortex of patients with schizophrenia. *Mol. Psychiatry* 8, 592–610. doi:10.1038/sj.mp.4001308
- West, M.J., 2012. Estimating volume in biological structures. *Cold Spring Harb. Protoc.* 7, 1129–1139. doi:10.1101/pdb.top071787
- Wolf, A., Bauer, B., Abner, E.L., Ashkenazy-Frolinger, T., Hartz, A.M.S., 2016. A comprehensive behavioral test battery to assess learning and memory in 129S6/Tg2576 mice. *PLoS One* 11, 1–23. doi:10.1371/journal.pone.0147733
- Wurmser, A.E., Palmer, T.D., Gage, F.H., Holland, P., 2004. Cellular interactions in the stem cell niche. *Science* (80-. ). 304, 1253–1255. doi:10.1126/science.1099344
- Xu, X., Yi, F., Pan, H., Duan, S., Ding, Z., Yuan, G., Qu, J., Zhang, H., Liu, G., 2013. Progress and prospects in stem cell therapy. *Acta Pharmacol. Sin.* 34, 741–746. doi:10.1038/aps.2013.77
- Yamada, K., Nabeshima, T., 2003. Brain-derived neurotrophic factor/TrkB signaling in memory processes. *J. Pharmacol. Sci.* 91, 267–270. doi:10.1254/jphs.91.267
- Yuliani, S., Mustofa, Partadiredja, G., 2017. Turmeric (*Curcuma longa* L.) extract may prevent the deterioration of spatial memory and the deficit of estimated total number of hippocampal pyramidal cells of trimethyltin-exposed rats. *Drug Chem. Toxicol.* 0545, 1–10. doi:10.1080/01480545.2017.1293087
- Zhang, L., Tan, X., Dong, C., Zou, L., Zhao, H., Zhang, X., Tian, M., Jin, G.,

2012. In vitro differentiation of human umbilical cord mesenchymal stem cells (hUCMSCs), derived from Wharton's jelly, into choline acetyltransferase (ChAT)-positive cells. *Int. J. Dev. Neurosci.* 30, 471–477. doi:10.1016/j.ijdevneu.2012.05.006
- Zhang, W., Su, X., Gao, Y., Sun, B., Yu, Y., Wang, X., Zhang, F., 2009. Berberine protects mesenchymal stem cells against hypoxia-induced apoptosis in vitro. *Biol. Pharm. Bull.* 32, 1335–1342. doi:JST.JSTAGE/bpb/32.1335 [pii]
- Zhao, Q., Hu, J., Xiang, J., Gu, Y., Jin, P., Hua, F., Zhang, Z.Z., Liu, Y., Zan, K., Zhang, Z.Z., Zu, J., Yang, X., Shi, H., Zhu, J., Xu, Y., Cui, G., Ye, X., 2015. Intranasal administration of human umbilical cord mesenchymal stem cells-conditioned medium enhances vascular remodeling after stroke. *Brain Res.* 1624, 489–496. doi:10.1016/j.brainres.2015.08.003
- Zheng, X., Liang, Y., He, Q., Yao, R., Bao, W., Bao, L., Wang, Y., Wang, Z., 2014. Current models of mammalian target of rapamycin complex 1 (mTORC1) activation by growth factors and amino acids. *Int. J. Mol. Sci.* 15, 20753–20769. doi:10.3390/ijms151120753
- Zhu, S.F., Zhong, Z.N., Fu, X.F., Peng, D.X., Lu, G.H., Li, W.H., Xu, H.Y., Hu, H.B., He, J.M., Su, W.Y., He, Y.L., 2013. Comparison of cell proliferation, apoptosis, cellular morphology and ultrastructure between human umbilical cord and placenta-derived mesenchymal stem cells. *Neurosci. Lett.* 541, 77–82. doi:10.1016/j.neulet.2013.03.018
- Zhu, W., Chen, J., Cong, X., Hu, S., Chen, X., 2006. Hypoxia and serum deprivation-induced apoptosis in mesenchymal stem cells. *Stem Cells* 24, 416–425. doi:10.1634/stemcells.2005-0121