UNIVERSITAS
GADJAH MADA
PEMANFAATAN PRODUK SEKRETOM SEL PUNCA MESENKIMAL DARI TALI PUSAT MANUSIA YANG DIKULTUR DALAM KONDISI HIPOKSIA PADA MODEL DEGENERASI HIPPOCAMPUS
VERONIKA M. SIDHARTA, dr. Ginus Partadiredja, M.Sc., Ph.D.; dr. Rina Susilowati, Ph.D.; dr. E. Henny Herningtyas
Universitas Gadjah Mada, 2018 | Diunduh dari http://etd.repository.ugm.ac.id/

KEPUSTAKAAN

Allen, S.J., Watson, J.J., Shoemark, D.K., Barua, N.U., Patel, N.K., 2013. GDNF, NGF and BDNF as therapeutic options for neurodegeneration. Pharmacol. Ther. 138, 155–175. doi:10.1016/j.pharmthera.2013.01.004

Aranha, A.M.F., Zhang, Z., Neiva, K.G., Costa, C.A.S., Hebling, J., Nör, J.E., 2010. Hypoxia enhances the angiogenic potential of human dental pulp cells. J. Endod. 36, 1633–1637. doi:10.1016/j.joen.2010.05.013

Barzilai, A., Yamamoto, K.I., 2004. DNA damage responses to oxidative stress. DNA Repair (Amst). 3, 1109–1115. doi:10.1016/j.dnarep.2004.03.002

Bathina, S., Das, U.N., 2015. Brain-derived neurotrophic factor and its clinical Implications. Arch. Med. Sci. 11, 1164–1178. doi:10.5114/aoms.2015.56342

Baziyar, Y., Edalatmanesh, M.A., Hosseini, S.A., Zar, A., 2016. The effects of endurance training and gallic acid on BDNF and TNF-a in male rats with Alzheimer. Int. J. Appl. Exerc. Physiol. 5, 45–54.

Block, F., 1999. Global ischemia and behavioural deficits. Prog. Neurobiol. 58, 279–295. doi:10.1016/S0301-0082(98)00085-9

Bono, F., Lamarche, I., Bornia, J., Savi, P., Valle, G. Della, Herbert, J.-M., 1999. Nerve growth factor (NGF) exerts its pro-apoptotic e ¡ ect via the P75 NTR receptor in a cell cycle-dependent manner. FEBS Lett. 457, 93–97.

Bouet, V., Freret, T., Ankri, S., Bezault, M., Renolleau, S., Boulouard, M., Jacotot, E., Chauvier, D., Schumann-Bard, P., 2010. Predicting sensorimotor and memory deficits after neonatal ischemic stroke with reperfusion in the rat. Behav. Brain Res. 212, 56–63. doi:10.1016/j.bbr.2010.03.043

Boyce, R.W., Dorph-Petersen, K.A., Lyck, L., Gundersen, H.J.G., 2010. Design-based stereology: introduction to basic concepts and practical approaches for estimation of cell number. Toxicol. Pathol. 38, 1011–1025. doi:10.1177/0192623310385140

Bromley-Brits, K., Deng, Y., Song, W., 2011. Morris water maze test for learning and memory deficits in Alzheimer's disease model mice. J. Vis. Exp. 2–6. doi:10.3791/2920

Buravkova, L.B., Andreeva, E.R., Gogvadze, V., Zhivotovsky, B., 2014. Mesenchymal stem cells and hypoxia: Where are we? Mitochondrion 19, 105–112. doi:10.1016/j.mito.2014.07.005

Busletta, C., Novo, E., Valfrè Di Bonzo, L., Povero, D., Paternostro, C., Ievolella, M., Mareschi, K., Ferrero, I., Cannito, S., Compagnone, A., Bandino, A., Colombatto, S., Fagioli, F., Parola, M., 2011. Dissection of the biphasic nature of hypoxia-induced motogenic action in bone marrow-derived human mesenchymal stem cells. Stem Cells 29, 952–963. doi:10.1002/stem.642

Cantinieaux, D., Quertainmont, R., Blacher, S., Rossi, L., Wanet, T., Noël, A., Brook, G., Schoenen, J., Franzen, R., 2013. Conditioned medium from bone marrow-derived mesenchymal stem cells improves recovery after spinal cord injury in rats: An original strategy to avoid cell transplantation. PLoS One 8, e69515. doi:10.1371/journal.pone.0069515

Ceccariglia, S., D'Altocolle, A., Del Fa', A., Pizzolante, F., Caccia, E., Michetti, F., Gangitano, C., 2011. Cathepsin D plays a crucial role in the trimethyltin-induced hippocampal neurodegeneration process. Neuroscience 174, 160–

PEMANFAATAN PRODUK SEKRETOM SEL PUNCA MESENKIMAL DARI TALI PUSAT MANUSIA YANG DIKULTUR DALAM KONDISI HIPOKSIA PADA MODEL DEGENERASI HIPPOCAMPUS
VERONIKA M. SIDHARTA, dr. Ginus Partadiredja, M.Sc., Ph.D.; dr. Rina Susilowati, Ph.D.; dr. E. Henny Herningtyas
Universitas Gadjah Mada, 2018 | Diunduh dari http://etd.repository.ugm.ac.id/

170. doi:10.1016/j.neuroscience.2010.11.024

Chacko, S.M., Ahmed, S., Selvendiran, K., Kuppusamy, M.L., Khan, M., Kuppusamy, P., 2010. Hypoxic preconditioning induces the expression of prosurvival and proangiogenic markers in mesenchymal stem cells. Am. J. Physiol. Physiol. 299, C1562–C1570. doi:10.1152/ajpcell.00221.2010

Chang, C.-P., Chio, C.-C., Cheong, C.-U., Chao, C.-M., Cheng, B.-C., Lin, M.-T., 2013. Hypoxic preconditioning enhances the therapeutic potential of the secretome from cultured human mesenchymal stem cells in experimental traumatic brain injury. Clin. Sci. 124, 165–76. doi:10.1042/CS20120226

Chang, W., Song, B.W., Lim, S., Song, H., Chi, Y.S., Cha, M.J., Dong, H.A., Jung, Y.G., Lee, D.H., Ji, H.C., Choi, K.D., Lee, S.K., Chung, N., Lee, S.K., Jang, Y., Hwang, K.C., 2009. Mesenchymal stem cells pretreated with delivered Hph-1-Hsp70 protein are protected from hypoxia-mediated cell death and rescue heart functions from myocardial injury. Stem Cells 27, 2283–2292. doi:10.1002/stem.153

Charan, J., Kantharia, N.D., 2013. How to calculate sample size in animal studies? J. Pharmacol. Pharmacother. 4, 303–6. doi:10.4103/0976-500X.119726

Corvino, V., Geloso, M.C., Cavallo, V., Guadagni, E., Passalacqua, R., Florenzano, F., Giannetti, S., Molinari, M., Michetti, F., 2005. Enhanced neurogenesis during trimethyltin-induced neurodegeneration in the hippocampus of the adult rat. Brain Res. Bull. 65, 471–477. doi:10.1016/j.brainresbull.2005.02.031

Crisostomo, P.R., Wang, Y., Markel, T.A., Wang, M., Lahm, T., Meldrum, D.R., 2008. Human mesenchymal stem cells stimulated by TNF-a , LPS, or hypoxia produce growth factors by an NFkB- but not JNK-dependent mechanism. Am. J. Physiol. Cell Physiol. 294, C675–C682. doi:10.1152/ajpcell.00437.2007

Cunha, C., Brambilla, R., Thomas, K.L., 2010. A simple role for BDNF in learning and memory? Front. Mol. Neurosci. 3, 1–14. doi:10.3389/neuro.02.001.2010

da Silva Meirelles, L., Fontes, A.M., Covas, D.T., Caplan, A.I., Meirelles, L.D.S., Fontes, A.M., Covas, D.T., Caplan, A.I., Da, L., Meirelles, S., Fontes, A.M., Covas, D.T., Caplan, A.I., 2009. Mechanisms involved in the therapeutic properties of mesenchymal stem cells. Cytokine Growth Factor Rev. 20, 419–427. doi:10.1016/j.cytogfr.2009.10.002

Dalous, J., Larghero, J., Baud, O., 2012. Transplantation of umbilical cord-derived mesenchymal stem cells as a novel strategy to protect the central nervous system: technical aspects, preclinical studies, and clinical perspectives. Pediatr. Res. 71, 482–490. doi:10.1038/pr.2011.67

Dember, W.N., Fowler, H., 1958. Spontaneous alternation behavior. Psychol. Bull. 55, 412–428. doi:10.1007/BF02686644

Deschepper, M., Oudina, K., David, B., Myrtil, V., Collet, C., Bensidhoum, M., Logeart-Avramoglou, D., Petite, H., 2011. Survival and function of mesenchymal stem cells (MSCs) depend on glucose to overcome exposure to long-term, severe and continuous hypoxia. J. Cell. Mol. Med. 15, 1505–1514. doi:10.1111/j.1582-4934.2010.01138.x

UNIVERSITAS GADJAH MADA
PEMANFAATAN PRODUK SEKRETOM SEL PUNCA MESENKIMAL DARI TALI PUSAT MANUSIA YANG DIKULTUR DALAM KONDISI HIPOKSIA PADA MODEL DEGENERASI HIPPOCAMPUS
VERONIKA M. SIDHARTA, dr. Ginus Partadiredja, M.Sc., Ph.D.; dr. Rina Susilowati, Ph.D.; dr. E. Henny Herningtyas
Universitas Gadjah Mada, 2018 | Diunduh dari http://etd.repository.ugm.ac.id/

Dominici, M., Le Blanc, K., Mueller, I., Slaper-Cortenbach, I., Marini, F.C., Krause, D.S., Deans, R.J., Keating, A., Prockop, D.J., Horwitz, E.M., 2006. Minimal criteria for defining multipotent mesenchymal stromal cells. The International Society for Cellular Therapy position statement. Cytotherapy 8, 315–317. doi:10.1080/14653240600855905

Drago, D., Cossetti, C., Iraci, N., Gaude, E., Musco, G., Bachi, A., Pluchino, S., 2013. The stem cell secretome and its role in brain repair. Biochimie 95, 2271–2285. doi:10.1016/j.biochi.2013.06.020

Drela, K., Sarnowska, A., Siedlecka, P., Szablowska-Gadomska, I., Wielgos, M., Jurga, M., Lukomska, B., Domanska-Janik, K., 2014. Low oxygen atmosphere facilitates proliferation and maintains undifferentiated state of umbilical cord mesenchymal stem cells in an hypoxia inducible factor-dependent manner. Cytotherapy 16, 881–892. doi:10.1016/j.jcyt.2014.02.009

Dyachok, J., Earnest, S., Iturraran, E.N., Cobb, M.H., Ross, E.M., 2016. Amino acids regulate mTORC1 by an obligate two-step mechanism. J. Biol. Chem. 291, 22414–22426. doi:10.1074/jbc.M116.732511

Ebert, R., Ulmer, M., Zeck, S., Meissner-Weigl, J., Schneider, D., Stopper, H., Schupp, N., Kassem, M., Jakob, F., 2006. Selenium supplementation restores the antioxidative capacity and prevents cell damage in bone marrow stromal cells in vitro. Stem Cells 24, 1226–1235. doi:10.1634/stemcells.2005-0117

Efimenko, A., Starostina, E., Kalinina, N., Stolzing, A., 2011. Angiogenic properties of aged adipose derived mesenchymal stem cells after hypoxic conditioning.pdf 1–13.

Eichenbaum, H., Stewart, C., Morris, R.G.M., 1990. Hippocampal representation in place learning. J. Neurosci. 10, 3531–3542. doi:10.1523/jneurosci.1068-06.2006

Ernfors, P., Ibáñez, C.F., Ebendal, T., Olson, L., Persson, H., 1990. Molecular cloning and neurotrophic activities of a protein with structural similarities to nerve growth factor: developmental and topographical expression in the brain. Proc. Natl. Acad. Sci. U. S. A. 87, 5454–5458. doi:10.1073/pnas.87.14.5454

Fan, C.-G., Zhang, Q., Zhou, J., 2011. Therapeutic potentials of mesenchymal stem cells derived from human umbilical cord. Stem Cell Rev. Reports 7, 195–207. doi:10.1007/s12015-010-9168-8

Forsythe, J.O.A., Jiang, B., Iyer, N. V., Agani, F., Leung, S.W., 1996. Activation of vascular endothelial growth factor gene transcription by hypoxia-inducible factor 1. Mol. Cell. Biol. 16, 4604–4612. doi:10.1128/MCB.16.9.4604

Friedenstein, A.J., Piatetzky-Shapiro, I.I., Petrakova, K.V., 1966. Osteogenesis in transplants of bone marrow cells. J. Embryol. Exp. Morphol. 16, 381–390.

Gage, F.H., 2000. Mammalian neural stem cells. Science (80-.). 287, 1433–1438. doi:10.1126/science.287.5457.1433

Galati, G., Pelle, G., Berthoz, A., Committeri, G., 2010. Multiple reference frames used by the human brain for spatial perception and memory. Exp. Brain Res. 206, 109–120. doi:10.1007/s00221-010-2168-8

Gasparova, Z., Janega, P., Stara, V., Ujhazy, E., Ujhazy, A.E., 2012. Early and late stage of neurodegeneration induced by trimethyltin in hippocampus and

PEMANFAATAN PRODUK SEKRETOM SEL PUNCA MESENKIMAL DARI TALI PUSAT MANUSIA YANG DIKULTUR DALAM KONDISI HIPOKSIA PADA MODEL DEGENERASI HIPPOCAMPUS
VERONIKA M. SIDHARTA, dr. Ginus Partadiredja, M.Sc., Ph.D.; dr. Rina Susilowati, Ph.D.; dr. E. Henny Herningtyas
Universitas Gadjah Mada, 2018 | Diunduh dari http://etd.repository.ugm.ac.id/

cortex of male Wistar rats. Neuroendocrinol. Lett. 33, 689–696.

Geloso, M.C., Corvino, V., Michetti, F., 2011. Trimethyltin-induced hippocampal degeneration as a tool to investigate neurodegenerative processes. Neurochem. Int. 58, 729–738. doi:10.1016/j.neuint.2011.03.009

Ghosh, A., Carnahan, J., Greenberg, M.E., 1994. Requirement for BDNF in activity-dependent survival of cortical neurons. Science (80-.). 263, 1618–1623. doi:10.1126/science.7907431

Gögel, S., Gubernator, M., Minger, S.L., 2011. Progress and prospects: Stem cells and neurological diseases. Gene Ther. 18, 1–6. doi:10.1038/gt.2010.130

Gómez-Palacio-Schjetnan, A., Escobar, M.L., 2013. Neurotrophins and synaptic plasticity. Curr. Top. Behav. Neurosci. 15, 117–36. doi:10.1007/7854

Govindasamy, V., Ronald, V.S., Totey, S., Binti Din, S., Mustafa, W.M.B.W., Totey, S., Zakaria, Z., Bhonde, R.R., 2010. Micromanipulation of culture niche permits long-term expansion of dental pulp stem cells-an economic and commercial angle. Vitr. Cell. Dev. Biol. - Anim. 46, 764–773. doi:10.1007/s11626-010-9332-0

Grayson, W.L., Zhao, F., Bunnell, B., Ma, T., 2007. Hypoxia enhances proliferation and tissue formation of human mesenchymal stem cells. Biochem. Biophys. Res. Commun. 358, 948–953. doi:10.1016/j.bbrc.2007.05.054

Gundersen, H.J.G., 1977. Notes on the estimation of numerical density of arbitrary profiles: the edge effect. J. Microsc. 111, 219–223.

Guzy, R.D., Hoyos, B., Robin, E., Chen, H., Liu, L., Mansfield, K.D., Simon, M.C., Hammerling, U., Schumacker, P.T., 2005. Mitochondrial complex III is required for hypoxia-induced ROS production and cellular oxygen sensing. Cell Metab. 1, 401–408. doi:10.1016/j.cmet.2005.05.001

Guzy, R.D., Schumacker, P.T., 2006. Oxygen sensing by mitochondria at complex III: The paradox of increased reactive oxygen species during hypoxia. Exp. Physiol. 91, 807–819. doi:10.1113/expphysiol.2006.033506

Ha, J.S., Jin, D.E., Park, S.K., Park, C.H., Seung, T.W., Bae, D.-W., Kim, D.-O., Heo, H.J., 2015. Antiamnesic effect of Actinidia arguta extract intake in a mouse model of TMT-induced learning and memory dysfunction. Evidence-Based Complement. Altern. Med. 2015, 1–13.

Hagmann, S., Moradi, B., Frank, S., Dreher, T., Kämmerer, P.W., Richter, W., Gotterbarm, T., 2013. Different culture media affect growth characteristics, surface marker distribution and chondrogenic differentiation of human bone marrow-derived mesenchymal stromal cells. BMC Musculoskelet. Disord. 14, 1–7. doi:10.1186/1471-2474-14-223

Hayashi, M., Mitsunaga, F., Ohira, K., Shimizu, K., Yamashita, A., 1999. Development of full-length Trk B-immunoreactive structures in the hippocampal formation of the macaque monkey. Anat. Embryol. 199, 529–537.

Hofer, M., Pagliusi, S.R., Hohn, A., Leibrock, J., Barde, Y.-A., 1990. Regional distribution of brain-derived neurotrophic factor mRNA in the adult mouse brain-derived neurotrophic factor/hippocampus/ in situ hybridization/nerve growth factor/trophic interactions. EMBO J. 9, 2459–2464.

UNIVERSITAS GADJAH MADA
PEMANFAATAN PRODUK SEKRETOM SEL PUNCA MESENKIMAL DARI TALI PUSAT MANUSIA YANG DIKULTUR DALAM KONDISI HIPOKSIA PADA MODEL DEGENERASI HIPPOCAMPUS
VERONIKA M. SIDHARTA, dr. Ginus Partadiredja, M.Sc., Ph.D.; dr. Rina Susilowati, Ph.D.; dr. E. Henny Herningtyas
Universitas Gadjah Mada, 2018 | Diunduh dari http://etd.repository.ugm.ac.id/

Holmes, D.I.R., Zachary, I., 2005. The vascular endothelial growth factor (VEGF) family: angiogenic factors in health and disease. Genome Biol. 6, 1–10. doi:10.1186/gb-2005-6-2-209

Hsieh, J.-Y., Wang, H.-W., Chang, S.-J., Liao, K.-H., Lee, I.-H., Lin, W.-S., Wu, C.-H., Lin, W.-Y., Cheng, S.-M., 2013. Mesenchymal stem cells from human umbilical cord express preferentially secreted factors related to neuroprotection, neurogenesis, and angiogenesis. PLoS One 8, 1–11. doi:10.1371/journal.pone.0072604

Hsieh, P.-S., Bochinski, D.J., Lin, G.T., Nunes, L., Lin, C.S., Lue, T.F., 2003. The effect of vascular endothelial growth factor and brain-derived neurotrophic factor on cavernosal nerve regeneration in a nerve-crush rat model. BJU Int. 92, 470–475. doi:10.1046/j.1464-410X.2003.04373.x

Jin, K., Zhu, Y., Sun, Y., Mao, X.O., Xie, L., Greenberg, D.A., 2002. Vascular endothelial growth factor (VEGF) stimulates neurogenesis in vitro and in vivo. Proc Natl Acad Sci U S A 99, 11946–11950. doi:10.1073/pnas.182296499

Joerger-Messerli, M.S., Marx, C., Oppliger, B., Mueller, M., Surbek, D. V., Schoeberlein, A., 2016. Mesenchymal stem cells from Wharton's jelly and amniotic fluid. Best Pract. Res. Clin. Obstet. Gynaecol. 31, 30–44. doi:10.1016/j.bpobgyn.2015.07.006

Johnson, E., Adamo-Villani, N., 2010. A study of the effects of immersion on short-term spatial memory. World Acad. Sci. Eng. Technol. 71, 582–587.

Jung, E.-Y., Lee, M.-S., Ahn, C.J., Cho, S.-H., Bae, H., Shim, I., 2013. The neuroprotective effect of gugijihwang-tang on trimethyltin-induced memory dysfunction in the rat. Evidence-Based Complement. Altern. Med. 2013, 1–6. doi:10.1155/2013/542081

Kang, H., Schuman, E.M., 1995. Long-lasting neurotrophin-induced enhancement of synaptic transmission in the adult hippocampus. Science (80-.). 267, 1658–1662.

Kang, J.Y., Park, S.K., Guo, T.J., Ha, J.S., Lee, D.S., Kim, J.M., Lee, U., Kim, D.O., Heo, H.J., 2016. Reversal of trimethyltin-induced learning and memory deficits by 3, 5-dicaffeoylquinic acid. Oxid. Med. Cell. Longev. 2016, 1–13.

Ke, X., Zhang, J., 2013. Changes in HIF-1 α , VEGF, NGF and BDNF levels in cerebrospinal fluid and their relationship with cognitive impairment in patients with cerebral infarction. J. Huazhong Univ. Sci. Technol. [Medical Sci. 33, 433–437.

Kesner, R.P., Hopkins, R.O., 2006. Mnemonic functions of the hippocampus: A comparison between animals and humans. Biol. Psychol., Translational Research: Parallels of Human and Animal Research in Biological Psychology 73, 3–18. doi:10.1016/j.biopsycho.2006.01.004

Kessels, R.P.C., Feijen, J., Postma, A., 2005. Implicit and explicit memory for spatial information in Alzheimer's disease. Dement. Geriatr. Cogn. Disord. 20, 184–191. doi:10.1159/000087233

Kim, J.W., Tchernyshyov, I., Semenza, G.L., Dang, C. V., 2006. HIF-1-mediated expression of pyruvate dehydrogenase kinase: A metabolic switch required

UNIVERSITAS GADJAH MADA
PEMANFAATAN PRODUK SEKRETOM SEL PUNCA MESENKIMAL DARI TALI PUSAT MANUSIA YANG DIKULTUR DALAM KONDISI HIPOKSIA PADA MODEL DEGENERASI HIPPOCAMPUS
VERONIKA M. SIDHARTA, dr. Ginus Partadiredja, M.Sc., Ph.D.; dr. Rina Susilowati, Ph.D.; dr. E. Henny Herningtyas
Universitas Gadjah Mada, 2018 | Diunduh dari http://etd.repository.ugm.ac.id/

for cellular adaptation to hypoxia. Cell Metab. 3, 177–185. doi:10.1016/j.cmet.2006.02.002

Kim, S.G., Buel, G.R., Blenis, J., 2013. Nutrient regulation of the mTOR complex 1 signaling pathway. Mol. Cells 35, 463–473. doi:10.1007/s10059-013-0138-2

Koda, T., Kuroda, Y., Imai, H., 2008. Protective effect of rutin against spatial memory impairment induced by trimethyltin in rats. Nutr. Res. 28, 629–634. doi:10.1016/j.nutres.2008.06.004

Korte, M., Carroll, P., Wolf, E., Brem, G., Thoenen, H., Bonhoeffer, T., 1995. Hippocampal long-term potentiation is impaired in mice lacking brain-derived neurotrophic factor. Proc. Natl. Acad. Sci. U. S. A. 92, 8856–8860. doi:10.1073/pnas.92.19.8856

Korte, M., Staiger, V., Griesbeck, O., Thoenen, H., Bonhoeffer, T., 1996. The involvement of brain-derived neurotrophic factor in hippocampal long-term potentiation revealed by gene targeting experiments. J. Physiol. Paris 90, 157–164. doi:10.1016/S0928-4257(97)81415-5

Kovalchuk, Y., Hanse, E., Kafitz, K.W., Konnerth, A., 2002. Postsynaptic induction of BDNF-mediated long-term potentiation. Science (80-.). 295, 1729–1734. doi:10.1126/science.1067766

Lavoie, J.R., Rosu-Myles, M., 2013. Uncovering the secretes of mesenchymal stem cells. Biochimie 95, 2212–2221. doi:10.1016/j.biochi.2013.06.017

Lavrentieva, A., Majore, I., Kasper, C., Hass, R., 2010. Effects of hypoxic culture conditions on umbilical cord-derived human mesenchymal stem cells. Cell Commun. Signal. 8, 18. doi:10.1186/1478-811X-8-18

Lee, B., Sur, B., Cho, S.-G., Yeom, M., Shim, I., Lee, H., Hahm, D.-H., 2016. Wogonin attenuates hippocampal neuronal loss and cognitive dysfunction in trimethyltin-intoxicated rats. Biomol. Ther. 24, 328–337. doi:10.4062/biomolther.2015.152

Lee, C.Y., Kang, J.Y., Lim, S., Ham, O., Chang, W., Jang, D.-H., 2016. Hypoxic conditioned medium from mesenchymal stem cells promotes lymphangiogenesis by regulation of mitochondrial-related proteins. Stem Cell Res. Ther. 7, 38. doi:10.1186/s13287-016-0296-1

Lee, E., Son, H., 2009. Adult hippocampal neurogenesis and related neurotrophic factors. BMB Rep. 42, 239–244. doi:10.5483/BMBRep.2009.42.5.239

Lee, H.J., Ryu, J.M., Jung, Y.H., Oh, S.Y., Lee, S.J., Han, H.J., 2015. Novel pathway for hypoxia-induced proliferation and migration in human mesenchymal stem cells: Involvement of HIF-1α, FASN, and mTORC1. Stem Cells 33, 2182–2195. doi:10.1002/stem.2020

Lee, S.C., Kim, K.H., Kim, O.H., Lee, S.K., Hong, H.E., Won, S.S., Jeon, S.J., Choi, B.J., Jeong, W., Kim, S.J., 2017. Determination of optimized oxygen partial pressure to maximize the liver regenerative potential of the secretome obtained from adipose-derived stem cells. Stem Cell Res. Ther. 8, 1–12. doi:10.1186/s13287-017-0635-x

Lee, S.H., Lee, Y.J., Song, C.H., Ahn, Y.K., Han, H.J., 2010. Role of FAK phosphorylation in hypoxia-induced hMSCS migration: involvement of VEGF as well as MAPKS and eNOS pathways. Am. J. Physiol. Cell Physiol.

UNIVERSITAS
GADJAH MADA
PEMANFAATAN PRODUK SEKRETOM SEL PUNCA MESENKIMAL DARI TALI PUSAT MANUSIA YANG DIKULTUR DALAM KONDISI HIPOKSIA PADA MODEL DEGENERASI HIPPOCAMPUS
VERONIKA M. SIDHARTA, dr. Ginus Partadiredja, M.Sc., Ph.D.; dr. Rina Susilowati, Ph.D.; dr. E. Henny Herningtyas
Universitas Gadjah Mada, 2018 | Diunduh dari http://etd.repository.ugm.ac.id/

298, C847–C856. doi:10.1152/ajpcell.00418.2009

Levine, E.S., Dreyfus, C.F., Black, I.B., Plummer, M.R., 1995. Brain-derived neurotrophic factor rapidly enhances synaptic transmission in hippocampal neurons via postsynaptic tyrosine kinase receptors. Proc. Natl. Acad. Sci. 92, 8074–8077. doi:10.1073/pnas.92.17.8074

Li, X., Bai, J., Ji, X., Li, R., Xuan, Y., Wang, Y., 2014. Comprehensive characterization of four different populations of human mesenchymal stem cells as regards their immune properties, proliferation and differentiation. Int. J. Mol. Med. 34, 695–704. doi:10.3892/ijmm.2014.1821

Lin, C., McGough, R., Aswad, B., Block, J., Terek, R., 2004. Hypoxia induces HIF-1-α and VEGF expression in chondrosarcoma cells and chondrocytes. J. Orthop. Res. 22, 1175–1181. doi:10.1016/j.orthres.2004.03.002

Lindholm, D., Carroll, P., Tzimagiorgis, G., Thoenen, H., 1996. Autocrine-paracrine regulation of hippocampal neuron survival by IGF-1 and the neurotrophins BDNF, NT-3 and NT-4. Eur. J. Neurosci. 8, 1452–1460. doi:10.1111/j.1460-9568.1996.tb01607.x

Lindvall, O., Kokaia, Z., 2010. Stem cells in human neurodegenerative disorders — time for clinical translation? J. Clin. Invest. 120, 29–40. doi:10.1172/JCI40543.patients

Liu, L., Gao, J., Yuan, Y., Chang, Q., Liao, Y., Lu, F., 2013. Hypoxia preconditioned human adipose derived mesenchymal stem cells enhance angiogenic potential via secretion of increased VEGF and bFGF. Cell Biol. Int. 37, 551–560. doi:10.1002/cbin.10097

Lowenstein, D.H., Arsenault, L., 1996. The effects of growth factors on the survival and differentiation of cultured dentate gyrus neurons. J. Neurosci. 16, 1759–1769.

Lührs, M.-L., Dammhahn, M., Kappeler, P.M., Fichtel, C., 2009. Spatial memory in the grey mouse lemur (Microcebus murinus). Anim. Cogn. 12, 599–609. doi:http://dx.doi.org/10.1007/s10071-009-0219-y

Ma, Y.L., Wang, H.L., Wu, H.C., Wei, C.L., Lee, E.H., 1998. Brain-derived neurotrophic factor antisense oligonucleotide impairs memory retention and inhibits long-term potentiation in rats. Neuroscience 82, 957–967. doi:10.1016/S0306-4522(97)00325-4

Majumdar, D., Bhonde, R., Datta, I., 2013. Influence of ischemic microenvironment on human Wharton's jelly mesenchymal stromal cells. Placenta 34, 642–649. doi:10.1016/j.placenta.2013.04.021

Maltman, D.J., Hardy, S. a., Przyborski, S. a., 2011. Role of mesenchymal stem cells in neurogenesis and nervous system repair. Neurochem. Int. 59, 347–356. doi:10.1016/j.neuint.2011.06.008

Martens, L.K., Kirschner, K.M., Warnecke, C., Scholz, H., 2007. Hypoxia-inducible factor-1 (HIF-1) is a transcriptional activator of the TrkB neurotrophin receptor gene. J. Biol. Chem. 282, 14379–14388. doi:10.1074/jbc.M609857200

McAllister, A.K., Katz, L.C., Lo, D.C., 1999. Neurotrophins and synaptic plasticity neurotrophins and synaptic plasticity. Annu. Rev. Neurosci. 22, 295–318. doi:10.1007/7854

PEMANFAATAN PRODUK SEKRETOM SEL PUNCA MESENKIMAL DARI TALI PUSAT MANUSIA YANG DIKULTUR DALAM KONDISI HIPOKSIA PADA MODEL DEGENERASI HIPPOCAMPUS
VERONIKA M. SIDHARTA, dr. Ginus Partadiredja, M.Sc., Ph.D.; dr. Rina Susilowati, Ph.D.; dr. E. Henny Herningtyas
Universitas Gadjah Mada, 2018 | Diunduh dari http://etd.repository.ugm.ac.id/

McKenzie, S., Eichenbaum, H., 2011. Consolidation and reconsolidation: Two lives of memories? Neuron 71, 224–233. doi:http://dx.doi.org/10.1016/j.neuron.2011.06.037

Meirelles, L. da S., Chagastelles, P.C., Nardi, N.B., 2006. Mesenchymal stem cells reside in virtually all post-natal organs and tissues. J. Cell Sci. 119, 2204–2213. doi:10.1242/jcs.02932

Messaoudi, E., Bårdsen, K., Srebro, B., Bramham, C.R., 1998. Acute intrahippocampal infusion of BDNF induces lasting potentiation of synaptic transmission in the rat dentate gyrus. J. Neurophysiol. 79, 496–499. doi:10.1152/jn.1998.79.1.496

Mignini, F., Nasuti, C., Artico, M., Giovannetti, F., Fumagalli, L., Iannetti, G., Pompili, E., Fabrizi, C., 2012. Effects of trimethyltin on hippocampal dopaminergic markers and cognitive behaviour. Int. J. Immunopathol. Pharmacol. 25, 1107–1119. doi:10.1177/039463201202500428

Miki, T., Satriotomo, I., Li, H.-P., Matsumoto, Y., Gu, H., Yokoyama, T., Lee, K.-Y., Bedi, K.S., Takeuchi, Y., 2005. Application of the physical disector to the central nervous system: Estimation of the total number of neurons in subdivisions of the rat hippocampus. Anat. Sci. Int. 80, 153–162. doi:10.1111/j.1447-073x.2005.00121.x

Ming, G., Song, H., 2011. Adult neurogenesis in the mammalian brain: Significant answers and significant questions. Neuron 70, 687–702. doi:10.1016/j.neuron.2011.05.001

Mizuno, M., Yamada, K., Olariu, A., Nawa, H., Nabeshima, T., 2000. Involvement of brain-derived neurotrophic factor in spatial memory formation and maintenance in a radial arm maze test in rats. J. Neurosci. 20, 7116–21. doi:20/18/7116 [pii]

Mohyeldin, A., Garzón-Muvdi, T., Quiñones-Hinojosa, A., 2010. Oxygen in stem cell biology: A critical component of the stem cell niche. Cell Stem Cell 7, 150–161. doi:10.1016/j.stem.2010.07.007

More, S., Kumar, H., Cho, D.-Y., Yun, Y.-S., Choi, D.-K., 2016. Toxin-induced experimental models of learning and memory impairment. Int. J. Mol. Sci. 17, 1–34. doi:10.3390/ijms17091447

Morris, R.G.M., 1981. Spatial localization does not require the presence of local cues. Learn. Motiv. 12, 239–260. doi:http://dx.doi.org/10.1016/0023-9690(81)90020-5

Nakatomi, H., Kuriu, T., Okabe, S., Yamamoto, S. ichi, Hatano, O., Kawahara, N., Tamura, A., Kirino, T., Nakafuku, M., 2002. Regeneration of hippocampal pyramidal neurons after ischemic brain injury by recruitment of endogenous neural progenitors. Cell 110, 429–441. doi:10.1016/S0092-8674(02)00862-0

Nakazawa, K., Sun, L.D., Quirk, M.C., Rondi-Reig, L., Wilson, M.A., Tonegawa, S., 2003. Hippocampal CA3 NMDA receptors are crucial for memory acquisition of one-time experience. Neuron 38, 305–315. doi:http://dx.doi.org/10.1016/S0896-6273(03)00165-X

Neha, Sodhi, R.K., Jaggi, A.S., Singh, N., 2014. Animal models of dementia and cognitive dysfunction. Life Sci. 109, 73–86. doi:10.1016/j.lfs.2014.05.017

UNIVERSITAS GADJAH MADA
PEMANFAATAN PRODUK SEKRETOM SEL PUNCA MESENKIMAL DARI TALI PUSAT MANUSIA YANG DIKULTUR DALAM KONDISI HIPOKSIA PADA MODEL DEGENERASI HIPPOCAMPUS
VERONIKA M. SIDHARTA, dr. Ginus Partadiredja, M.Sc., Ph.D.; dr. Rina Susilowati, Ph.D.; dr. E. Henny Herningtyas
Universitas Gadjah Mada, 2018 | Diunduh dari http://etd.repository.ugm.ac.id/

Nekanti, U., Dastidar, S., Venugopal, P., Totey, S., Ta, M., 2010. Increased proliferation and analysis of differential gene expression in human Wharton's jelly-derived mesenchymal stromal cells under hypoxia. Int. J. Biol. Sci. 6, 499–512. doi:10.7150/ijbs.6.499

Nie, Y., Han, B.M., Liu, X. Bin, Yang, J.J., Wang, F., Cong, X.F., Chen, X., 2011. Identification of micrornas involved in hypoxia- and serum deprivation-induced apoptosis in mesenchymal stem cells. Int. J. Biol. Sci. 7, 762–768. doi:10.7150/ijbs.7.762

Nishina, A., Kimura, H., Kozawa, K., Sommen, G., Nakamura, T., Heimgartner, H., Koketsu, M., Furukawa, S., 2011. A superoxide anion-scavenger, 1,3-selenazolidin-4-one suppresses serum deprivation-induced apoptosis in PC12 cells by activating MAP kinase. Toxicol. Appl. Pharmacol. 257, 388–395. doi:10.1016/j.taap.2011.09.022

Nyengaard, J.R., 1999. Stereologic methods and their application in kidney research. J. Am. Soc. Nephrol. 10, 1100–1123.

Ogita, K., Nishiyama, N., Sugiyama, C., Higuchi, K., Yoneyama, M., Yoneda, Y., 2005. Regeneration of granule neurons after lesioning of hippocampal dentate gyrus: Evaluation using adult mice treated with trimethyltin chloride as a model. J. Neurosci. Res. 82, 609–621. doi:10.1002/jnr.20678

Palomäki, S., Pietilä, M., Laitinen, S., Pesälä, J., Sormunen, R., Lehenkari, P., Koivunen, P., 2013. HIF-1α is upregulated in human mesenchymal stem cells. Stem Cells 31, 1902–1909. doi:10.1002/stem.1435

Papandreou, I., Cairns, R.A., Fontana, L., Lim, A.L., Denko, N.C., 2006. HIF-1 mediates adaptation to hypoxia by actively downregulating mitochondrial oxygen consumption. Cell Metab. 3, 187–197. doi:10.1016/j.cmet.2006.01.012

Park, H.-J., Shim, H.S., Choi, W.K., Kim, K.S., Bae, H., Shim, I., 2011. Neuroprotective effect of Lucium chinense fruit on trimethyltin-induced learning and memory deficits in the rats. Exp. Neurobiol. 20, 137–143. doi:10.5607/en.2011.20.3.137

Park, H.J., Lee, M.S., Shim, H.S., Lee, G.R., Chung, S.Y., Kang, Y.M., Lee, B.J., Seo, Y.B., Kim, K.S., Shim, I., 2016. Fermented Saccharina japonica (Phaeophyta) improves neuritogenic activity and TMT-induced cognitive deficits in rats. Algae 31, 73–84. doi:10.4490/algae.2016.31.11.10

Park, I.H., Kim, K.H., Choi, H.K., Shim, J.S., Whang, S.Y., Hahn, S.J., Kwon, O.J., Oh, I.H., 2013. Constitutive stabilization of hypoxia-inducible factor alpha selectively promotes the self-renewal of mesenchymal progenitors and maintains mesenchymal stromal cells in an undifferentiated state. Exp. Mol. Med. 45, e44. doi:10.1038/emm.2013.87

Park, S., Koh, S.E., Maeng, S., Lee, W.D., Lim, J., Lee, Y.J., 2011. Neural progenitors generated from the mesenchymal stem cells of first-trimester human placenta matured in the hypoxic-ischemic rat brain and mediated restoration of locomotor activity. Placenta 32, 269–276. doi:10.1016/j.placenta.2010.12.027

Partadiredja, G., Bedi, K.S., 2010. Undernutrition during the gestation and suckling periods does not cause any loss of pyramidal neurons in the CA2-

UNIVERSITAS GADJAH MADA
PEMANFAATAN PRODUK SEKRETOM SEL PUNCA MESENKIMAL DARI TALI PUSAT MANUSIA YANG DIKULTUR DALAM KONDISI HIPOKSIA PADA MODEL DEGENERASI HIPPOCAMPUS
VERONIKA M. SIDHARTA, dr. Ginus Partadiredja, M.Sc., Ph.D.; dr. Rina Susilowati, Ph.D.; dr. E. Henny Herningtyas
Universitas Gadjah Mada, 2018 | Diunduh dari http://etd.repository.ugm.ac.id/

CA3 region of the rat hippocampus. Nutr. Neurosci. 13, 102–108. doi:10.1179/147683010X12611460764165

Patterson, S.L., Abel, T., Deuel, T.A.S., Martin, K.C., Rose, J.C., Kandel, E.R., 1996. Recombinant BDNF rescues deficits in basal synaptic transmission and hippocampal LTP in BDNF knockout mice. Neuron 16, 1137–1145. doi:10.1016/S0896-6273(00)80140-3

Paul, G., Anisimov, S. V., 2013. The secretome of mesenchymal stem cells: Potential implications for neuroregeneration. Biochimie 95, 2246–2256. doi:10.1016/j.biochi.2013.07.013

Peterson, K.M., Aly, A., Lerman, A., Lerman, L.O., Rodriguez-Porcel, M., 2011. Improved survival of mesenchymal stromal cell after hypoxia preconditioning: Role of oxidative stress. Life Sci. 88, 65–73. doi:10.1016/j.lfs.2010.10.023

Poldrack, R.A., Clark, J., Pare-Blagoev, E.J., Shohamy, D., 2001. Interactive memory systems in the human brain. Nature 414, 546–550. doi:http://dx.doi.org/10.1038/35107080

Pulungan, Z.S.A., Sofro, Z.M., Partadiredja, G., 2016. Sodium fluoride does not affect the working memory and number of pyramidal cells in rat medial prefrontal cortex. Anat. Sci. Int. 93, 1–11. doi:10.1007/s12565-016-0384-4

Puzzo, D., Lee, L., Palmeri, A., Calabrese, G., Arancio, O., 2014. Behavioral assays with mouse models of Alzheimer's disease: practical considerations and guidelines. Biochem. Pharmacol. 88, 450–467. doi:10.1016/j.bcp.2014.01.011

Razban, V., Lotfi, A.S., Soleimani, M., Ahmadi, H., Massumi, M., Khajeh, S., Ghaedi, M., Arjmand, S., Najavand, S., Khoshdel, A., 2012. HIF-1α overexpression induces angiogenesis in mesenchymal stem cells. Biores. Open Access 1, 174–183. doi:10.1089/biores.2012.9905

Ribeiro, C.A., Fraga, J.S., Grãos, M., Neves, N.M., Reis, R.L., Gimble, J.M., Sousa, N., Salgado, A.J., 2012. The secretome of stem cells isolated from the adipose tissue and Wharton jelly acts differently on central nervous system derived cell populations. Stem Cell Res. Ther. 3, 1–7. doi:10.1186/scrt109

Romanczyk, T.B., Weickert, C.S., Webster, M.J., Herman, M.M., Akil, M., Kleinman, J.E., 2002. Alterations in trkB mRNA in the human prefrontal cortex throughout the lifespan. Eur. J. Neurosci. 15, 269–280. doi:10.1046/j.0953-816x.2001.01858.x

Rosova, I., Dao, M., Capoccia, B., Link, D., Nolta, J. a., 2008. Hypoxic preconditioning results in increased motility and improved therapeutic potential of human mesenchymal stem cells. Stem Cells 26, 2173–2182. doi:10.1634/stemcells.2007-1104.Hypoxic

Sakhaie, M.H., Soleimani, M., Pirhajati, V., Soleimani Asl, S., Madjd, Z., Mehdizadeh, M., 2016. Coenzyme Q10 ameliorates trimethyltin chloride neurotoxicity in experimental model of injury in dentate gyrus of hippocampus: A histopathological and behavioral study. Iran. Red Crescent Med. J. 18, e30297. doi:10.5812/ircmj.30297

Saki, N., Jalalifar, M.A., Soleimani, M., Hajizamani, S., Rahim, F., 2013. Adverse effect of high glucose concentration on stem cell therapy. Int. J. Hematol.

UNIVERSITAS GADJAH MADA
PEMANFAATAN PRODUK SEKRETOM SEL PUNCA MESENKIMAL DARI TALI PUSAT MANUSIA YANG DIKULTUR DALAM KONDISI HIPOKSIA PADA MODEL DEGENERASI HIPPOCAMPUS
VERONIKA M. SIDHARTA, dr. Ginus Partadiredja, M.Sc., Ph.D.; dr. Rina Susilowati, Ph.D.; dr. E. Henny Herningtyas
Universitas Gadjah Mada, 2018 | Diunduh dari http://etd.repository.ugm.ac.id/

Stem Cell Res. 7, 33–39.

Salgado, A.J., Fraga, J.S., Mesquita, A.R., Neves, N.M., Reis, R.L., Sousa, N., 2010. Role of human umbilical cord mesenchymal progenitors conditioned media in neuronal/glial cell densities, viability, and proliferation. Stem Cells Dev. 19, 1067–1074. doi:10.1089/scd.2009.0279

Schäbitz, W.-R., Steigleder, T., Cooper-Kuhn, C.M., Schwab, S., Sommer, C., Schneider, A., Kuhn, H.G., 2007. Intravenous brain-derived neurotrophic factor enhances poststroke sensorimotor recovery and stimulates neurogenesis. Stroke 38, 2165–2172. doi:10.1161/STROKEAHA.106.477331

Semenza, G.L., 2004. Hydroxylation of HIF-1: Oxygen sensing at the molecular level. Physiology 19, 176–182. doi:10.1152/physiol.00001.2004

Semenza, G.L., Jiang, B.-H., Leung, S.W., Passantino, R., Concorde, J.-P., Maire, P., Giallongo, A., 1996. Hypoxia response elements in the aldolase A, enolase 1, and lactate dehydrogenase A gene promoters contain essential binding sites for hypoxia-inducible factor 1. J. Biol. Chem. 271, 32529–32537. doi:10.1074/jbc.271.51.32529

Silhol, M., Arancibia, S., Maurice, T., Tapia-Arancibia, L., 2007. Spatial memory training modifies the expression of brain-derived neurotrophic factor tyrosine kinase receptors in young and aged rats. Neuroscience 146, 962–973. doi:10.1016/j.neuroscience.2007.02.013

Silhol, M., Bonnichon, V., Rage, F., Tapia-Arancibia, L., 2005. Age-related changes in brain-derived neurotrophic factor and tyrosine kinase receptor isoforms in the hippocampus and hypothalamus in male rats. Neuroscience 132, 613–624. doi:10.1016/j.neuroscience.2005.01.008

Simon, H.-U., Haj-Yehia, A., Levi-Schaffer, F., 2000. Role of reactive oxygen species (ROS) in apoptosis induction. Apoptosis 5, 415–418. doi:10.1023/A:1009616228304

Spence, K.W., Lippitt, R., 1946. An experimental test of the sign-gestalt theory of trial and error learning. J. Exp. Psychol. 36, 491–502. doi:10.1037/h0062419

Sterio, D.C., 1984. The unbiased estimation of number and sizes of arbitrary particles using the disector. J. Microsc. 134, 127–136. doi:10.1111/j.1365-2818.1984.tb02501.x

Stolzing, A., Coleman, N., Scutt, A., 2006. Glucose-induced replicative senescence in mesenchymal stem cells. Rejuvenation Res. 9, 31–35. doi:10.1089/rej.2006.9.31

Teixeira, F.G., Carvalho, M.M., Neves-Carvalho, A., Panchalingam, K.M., Behie, L.A., Pinto, L., Sousa, N., Salgado, A.J., 2015. Secretome of mesenchymal progenitors from the umbilical cord acts as modulator of neural/glial proliferation and differentiation. Stem Cell Rev. Reports 11, 288–297. doi:10.1007/s12015-014-9576-2

Tsai, M.-J., Tsai, S.-K., Hu, B.-R., Liou, D.-Y., Huang, S.-L., Huang, M.-C., Huang, W.-C., Cheng, H., Huang, S.-S., 2014. Recovery of neurological function of ischemic stroke by application of conditioned medium of bone marrow mesenchymal stem cells derived from normal and cerebral ischemia rats. J. Biomed. Sci. 21, 5. doi:10.1186/1423-0127-21-5

UNIVERSITAS GADJAH MADA
PEMANFAATAN PRODUK SEKRETOM SEL PUNCA MESENKIMAL DARI TALI PUSAT MANUSIA YANG DIKULTUR DALAM KONDISI HIPOKSIA PADA MODEL DEGENERASI HIPPOCAMPUS
VERONIKA M. SIDHARTA, dr. Ginus Partadiredja, M.Sc., Ph.D.; dr. Rina Susilowati, Ph.D.; dr. E. Henny Herningtyas
Universitas Gadjah Mada, 2018 | Diunduh dari http://etd.repository.ugm.ac.id/

Uccelli, A., 2013. Mesenchymal stem cells exert a remarkable regenerative effect requiring minimal CNS integration. Commentary on: "Mesenchymal stem cells protect CNS neurons against glutamate excitotoxicity by inhibiting glutamate receptor expression and function" by Voulg. Exp. Neurol. 247, 292–295. doi:10.1016/j.expneurol.2013.01.028

Ullman, M.T., 2001. The declarative/procedural model of lexicon and grammar. J. Psycholinguist. Res. 30, 37–69. doi:http://dx.doi.org/10.1023/A:1005204207369

Vizoso, F.J., Eiro, N., Cid, S., Schneider, J., Perez-Fernandez, R., 2017. Mesenchymal stem cell secretome: Toward cell-free therapeutic strategies in regenerative medicine. Int. J. Mol. Sci. doi:10.3390/ijms18091852

Vorhees, C. V., Williams, M.T., 2014. Assessing spatial learning and memory in rodents. ILAR J. 55, 310–332. doi:10.1093/ilar/ilu013

Vorhees, C. V, Williams, M.T., 2006. Morris water maze: procedures for assessing spatial and related forms of learning and memory. Nat. Protoc. 1, 848–858. doi:10.1038/nprot.2006.116

Wang, G.L., Semenza, G.L., 1993. General involvement of hypoxia-inducible factor 1 in transcriptional response to hypoxia. Genetics 90, 4304–4308. doi:10.1073/pnas.90.9.4304

Wang, H.-S., Hung, S.-C., Peng, S.-T., Huang, C.-C., Wei, H.-M., Guo, Y.-J., Fu, Y.-S., Lai, M.-C., Chen, C.-C., 2004. Mesenchymal stem cells in the Wharton's jelly of the human umbilical cord. Stem Cells 22, 1330–1337. doi:10.1634/stemcells.2004-0013

Weickert, C.S., Hyde, T.M., Lipska, B.K., Herman, M.M., Weinberger, D.R., Kleinman, J.E., 2003. Reduced brain-derived neurotrophic factor in prefrontal cortex of patients with schizophrenia. Mol. Psychiatry 8, 592–610. doi:10.1038/sj.mp.4001308

West, M.J., 2012. Estimating volume in biological structures. Cold Spring Harb. Protoc. 7, 1129–1139. doi:10.1101/pdb.top071787

Wolf, A., Bauer, B., Abner, E.L., Ashkenazy-Frolinger, T., Hartz, A.M.S., 2016. A comprehensive behavioral test battery to assess learning and memory in 129S6/Tg2576 mice. PLoS One 11, 1–23. doi:10.1371/journal.pone.0147733

Wurmser, A.E., Palmer, T.D., Gage, F.H., Holland, P., 2004. Cellular interactions in the stem cell niche. Science (80-.). 304, 1253–1255. doi:10.1126/science.1099344

Xu, X., Yi, F., Pan, H., Duan, S., Ding, Z., Yuan, G., Qu, J., Zhang, H., Liu, G., 2013. Progress and prospects in stem cell therapy. Acta Pharmacol. Sin. 34, 741–746. doi:10.1038/aps.2013.77

Yamada, K., Nabeshima, T., 2003. Brain-derived neurotrophic factor/TrkB signaling in memory processes. J. Pharmacol. Sci. 91, 267–270. doi:10.1254/jphs.91.267

Yuliani, S., Mustofa, Partadiredja, G., 2017. Turmeric (Curcuma longa L.) extract may prevent the deterioration of spatial memory and the deficit of estimated total number of hippocampal pyramidal cells of trimethyltin-exposed rats. Drug Chem. Toxicol. 0545, 1–10. doi:10.1080/01480545.2017.1293087

Zhang, L., Tan, X., Dong, C., Zou, L., Zhao, H., Zhang, X., Tian, M., Jin, G.,

UNIVERSITAS
GADJAH MADA
PEMANFAATAN PRODUK SEKRETOM SEL PUNCA MESENKIMAL DARI TALI PUSAT MANUSIA YANG DIKULTUR DALAM KONDISI HIPOKSIA PADA MODEL DEGENERASI HIPPOCAMPUS
VERONIKA M. SIDHARTA, dr. Ginus Partadiredja, M.Sc., Ph.D.; dr. Rina Susilowati, Ph.D.; dr. E. Henny Herningtyas
Universitas Gadjah Mada, 2018 | Diunduh dari http://etd.repository.ugm.ac.id/

2012. In vitro differentiation of human umbilical cord mesenchymal stem cells (hUCMSCs), derived from Wharton's jelly, into choline acetyltransferase (ChAT)-positive cells. Int. J. Dev. Neurosci. 30, 471–477. doi:10.1016/j.ijdevneu.2012.05.006

Zhang, W., Su, X., Gao, Y., Sun, B., Yu, Y., Wang, X., Zhang, F., 2009. Berberine protects mesenchymal stem cells against hypoxia-induced apoptosis in vitro. Biol. Pharm. Bull. 32, 1335–1342. doi:JST.JSTAGE/bpb/32.1335 [pii]

Zhao, Q., Hu, J., Xiang, J., Gu, Y., Jin, P., Hua, F., Zhang, Z.Z., Liu, Y., Zan, K., Zhang, Z.Z., Zu, J., Yang, X., Shi, H., Zhu, J., Xu, Y., Cui, G., Ye, X., 2015. Intranasal administration of human umbilical cord mesenchymal stem cells-conditioned medium enhances vascular remodeling after stroke. Brain Res. 1624, 489–496. doi:10.1016/j.brainres.2015.08.003

Zheng, X., Liang, Y., He, Q., Yao, R., Bao, W., Bao, L., Wang, Y., Wang, Z., 2014. Current models of mammalian target of rapamycin complex 1 (mTORC1) activation by growth factors and amino acids. Int. J. Mol. Sci. 15, 20753–20769. doi:10.3390/ijms151120753

Zhu, S.F., Zhong, Z.N., Fu, X.F., Peng, D.X., Lu, G.H., Li, W.H., Xu, H.Y., Hu, H.B., He, J.M., Su, W.Y., He, Y.L., 2013. Comparison of cell proliferation, apoptosis, cellular morphology and ultrastructure between human umbilical cord and placenta-derived mesenchymal stem cells. Neurosci. Lett. 541, 77–82. doi:10.1016/j.neulet.2013.03.018

Zhu, W., Chen, J., Cong, X., Hu, S., Chen, X., 2006. Hypoxia and serum deprivation-induced apoptosis in mesenchymal stem cells. Stem Cells 24, 416–425. doi:10.1634/stemcells.2005-0121