

REFERENCES

- Anastassiades, M., Lehotay, S.J., Štajnbaher, D., Schenck, F.J., 2003. Fast and easy multiresidue method employing acetonitrile extraction/partitioning and “dispersive solid-phase extraction” for the determination of pesticide residues in produce. *J. AOAC Int.* 86, 412–431.
- Arslan, D., 2012. Physico-chemical characteristics of olive fruits of Turkish varieties from the province of Hatay. *Grasas y Aceites* 63, 158–166.
- Baccouri, O., Guerfel, M., Baccouri, B., Cerretani, L., Bendini, A., Lercker, G., Zarrouk, M., Miled, D.D. Ben, 2008. Chemical composition and oxidative stability of Tunisian monovarietal virgin olive oils with regard to fruit ripening. *Food Chem.* 109, 743–754.
- Ballus, C.A., Meinhart, A.D., de Souza Campos, F.A., da Silva, L.F. de O., de Oliveira, A.F., Godoy, H.T., 2014. A quantitative study on the phenolic compound, tocopherol and fatty acid contents of monovarietal virgin olive oils produced in the southeast region of Brazil. *Food Res. Int.* 62, 74–83.
- Ben Brahim, S., Kelebek, H., Ammar, S., Abichou, M., Bouaziz, M., 2017. LC–MS phenolic profiling combined with multivariate analysis as an approach for the characterization of extra virgin olive oils of four rare Tunisian cultivars during ripening. *Food Chem.* 229, 9–19.
- Borges, T.H., López, L.C., Pereira, J.A., Cabrera, C., Vique, –, Seiquer, I., 2017. Comparative analysis of minor bioactive constituents (CoQ 10, tocopherols and phenolic compounds) in Arbequina extra virgin olive oils from Brazil and Spain. *J. Food Compos. Anal.* 63, 47–54.
- Bouaziz, M., Chamkha, M., Sayadi, S., 2004. Comparative study on phenolic content and antioxidant activity during maturation of the olive cultivar Chemlali from Tunisia. *J. Agric. Food Chem.* 52, 5476–5481.
- Brahmi, F., Mechri, B., Dhibi, M., Hammami, M., 2015. Antioxidant capacity and phenolic content of olive fruits from Tunisia, extracted with different solvents. *J. Plant Sci.* 2015; 3, 8–12.
- Brenes, M., Hidalgo, F.J., García, A., Rios, J.J., García, P., Zamora, R., Garrido, A., 2000. Pinoresinol and 1-acetoxypinoresinol, two new phenolic compounds identified in olive oil. *J. Am. Oil Chem. Soc.* 77, 715–720.
- Briante, R., Cara, F. La, Tonziello, M.P., Febbraio, F., Nucci, R., 2001. Antioxidant activity of the main bioactive derivatives from oleuropein hydrolysis by hyperthermophilic β -glycosidase. *J. Agric. Food Chem.* 49, 3198–3203.
- Briante, R., Patumi, M., Limongelli, S., Febbraio, F., Vaccaro, C., Di Salle, A., La Cara, F., Nucci, R., 2002. Changes in phenolic and enzymatic activities content during fruit ripening in two Italian cultivars of *Olea europaea* L. *Plant*

Sci. 162, 791–798.

- Capogna, D., Gomez, M.I., 2016. Olive oil: an overview of the Japanese market. *Oilseeds fats Crop. Lipids* 23, 21–23.
- Caporaso, N., Savarese, M., Paduano, A., Guidone, G., De Marco, E., Sacchi, R., 2015. Nutritional quality assessment of extra virgin olive oil from the Italian retail market: Do natural antioxidants satisfy EFSA health claims? *J. Food Compos. Anal.* 40, 154–162.
- Capriotti, A.L., Cavaliere, C., Crescenzi, C., Foglia, P., Nescatelli, R., Samperi, R., Laganà, A., 2014. Comparison of extraction methods for the identification and quantification of polyphenols in virgin olive oil by ultra-HPLC-QToF mass spectrometry. *Food Chem.* 158, 392–400.
- Carrasco-Pancorbo, A., Cerretani, L., Bendini, A., Segura-Carretero, A., Del Carlo, M., Gallina-Toschi, T., Lercker, G., Compagnone, D., Fernández-Gutiérrez, A., 2005. Evaluation of the antioxidant capacity of individual phenolic compounds in virgin olive oil. *J. Agric. Food Chem.* 53, 8918–8925.
- Cheng, Z., Zhan, M., Yang, Z., Zumstein, K., Chen, H., Huang, Q., 2017. The major qualitative characteristics of olive (*Olea europaea* L.) cultivated in Southwest China. *Front. Plant Sci.* 8, 1–8.
- Covas, M.I., Ruiz-Gutierrez, V., de la Torre, R., Kafatos, A., Lamuela-Raventos, R.M., Osada, J., Owen, R.W., Visioli, F., 2006. Monir Components of Olive Oil: Evidrnce to Date of Health Benefits in Humans. *Nutr. Rev.* 64, S20–S31.
- De Boer, A., Vos, E., Bast, A., 2014. Implementation of the nutrition and health claim regulation - The case of antioxidants. *Regul. Toxicol. Pharmacol.* 68, 475–487.
- De La Torre-Carbot, K., Jauregui, O., Gimeno, E., Castellote, A.I., Lamuela-Raventós, R.M., López-Sabater, M.C., 2005. Characterization and quantification of phenolic compounds in olive oils by solid-phase extraction, HPLC-DAD, and HPLC-MS/MS. *J. Agric. Food Chem.* 53, 4331–4340.
- Dekdouk, N., Malafrente, N., Russo, D., Faraone, I., De Tommasi, N., Ameddah, S., Severino, L., Milella, L., 2015. Phenolic Compounds from *Olea europaea* L. Possess Antioxidant Activity and Inhibit Carbohydrate Metabolizing Enzymes in Vitro. *Evidence-based Complement. Altern. Med.* 2015.
- Demopoulos, V., Karkoula, E., Magiatis, P., Melliou, E., Kotsiras, A., Mouroutoglou, C., 2015. Correlation of oleocanthal and oleacein concentration with pungency and bitterness in “koroneiki” virgin olive oil. *Acta Hortic.* 1099, 219–224.
- Deng, J., Xu, Z., Xiang, C., Liu, J., Zhou, L., Li, T., Yang, Z., Ding, C., 2017. Comparative evaluation of maceration and ultrasonic-assisted extraction of phenolic compounds from fresh olives. *Ultrason. Sonochem.* 37, 328–334.

- Dierkes, G., Krieger, S., Du, R., Bongartz, A., Schmitz, O.J., Hayen, H., 2012. High-Performance Liquid Chromatography – Mass Spectrometry Profiling of Phenolic Compounds for Evaluation of Olive Oil Bitterness and Pungency. *J. Agric. Food Chem.* 60, 7597–7606.
- Dolan, J., 2004. A Guide to HPLC - ACE Guide 16.
- El Riachy, M., Priego-Capote, F., León, L., Rallo, L., Luque de Castro, M.D., 2011. Hydrophilic antioxidants of virgin olive oil. Part 2: Biosynthesis and biotransformation of phenolic compounds in virgin olive oil as affected by agronomic and processing factors. *Eur. J. Lipid Sci. Technol.* 113, 692–707.
- Fang, X., Wang, J., Hao, J., Li, X., Guo, N., 2015. Simultaneous extraction, identification and quantification of phenolic compounds in *Eclipta prostrata* using microwave-assisted extraction combined with HPLC-DAD-ESI-MS/MS. *Food Chem.* 188, 527–536.
- Fiorini, D., Boarelli, M.C., Conti, P., Alfei, B., Caprioli, G., Ricciutelli, M., Sagratini, G., Fedeli, D., Gabbianelli, R., Pacetti, D., 2018. Chemical and sensory differences between high price and low price extra virgin olive oils. *Food Res. Int.* 105, 65–75.
- Fitri, A., Andriani, M., Sudarman, A., Toharmat, T., Yonekura, L., Tamura, H., Ramli, N., 2016. Screening of antioxidant activities and their bioavailability of tropical fruit by products from Indonesia. *Int. J. Pharm. Pharm. Sci.* 8, 5–9.
- Franco, M.N., Galeano-Díaz, T., López, Ó., Fernández-Bolaños, J.G., Sánchez, J., De Miguel, C., Gil, M.V., Martín-Vertedor, D., 2014. Phenolic compounds and antioxidant capacity of virgin olive oil. *Food Chem.* 163, 289–298.
- García-Villalba, R., Carrasco-Pancorbo, A., Oliveras-Ferraro, C., Vázquez-Martín, A., Menéndez, J.A., Segura-Carretero, A., Fernández-Gutiérrez, A., 2010a. Characterization and quantification of phenolic compounds of extra-virgin olive oils with anticancer properties by a rapid and resolute LC-ESI-TOF MS method. *J. Pharm. Biomed. Anal.* 51, 416–429.
- García-Villalba, R., Carrasco-Pancorbo, A., Zurek, G., Behrens, M., Bäßmann, C., Segura-Carretero, A., Fernández-Gutiérrez, A., 2010b. Nano and rapid resolution liquid chromatography-electrospray ionization-time of flight mass spectrometry to identify and quantify phenolic compounds in olive oil. *J. Sep. Sci.* 33, 2069–2078.
- García-Villalba, R., Pacchiarotta, T., Carrasco-Pancorbo, A., Segura-Carretero, A., Fernández-Gutiérrez, A., Deelder, A.M., Mayboroda, O.A., 2011. Gas chromatography-atmospheric pressure chemical ionization-time of flight mass spectrometry for profiling of phenolic compounds in extra virgin olive oil. *J. Chromatogr. A* 1218, 959–971.
- Gilbert-López, B., Valencia-Reyes, Z.L., Yufra-Picardo, V.M., García-Reyes, J.F., Ramos-Martos, N., Molina-Díaz, A., 2014. Determination of Polyphenols in

- Commercial Extra Virgin Olive Oils from Different Origins (Mediterranean and South American Countries) by Liquid Chromatography–Electrospray Time-of-Flight Mass Spectrometry. *Food Anal. Methods* 7, 1824–1833.
- Goulas, V., Charisiadis, P., P. Gerothanassis, I., A. Manganaris, G., 2012. Classification, Biotransformation and Antioxidant Activity of Olive Fruit Biophenols: A Review. *Curr. Bioact. Compd.* 8, 232–239.
- Gupta, D.K., Verma, M.K., Anand, R., Khajuria, R.K., 2013. Development of a validated UPLC-qTOF-MS/MS method for determination of bioactive constituent from *Glycyrrhiza glabra*. *J. Pharm. Anal.* 3, 205–210.
- Gutierrez-Rosales, F., Romero, M.P., Casanovas, M., Motilva, M.J., Mínguez-Mosquera, M.I., 2010. Metabolites Involved in Oleuropein Accumulation and Degradation in Fruits of *Olea europaea* L.: Hojiblanca and Arbequina Varieties. *J. Agric. Food Chem.* 58, 12924–12933.
- Gutierrez-Rosales, F., Romero, M.P., Casanovas, M., Motilva, M.J., Mínguez-Mosquera, M.I., 2012. β -Glucosidase involvement in the formation and transformation of oleuropein during the growth and development of olive fruits (*Olea europaea* L. cv. Arbequina) grown under different farming practices. *J. Agric. Food Chem.* 60, 4348–4358.
- Guzmán, E., Baeten, V., Pierna, J.A.F., García-Mesa, J.A., 2015. Determination of the olive maturity index of intact fruits using image analysis. *J. Food Sci. Technol.* 52, 1462–1470.
- Hrcirik, K., Fritsche, S., 2004. Comparability and reliability of different techniques for the determination of phenolic compounds in virgin olive oil. *Eur. J. Lipid Sci. Technol.* 106, 540–549.
- Jemai, H., Bouaziz, M., Sayadi, S., 2009. Phenolic composition, sugar contents and antioxidant activity of tunisian sweet olive cultivar with regard to fruit ripening. *J. Agric. Food Chem.* 57, 2961–2968.
- Jerman Klen, T., Golc Wondra, A., Vrhovšek, U., Sivilotti, P., Vodopivec, B.M., 2015. Olive fruit phenols transfer, transformation, and partition trail during laboratory-scale olive oil processing. *J. Agric. Food Chem.* 63, 4570–4579.
- Kedare, S.B., Singh, R.P., 2011. Genesis and development of DPPH method of antioxidant assay. *J. Food Sci. Technol.* 48, 412–422.
- Kim, J., Kim, J. Il, Lee, C.W., 2016. Development and validation of a modified QuEChERS method coupled with LC-MS / MS to determine arbutin in pear peels. *Food Sci. Biotechnol.* 25, 987–992.
- Lafka, T.-I., Lazou, A., Sinanoglou, V., Lazos, E., 2013. Phenolic Extracts from Wild Olive Leaves and Their Potential as Edible Oils Antioxidants. *Foods* 2, 18–31.
- Morelló, J.R., Romero, M.P., Motilva, M.J., 2004. Effect of the maturation of the

- olive fruit on the phenolic fraction of drupes and oils from Arbequina, Farga, and Morrut cultivars. *J. Agric. Food Chem.* 52, 6002–6009.
- Morelló, J.R., Vuorela, S., Romero, M.P., Motilva, M.J., Heinonen, M., 2005. Antioxidant activity of olive pulp and olive oil phenolic compounds of the arbequina cultivar. *J. Agric. Food Chem.* 53, 2002–2008.
- Mtimet, N., Kashiwagi, A.K., Zaibet, L., Masakazu, N., 2008. Exploring Japanese olive oil consumer behavior 26–29.
- Pizarro, M.L., Becerra, M., Sayago, A., Beltrán, M., Beltrán, R., 2013. Comparison of Different Extraction Methods to Determine Phenolic Compounds in Virgin Olive Oil. *Food Anal. Methods* 6, 123–132.
- Reboredo-Rodriguez, P., Valli, E., Bendini, A., Di Lecce, G., Simal-Gandara, J., Gallina Toschi, T., 2016. A widely used spectrophotometric assay to quantify olive oil biophenols according to the health claim (EU Reg. 432/2012). *Eur. J. Lipid Sci. Technol.* 118, 1593–1599.
- Rejczak, T., Tuzimski, T., 2015. A review of recent developments and trends in the QuEChERS sample preparation approach. *Open Chem.* 13, 980–1010.
- Ricciutelli, M., Marconi, S., Boarelli, M.C., Caprioli, G., Sagratini, G., Ballini, R., Fiorini, D., 2017. Olive oil polyphenols: A quantitative method by high-performance liquid-chromatography-diode-array detection for their determination and the assessment of the related health claim. *J. Chromatogr. A* 1481, 53–63.
- Ríos, J.J., Gil, M.J., Gutiérrez-Rosales, F., 2005. Solid-phase extraction gas chromatography-ion trap-mass spectrometry qualitative method for evaluation of phenolic compounds in virgin olive oil and structural confirmation of oleuropein and ligstroside aglycons and their oxidation products. *J. Chromatogr. A* 1093, 167–176.
- Romero, C., Brenes, M., Garcia, P., Garcia, A., Garrido, A., 2004. Polyphenol Changes during Fermentation of Naturally Black Olives. *J. Agric. Food Chem.* 52, 1973–1979.
- Rotondi, A., Bendini, A., Cerretani, L., Mari, M., Lercker, G., Toschi, T.G., 2004. Effect of olive ripening degree on the oxidative stability and organoleptic properties of cv. Nostrana di Brisighella extra virgin olive oil. *J. Agric. Food Chem.* 52, 3649–3654.
- Ryan, D., Prenzler, P.D., Lavee, S., Antolovich, M., Robards, K., 2003. Quantitative Changes in Phenolic Content during Physiological Development of the Olive (*Olea europaea*) Cultivar Hardy's Mammoth. *J. Agric. Food Chem.* 51, 2532–2540.
- Sato, A., Shinozaki, N., Tamura, H., 2014. Secoiridoid type of antiallergic substances in olive waste materials of three japanese varieties of *olea europaea*. *J. Agric. Food Chem.* 62, 7787–7795.

- Sato, A., Zhang, T., Yonekura, L., Tamura, H., 2015. Antiallergic activities of eleven onions (*Allium cepa*) were attributed to quercetin 4'-glucoside using QuEChERS method and Pearson's correlation coefficient. *J. Funct. Foods* 14, 581–589.
- Shibasaki, H., 2005. Influence of Fruit Ripening on Chemical Properties of "Mission" Variety Olive Oil in Japan. *Food Sci. Technol. Res.* 11, 9–12.
- Talhaoui, N., Gómez-Caravaca, A.M., León, L., De La Rosa, R., Fernández-Gutiérrez, A., Segura-Carretero, A., 2015. Pattern of Variation of Fruit Traits and Phenol Content in Olive Fruits from Six Different Cultivars. *J. Agric. Food Chem.* 63, 10466–10476.
- Talhaoui, N., Gómez-Caravaca, A.M., León, L., De La Rosa, R., Fernández-Gutiérrez, A., Segura-Carretero, A., 2016. From olive fruits to olive Oil: Phenolic compound transfer in six different olive cultivars grown under the same agronomical conditions. *Int. J. Mol. Sci.* 17, 1–14.
- Talhaoui, N., Gómez-Caravaca, A.M., León, L., De la Rosa, R., Segura-Carretero, A., Fernández-Gutiérrez, A., 2014. Determination of phenolic compounds of "Sikitita" olive leaves by HPLC-DAD-TOF-MS. Comparison with its parents "Arbequina" and "Picual" olive leaves. *LWT - Food Sci. Technol.* 58, 28–34.
- Tasioula-Margari, M., Tsabolatidou, E., 2015. Extraction, Separation, and Identification of Phenolic Compounds in Virgin Olive Oil by HPLC-DAD and HPLC-MS. *Antioxidants* 4, 548–562.
- Tripoli, E., Giammanco, M., Tabacchi, G., Majo, D. Di, 2017. The phenolic compounds of olive oil : structure , biological activity and beneficial effects on human health 98–112.
- Tuberoso, C.I.G., Jerković, I., Maldini, M., Serreli, G., 2016. Phenolic Compounds, Antioxidant Activity, and Other Characteristics of Extra Virgin Olive Oils from Italian Autochthonous Varieties Tonda di Villacidro, Tonda di Cagliari, Semidana, and Bosana. *J. Chem.* 2016.
- Valente, I.M., Santos, C.M., Moreira, M.M., Rodrigues, J.A., 2013. New application of the QuEChERS methodology for the determination of volatile phenols in beverages by liquid chromatography. *J. Chromatogr. A* 1271, 27–32.
- Vinha, A.F., Ferreres, F., Silva, B.M., Valentão, P., Gonçalves, A., Pereira, J.A., Oliveira, M.B., Seabra, R.M., Andrade, P.B., 2005. Phenolic profiles of Portuguese olive fruits (*Olea europaea* L.): Influences of cultivar and geographical origin. *Food Chem.* 89, 561–568.
- Vossen, P., 2007. Olive Oil: History, Production, and Characteristics of the World's Classic Oils. *Hortic. Sci.* 42, 1093–1100.
- Yanık, D.K., 2017. Alternative to traditional olive pomace oil extraction systems:

Microwave-assisted solvent extraction of oil from wet olive pomace. *LWT - Food Sci. Technol.* 77, 45–51.

Yousfi, K., Cert, R.M., Garc, J.M., 2006. Changes in quality and phenolic compounds of virgin olive oils during objectively described fruit maturation. *Eur. Food Res. Technol.* 223, 117–124.

Zaringhalami, S., Ebrahimi, M., Piravi Vanak, Z., Ganjloo, A., 2015. Effects of cultivar and ripening stage of Iranian olive fruit on bioactive compounds and antioxidant activity of its virgin oil. *Int. Food Res. J.* 22, 1961–1967.