

DAFTAR PUSTAKA

- Abeyrathne, E.D.N.S., Lee, H.Y., and Ahn, D.U., 2013, Egg white proteins and their potential use in food processing or as nutraceutical and pharmaceutical agents – a review., *Poult. Sci.*, 92, 3292–9.
- Abtahi, S. and Aminlari, M., 1997, Effect of Sodium Sulfite, Sodium Bisulfite, Cysteine, and pH on Protein Solubility and Sodium Dodecyl Sulfate-Polyacrylamide Gel Electrophoresis of Soybean Milk Base, *J. Agric. Food Chem.*, 45, 4768–4772.
- Bailey, J.L. and Cole, R.D., 1959, Studies on the Reaction of Sulfite with Proteins, *J. Biol. Chem.*, 234, 1733–1739.
- Beauchamp, D.L. and Khajehpour, M., 2012, Studying salt effects on protein stability using ribonuclease t1 as a model system, *Biophys. Chem.*, 161, 29–38.
- Berkelman, T. and Stenstedt, T.T., 1998, 2-D Electrophoresis using immobilized pH gradients, AB. Amersham Biosciences, Munich.
- Bernhisel-Broadbent, J., Dintzis, H.M., Dintzis, R.Z., and Sampson, H.A., 1994, Allergenicity and antigenicity of chicken egg ovomucoid (Gal d III) compared with ovalbumin (Gal d I) in children with egg allergy and in mice, *J. Allergy Clin. Immunol.*, 93, 1047–1059.
- Carpentier, S.C., Panis, B., Vertommen, A., Swennen, R., Sergeant, K., Renaut, J., et al., 2008, PROTEOME ANALYSIS OF NON-MODEL PLANTS: A CHALLENGING BUT POWERFUL APPROACH, *Mass Spectrom. Rev.*, 27, 354–377.
- Chou, D.K., Krishnamurthy, R., Randolph, T.W., Carpenter, J.F., and Manning, M.C., 2005, Effects of Tween 20® and Tween 80® on the stability of Albutropin during agitation, *J. Pharm. Sci.*, 94, 1368–1381.
- Desert, C., Guérin-Dubiard, C., Nau, F., Jan, G., Val, F., and Mallard, J., 2001, Comparison of different electrophoretic separations of hen egg white proteins, *J. Agric. Food Chem.*, 49, 4553–4561.
- Fæste, C.K., Løvberg, K.E., Lindvik, H., and Egaas, E., 2007, Extractability, stability, and allergenicity of egg white proteins in differently heat-processed foods, *J. AOAC Int.*, 90, 427–436.
- Fernández, S.S., Padilla, A.P., and Mucciarelli, S., 1999, Protein extraction from Atriplex lampa leaves: Potential use as forage for animals used for human diets, *Plant Foods Hum. Nutr.*, 54, 251–259.
- Forsythe, E.L. and Pusey, M.L., 1996, The Effects of Acetate Buffer

Concentration on Lysozyme Solubility, *J. Cryst. Growth*, 168, 112–117.

Goklen, K.E. and Hatton, T.A., 1987, Liquid-Liquid Extraction of Low Molecular-Weight Proteins by Selective Solubilization in Reversed Micelles, *Sep. Sci. Technol.*, 22, 831–841.

Goklen, K.E. and Hatton, T.A., 1985, Protein extraction using reverse micelles, *Biotechnol. Prog.*, 1, 69–74.

Görg, A., Weiss, W., and Dunn, M.J., 2004, Current two-dimensional electrophoresis technology for proteomics, *Proteomics*, 4, 3665–3685.

Guérin-Dubiard, C., Pasco, M., Hietanen, A., Quiros Del Bosque, A., Nau, F., and Croguennec, T., 2005, Hen egg white fractionation by ion-exchange chromatography, *J. Chromatogr. A*, 1090, 58–67.

He, Z., Tan, J.S., Lai, O.M., and Ariff, A.B., 2015, Optimization of conditions for the single step IMAC purification of miraculin from *Synsepalum dulcificum*, *Food Chem.*, 181, 19–24.

Herbert, B. and Ryde, N., 1999, Review Advances in protein solubilisation for two-dimensional electrophoresis Proteomics and 2-DE, *Electrophoresis*, 20, 660–663.

Hildebrandt, S., Steinhart, H., and Paschke, A., 2008, Comparison of different extraction solutions for the analysis of allergens in hen's egg, *Food Chem.*, 108, 1088–1093.

Hirose, J., Kitabatake, N., Kimura, A., and Narita, H., 2004, Recognition of native and/or thermally induced denatured forms of the major food allergen, ovomucoid, by human IgE and mouse monoclonal IgG antibodies, *Biosci Biotechnol Biochem*, 68, 2490–2497.

Hoffmann, C., Blume, A., Miller, I., and Garidel, P., 2009, Insights into protein-polysorbate interactions analysed by means of isothermal titration and differential scanning calorimetry, *Eur. Biophys. J.*, 38, 557–568.

Huang, W. and Sun, X., 2000, Adhesive properties of soy proteins modified by sodium dodecyl sulfate and sodium dodecylbenzene sulfonate, *JAACS, J. Am. Oil Chem. Soc.*, 77, 705–708.

James, D.H., Castor, W.M., Khan, A.Z., Ruckenstein, E., Chen, S.S., Rowe, S.M., et al., 2003, Surfactants, *Ullmann's Encycl. Ind. Chem.*, 1, 1–32.

Jarudilokkul, S., Poppenborg, L.H., and Stuckey, D.C., 1999, Backward extraction of reverse micellar encapsulated proteins using a counterionic surfactant, *Biotechnol. Bioeng.*, 62, 593–601.

- Jelinska, A., Zagozdzon, A., Recki, M.G., Wisniewska, A., Frelek, J., and Holyst, R., 2017, Denaturation of proteins by surfactants studied by the taylor dispersion analysis, *PLoS One*, 12, 1–11.
- Jensen, J., 1999, Fate and effects of linear alkylbenzene sulphonates (LAS) in the terrestrial environment, *Sci. Total Environ.*, 226, 93–111.
- Johnson, D.H., Wilson, W.W., and DeLucas, L.J., 2014, Protein solubilization: A novel approach, *J. Chromatogr. B Anal. Technol. Biomed. Life Sci.*, 971, 99–106.
- Jolival, C., Minier, M., and Renon, H., 1993, Extraction of Cytochrome c in Sodium Dodecylbenzenesulfonate Microemulsions, *Biotechnol. Prog.*, 9, 456–461.
- Lad, M.D., Ledger, V.M., Briggs, B., Green, R.J., and Frazier, R.A., 2003, Analysis of the SDS - Lysozyme Binding Isotherm, *Langmuir*, 19, 5098–5103.
- Langeland, T., 1983, A clinical and immunological study of allergy to hen's egg white. I. A clinical study of egg allergy., *Clin. Allergy*, 13, 371–382.
- Liljas, L., Lundahl, P., and Hjerten, S., 1974, Human erythrocyte membranes Water-extractable proteins Solubilization of the membrane residue with Tween 20 Analytical polyacrylamide-gel electrophoresis, *Biochim. Biophys. Acta*, 352, 327–337.
- Lim, S., Lee, J.-H., Shin, D.-H., and Lim, H.S., 1999, Comparison of Protein Extraction Solutions for Rice Starch Isolation and Effects of Residual Protein Content on Starch Pasting Properties, *Starch - Stärke*, 51, 120–125.
- Lundgren, H.P., 1945, Synthetic Protein Fibers from Protein-Detergent Complexes, *Text. Res. J.*, 15, 335–353.
- Mackay, D., Di Guardo, A., Paterson, S., Kicsi, G., Cowan, C.E., and Kane, D.M., 1996, Assessment of chemical fate in the environment using evaluative, regional and local-scale models: Illustrative application to chlorobenzene and linear alkylbenzene sulfonates, *Environ. Toxicol. Chem.*, 15, 1638–1648.
- Minkiewicz, P. and Dziuba, J., 2008, Food peptidomics, *Food Technol. Biotechnol.*, 46, 1–10.
- Negrusz, A. and Cooper, G.A.A., 2013, Clarke's Analytical Forensic Toxicology, 2nd ed. Pharmaceutical Press, London.
- Pastorello, E.A. and Trambaioli, C., 2001, Isolation of food allergens, *J. Chromatogr. B Biomed. Sci. Appl.*, 756, 71–84.

- Pedreschi, R., Maarten, H., Lilley, K.S., and Bart, N., 2010, Proteomics for the food industry: Opportunities and challenges, *Crit. Rev. Food Sci. Nutr.*, 50, 680–692.
- Pelegrine, D.H.G. and Gasparetto, C.A., 2005, Whey proteins solubility as function of temperature and pH, *LWT - Food Sci. Technol.*, 38, 77–80.
- Piergiovanni, A.R., 2007, Extraction and separation of water-soluble proteins from different wheat species by acidic capillary electrophoresis. In, *Journal of Agricultural and Food Chemistry.*, pp. 3850–3856.
- Raikos, V., Hansen, R., Campbell, L., and Euston, S.R., 2006, Separation and identification of hen egg protein isoforms using SDS-PAGE and 2D gel electrophoresis with MALDI-TOF mass spectrometry, *Food Chem.*, 99, 702–710.
- Rossky, P.J., 2008, Protein denaturation by urea: Slash and bond, *Proc. Natl. Acad. Sci.*, 105, 16825–16826.
- Sampson, H.A. and Cooke, S.K., 1990, Food allergy and the potential allergenicity-antigenicity of microparticulated egg and cow's milk proteins, *J. Am. Coll. Nutr.*, 9, 410–417.
- Schmid, M., Prinz, T.K., Stäbler, A., and Sänglerlaub, S., 2017, Effect of Sodium Sulfite, Sodium Dodecyl Sulfate, and Urea on the Molecular Interactions and Properties of Whey Protein Isolate-Based Films, *Front. Chem.*, 4, 1–15.
- Tan, S.H., Mailer, R.J., Blanchard, C.L., and Agboola, S.O., 2011, Canola Proteins for Human Consumption: Extraction, Profile, and Functional Properties, *J. Food Sci.*, 76, 16–18.
- Toldrá, F. and Nollet, L.M.L., 2013, Proteomics in foods: Principles and applications, Springer, United States.
- Ugwu, S.O. and Apte, S.P., 2004, The Effect of Buffers on Protein Conformational Stability, *Pharm. Technol.*, 28, 86–109.
- Umesh Hebbar, H., Sumana, B., and Raghavarao, K.S.M.S., 2008, Use of reverse micellar systems for the extraction and purification of bromelain from pineapple wastes, *Bioresour. Technol.*, 99, 4896–4902.
- Ünsel, M., Sin, A.Z., Ardeniz, Ö., Erdem, N., Ersoy, R., Gulbahar, O., et al., 2007, New onset egg allergy in an adult, *J. Investig. Allergol. Clin. Immunol.*, 17, 55–58.
- Wang, J., MacNeil, J.D., and Kay, J.F., 2012, Chemical Analysis of Antibiotics, 2nd ed. John Wiley & Sons, Inc., Hoboken.

Witteaman, A.M., Akkerdaas, J.H., van Leeuwen, J., van der Zee, J.S., and Aalberse, R.C., 1994, Identification of a Cross-Reactive Allergen (Presumably Tropomyosin) in Shrimp, Mite and Insects, *Int. Arch. Allergy Immunol.*, 105, 56–61.

Zou, Q., Habermann-Rottinghaus, S.M., and Murphy, K.P., 1998, Urea effects on protein stability: Hydrogen bonding and the hydrophobic effect, *Proteins Struct. Funct. Genet.*, 31, 107–115.